

Editorial

Mechanisms Underlying Chemoprevention of Ovarian Cancer

Gordon B. Mills¹

Ovarian cancer remains the most lethal of the gynecological cancers. Although relatively uncommon, afflicting ~1 of 70 women in the United States, the high mortality rate makes this disease a major health concern. The high mortality rate arises from the lack of an effective screening approach combined with inadequate therapeutic approaches for advanced disease. Indeed, fewer than 25% of ovarian cancers are identified at an early curable stage. Other than family and personal history of cancer or the inheritance of abnormalities in the BRCA1 or BRCA2 or other ovarian cancer predisposition genes, there are few indicators that a woman has an elevated risk of development of cancer. Thus, it is difficult to identify women warranting prophylactic surgery or intensive screening. The high mortality rate combined with the lack of effective screening approaches make ovarian cancer a strong candidate for chemoprevention (1).

In this issue of "Clinical Cancer Research," Rodriguez-Burford *et al.* (2) explore the potential mechanisms underlying epidemiological observations that ovarian cancer occurs at a lower frequency in women exposed to NSAIDs.² Cramer *et al.* (3) in a population-based, case-control study demonstrated that a 6-month exposure to acetaminophen at least 1 day/week, but not aspirin or ibuprofen, resulted in a statistically significant 2-fold decrease in the lifetime risk of developing ovarian cancer. Moysich *et al.* (4) in a hospital-based, case-control study also demonstrated a time- and dose-dependent association of acetaminophen, but not aspirin use, with decreased risk of developing ovarian cancer. Rosenberg *et al.* (5) in a large-scale, case-control study failed to confirm the association of moderate use of acetaminophen and decreased risk of ovarian cancer demonstrated by Cramer *et al.* (3) but did identify a statistically significant decrease in development of ovarian cancer with prolonged acetaminophen use, at least ≥ 4 days/week for 5 years. Tavani *et al.* (6) has failed recently to detect an association of aspirin use with prevention of ovarian cancer. In contrast to these reports, Akhmedkhanov *et al.* (7) demonstrated a 2~3-fold decrease in epithelial ovarian cancer associated with aspirin use. Taken together, although controversial, these epidemiological observations suggest that an improved understanding of the mechanism(s) by which NSAIDs may decrease the development of ovarian cancer could lead to improved approaches for chemoprevention of this deadly disease.

The studies in this issue of "Clinical Cancer Research" identify several of the problems associated with attempting to use cell models to understand the mechanisms underlying epidemiological studies of chemoprevention of epithelial ovarian cancer. At this point, there is no readily manipulatable mammalian model of epithelial ovarian cancer. Indeed, the only extant animal model with reproducible epithelial ovarian cancer is a chicken model (8). This model, while contributing to our understanding of chemoprevention of ovarian cancer, is far from optimal. The mouse models of human cancer consortium, as well as members of the ovarian cancer research community, are investigating a number of approaches to develop practical animal models; however, at the present, it is necessary to rely on ovarian cancer cell lines, cultures of normal ovarian epithelium, or normal epithelium with enforced expression of genes, such as the large T antigen of SV40, telomerase, and oncogenes to attempt to explore the mechanisms underlying chemoprevention of epithelial ovarian cancer. These established cell culture systems and, in particular, ovarian cancer cell lines may not adequately reflect the transformation processes targeted by chemoprevention agents *in vivo*.

The studies by Rodriguez-Burford *et al.* (2) demonstrate that acetylsalicylic acid, acetaminophen, and a Cox-2 inhibitor, NS-398, can decrease the growth of fully transformed human epithelial ovarian cancer cells. The Cox-2 agent both decreased cell proliferation in established cell lines and induced apoptosis in freshly isolated ovarian cancer cells. This information implicates cellular proliferation and survival in the action of NSAIDs on the development of epithelial ovarian cancer. Although all three NSAIDs decreased the proliferation of ovarian cancer cell lines, concentrations far above the therapeutic range and well above the maximal tolerated dose and, in particular, the chemopreventative doses of the inhibitors were required to demonstrate antiproliferative effects *in vitro* (2). This is compatible with the observation that in contrast to bowel cancer cells, which are very sensitive to the effects of NSAIDs likely as a consequence of overexpression of Cox2 (9–12) and dependent on Cox2 function for growth and survival, human epithelial ovarian cancer cells appear to express Cox1 and Cox2 at low levels (2, 13). Thus, either a prolonged low-dose exposure to the NSAIDs has effects on Cox2 function in normal or partially transformed ovarian epithelium, or NSAIDs exert their effects through alternative mechanisms.

The observation that very high concentrations of NSAIDs are required to decrease the proliferation of ovarian cancer cell lines and that ovarian cancer cells express low levels of COX1 and COX2 (2, 13) suggests that the effects of NSAIDs on ovarian cancer prevention may be attributable to mechanisms independent of inhibition of COX1 and COX2 as reflected by the effects of NSAIDs on the proliferation or survival of established ovarian cancer cell lines. In this aspect, it is important to note that the epidemiological data most strongly implicate acetaminophen as compared with aspirin or ibuprofen in the prevention of ovarian cancer (2–7). Acetaminophen is a relatively poor inhibitor of cyclooxygenase and does not have

Received 9/28/01; accepted 10/3/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ To whom requests for reprints should be addressed, at Molecular Therapeutics, University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030.

² The abbreviations used are: NSAID, nonsteroidal anti-inflammatory agent; PI3K, phosphatidylinositol 3'-kinase; LPA, lysophosphatidic acid.




Fig. 1 Potential mechanism of Cox2 expression and NSAID action in the prevention of ovarian cancer. AKT has been demonstrated to be a potent regulator of Cox2 expression and action. Multiple mechanisms result in constitutive activation in ovarian cancer cells. Ovarian cancer cells can overexpress members of the human epidermal growth factor receptor kinase family (*HER*). The *HER* family efficiently couples to both the *PI3K* and *RAS* pathways, as well as to the *SRC* proto-oncogene. In addition, the LPA receptors 2 and 3 of the G protein-coupled receptor family are expressed at elevated levels in ovarian cancer cells. In ovarian cancer cells, these receptors couple efficiently to *PI3K*, *RAS*, and *SRC*. In the presence of LPA, which is itself increased in ovarian cancer, the LPA Multiple components of the *PI3K* pathway are amplified in ovarian cancer as a consequence of increased DNA copy number, including the *PI3K* catalytic subunit and *AKT1* and *2*. In addition, *PTEN*, which functions as a tumor suppressor gene to inhibit activation of *AKT* by *PI3K*, is expressed at decreased levels and can be mutated, particularly in endometrioid ovarian cancers. This renders the *PI3K* pathway hyper-responsive to ligand in ovarian cancer cells. The *RAS* signaling pathway also impinges on *AKT* activity by activating *PI3K*. *SRC*, which is increased in amount and activity in ovarian cancer cells, impinges on the *PI3K* *AKT* pathway by increasing *PI3K* activity, as well as by phosphorylating and decreasing *PTEN* activity. NSAIDs inhibit Cox2, decreasing both production of matrix metalloproteinases and apoptosis. An ability to invade and bypass apoptosis is required for activation of the metastatic cascade. NSAIDs either through Cox2 or through as-yet unknown mechanisms can inhibit Lox15, Par4, ovulation, follicle-stimulating hormone production, activation of caspase proapoptotic mediators, *AKT* activation, and function of the *BCL-2* antiapoptotic mediator. In addition to these effects, acetaminophen, which is a relatively weak Cox2 inhibitor, may decrease the production of ovarian steroids and follicle-stimulating hormone, which can decrease ovulation and alter neoangiogenesis, respectively, thus decreasing tumor development.

significant chemopreventative activity against bowel cancer (2, 3). Given that other NSAIDs and particularly Cox2 inhibitors appear to demonstrate chemopreventative activity against bowel cancer (9–12), it seems likely that acetaminophen is exerting its effects through a non-Cox2-dependent mechanism.

The development of ovarian cancer has been strongly linked to the frequency of ovulation and also to the high concentrations of gonadotropins present postmenopausally (14–21). The association with ovulation has been linked to the proliferation of the ovarian epithelium, and potentially, the entrapment of ovarian surface epithelium cells within the stroma associated with repair of the ovulatory defect. Gonadotropins, in contrast, have been demonstrated to regulate neoangiogenesis in ovarian cancer models, a critical component of the development and metastases of tumors in many lineages.

The first and strongest epidemiological evidence demon-

strating that chemoprevention was effective at inhibiting the development of any major life-threatening cancer was derived from studies of ovarian cancer. Prolonged (5 years) use of the oral contraceptive pill significantly reduces the risk of developing ovarian cancer (odds ratio, 0.25–0.8; Refs. 22–27). Strikingly, decreased risk can be observed for at least 10 years after discontinuation of oral contraceptives. Although there are multiple potential causes, the most likely mechanism by which oral contraceptives prevent ovarian cancer is through decreasing the frequency of ovulation and, thus, the mutagenic effects of cellular proliferation and entrapment of the ovarian epithelium within the stroma.

The epidemiological evidence linking ovulation and gonadotropin levels to development of ovarian cancer suggests alternative mechanisms by which NSAIDs and, in particular, acetaminophen could impact the development of ovarian cancer (Fig. 1). Acetaminophen contains structural moieties

reminiscent of steroidal hormones potentially contributing to its ability, at high doses, to induce uterine, ovarian, and testicular atrophy in rats (2, 3, 28). This suggests that acetaminophen may decrease ovulation. In support of this contention, several studies have indicated that NSAIDs can decrease productive ovulation in animal and human models (29–32). Thus, acetaminophen could decrease the frequency of ovulation and the “wear-and-tear” damage on the at-risk tissue, the ovarian epithelium, that contributes to the accumulation of mutations in the epithelium and the development of ovarian cancer. Acetaminophen also alters glutathione metabolism potentially leading to alterations in gonadotropin levels (33). Thus, NSAIDs may exhibit activities on ovarian epithelium through multiple mechanisms unrelated to proliferation and survival.

Ovarian cancer, like all cancers, occurs as a consequence of either germ-line or acquired somatic changes in genetic function (14, 34–36). The required alterations in gene expression or function can occur as a result of mutations or of epigenetic alterations, such as changes in methylation (14, 34–36). One important challenge is to link the genotypic changes that occur in ovarian cancer cells to the phenotypic and biological changes observed in human tumors. Linking epidemiology, mechanistic studies of potential chemoprevention agents, and studies of the underlying genetic events leading to the development of ovarian cancer is critical for our understanding of the pathophysiology of ovarian cancer. Development of practical animal models to test hypotheses generated through genetic and epidemiological studies of human ovarian cancer should greatly facilitate the development of effective chemopreventative approaches for this devastating disease.

References

1. NIH Consensus Development Panel on Ovarian Cancer. Ovarian cancer: screening, treatment and follow-up. *JAMA*, 273: 491–497, 1995.
2. Rodriguez-Burford, C., Barnes, M. N., Oelschlager, D. K., Myers, R. B., Talley, L. I., Partridge, E. E., and Grizzle, W. E. *Clin. Cancer Res.*, 8: 202–209, 2001.
3. Cramer, D., Harlow, B., Titus-Ernstoff, L., Bohike, K., Welsh, W., and Greenberg, E. Over the counter analgesics and risk of ovarian cancer. *Lancet*, 351: 104–107, 1998.
4. Moyisch, K. B., Mettlin, C., Piver, M. S., Natarajan, N., Menezes, R. J., and Swede, H. Regular use of analgesic drugs and ovarian cancer risk. *Cancer Epidemiol. Biomarkers Prev.*, 10: 903–906, 2001.
5. Rosenberg, L., Palmer, J. R., Rao, R. S., Coogan, P. F., Strom, B. L., Zaubler, A. G., Stolley, P. D., and Shapiro, S. A case-control study of analgesic use and ovarian cancer. *Cancer Epidemiol. Biomarkers Prev.*, 9: 933–937, 2000.
6. Tavani, A., Gallus, S., La Vecchia, C., Conti, E., Montella, M., and Franceschi, S. Aspirin and ovarian cancer: an Italian case-control study. *Ann. Oncol.*, 11: 1171–1173, 2001.
7. Akhmedkhanov, A. A., Toniolo, P., Zeleniuch-Jacquotte, A., Kato, I., Koenig, K. L., and Shore, R. E. Aspirin and risk of epithelial cancer. *Preventive Medicine*, 33: 682–687, 2001.
8. Fredrickson, T. N. Ovarian tumors of the hen. *Environ. Health Perspect.*, 73: 35–51, 1987.
9. Hong, W., and Sporn, M. Recent advances in chemoprevention of cancer. *Science (Wash. DC)*, 278: 1073–1077, 1997.
10. Reddy, B. S., Rao, C. V., and Seibert, K. Evaluation of cyclooxygenase-2 inhibitor for potential chemopreventive properties in colon carcinogenesis. *Cancer Res.*, 56: 4566–4569, 1996.
11. Kawamori, T., Rao, C., Seibert, K., and Reddy, B. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, against colon carcinogenesis. *Cancer Res.*, 58: 409–412, 1998.
12. Giovannucci, E., Egan, K. M., Hunter, D. J., Stampfer, M. J., Colditz, G. A., Willett, W. C., and Speizer, F. E. Aspirin and the risk of colorectal cancer in women. *N. Engl. J. Med.*, 333: 609–614, 1995.
13. Ristimake, A., Honkanen, N., Nankala, H., Sipponen, P., and Harkonen, M. Expression of cyclooxygenase 2 in human gastric cancer. *Cancer Res.*, 57: 1276–1280, 1997.
14. Mills, G. B., Fang, X., Lu, Y., Hasegawa, Y., Eder, A., Erickson, J., Brewer, M., Fidler, I. J., Schmandt, R., Lu, K., Dubeau, L., Kavanagh, J., Gershenson, D., Shaw, P. A., Auersperg, N., Jaffe, R. B., Kohn, E., Gray, J., and Bast, R. C. *In: R. Freedman and J. Kavanagh (eds.), Biology of Ovarian Cancer*. New York: ISIS Press, in press, 2001.
15. Hildreth, N. G., Kelsey, J. L., LiVolsi, V. A., Fischer, D. B., Holford, T. R., Mostow, E. D., Schwartz, P. E., and White, C. An epidemiologic study of epithelial carcinoma of the ovary. *Am. J. Epidemiol.*, 114: 398–405, 1981.
16. Kvale, G., Heuch, I., Nilssen, S., and Beral, V. Reproductive factors and risk of ovarian cancer. A prospective study. *Int. J. Cancer*, 42: 246–251, 1988.
17. Mori, M., Harabuchi, I., Miyake, H., Casagrande, J. T., Henderson, B. E., and Ross, R. K. Reproductive, genetic, and dietary risk factors for ovarian cancer. *Am. J. Epidemiol.*, 128: 771–777, 1988.
18. Schildkraut, J. M., Bastos, E., and Berchuck, A. Relationship between lifetime ovulatory cycles and an overexpression of mutant p53 in epithelial ovarian cancer. *J. Natl. Cancer Inst. (Bethesda)*, 89: 932–938, 1997.
19. Webb, P. M., Green, A., Cummings, M. C., Purdie, D. M., Walsh, M. D., and Chenevix-Trench, G. Relationship between number of ovulatory cycles and accumulation of mutant p53 in epithelial ovarian cancer. *J. Natl. Cancer Inst. (Bethesda)*, 90: 1729–1734, 1998.
20. Yin, H., Cheng, K. W., Hwa, H. L., Peng, C., Auersperg, N., and Leung, P. C. Expression of the messenger RNA for gonadotropin-releasing hormone and its receptor in human cancer cell lines. *Life Sci.*, 62: 2015–2023, 1998.
21. Schiffenbauer, Y. S., Abramovitch, R., Meir, G., Nevo, N., Holzinger, M., Itin, A., Keshet, E., and Neeman, M. Loss of ovarian function promotes angiogenesis in human ovarian carcinoma. *Proc. Natl. Acad. Sci. USA*, 94: 13203–13208, 1997.
22. Harlap, S. Cancer benefits and risks of oral contraceptives. *In: V. DeVita, Jr., S. Hellman, and S. A. Rosenberg (eds). Cancer Prevention*, pp. 1–16. Philadelphia: J. B. Lippincott Co., 1992.
23. Prentice, R. L., and Thomas, D. B. On the epidemiology of oral contraceptives and disease. *Adv. Cancer Res.*, 49: 285–401, 1987.
24. Whittemore, A. S., Harris, R., Intyre, J. and Collaborative Ovarian Cancer Group. Characteristics relating to ovarian cancer risk: collaborative analysis of 12 U.S. case-control studies. II. Invasive epithelial ovarian cancers in white women. *Am. J. Epidemiol.*, 136: 1184–1203, 1992.
25. Franceschi, S., Parazzini, F., Negri, E., Booth, M., La Vecchia, C., Beral, V., Tzonou, A., and Trichopoulos, D. Pooled analysis of 3 European case-control studies. III. Oral contraceptive use. *Int. J. Cancer*, 49: 61–65, 1991.
26. Centers for Disease Control Cancer and Steroid Hormone Study. Oral contraceptive use and the risk of ovarian cancer. *JAMA*, 249: 1596–1599, 1983.
27. Centers for Disease Control Cancer and Steroid Hormone Study. The reduction in risk of ovarian cancer associated with oral contraceptive use. *N. Engl. J. Med.*, 316: 650–655, 1987.
28. National Toxicology Program. Toxicology and carcinogenesis studies of acetaminophen in F344/N rats and B6C3F1 mice Technical report

series no. 394. NIH publication no. 93-2849, Research Triangle Park, NC, 1993.

29. Ando, M., Kol, S., Irahara, M., Sirois, J., and Adashi, E. Y. Non-steroidal anti-inflammatory drugs (NSAIDs) block the late prostanoic-dependent/ceramide-independent component of ovarian IL-1 action: implications for the ovulatory process. *Mol. Cell. Endocrinol.*, 157: 21-30, 1999.

30. Mendonca, L. L., Khamashta, M. A., Nelson-Piercy, C., Hunt, B. J., and Hughes, G. R. Non-steroidal anti-inflammatory drugs as a possible cause for reversible infertility. *Rheumatology*, 39: 880-882, 2000.

31. Smith, G., Roberts, R., Hall, C., and Nuki, G. Reversible ovulatory failure associated with the development of leutenized unruptured follicles in women with inflammatory arthritis taking non-steroidal anti-inflammatory drugs. *Br. J. Rheumatol.*, 35: 458-462, 1996.

32. Ostensen, M., and Villiger, P. M. Nonsteroidal anti-inflammatory drugs in systemic lupus erythematosus. *Lupus*, 10: 135-139, 2001.

33. Pasha, K., and Vijayan, E. Glutathione distribution in rat brain at different ages and the effect of intraventricular glutathione on gonadotropin levels in ovariectomized steroid primed rats. *Brain Res. Bull.*, 22: 617-619, 1989.

34. Cahill, D. P., Kinzler, K. W., Vogelstein, B., and Lengauer, C. Genetic instability and Darwinian selection in tumours. *Trends Cell Biol.*, 9: 57-60, 1999.

35. Hanahan, D., and Weinberg, R. A. The hallmarks of cancer. *Cell*, 100: 57-70, 2000.

36. Lundberg, A. S., and Weinberg, R. A. Control of the cell cycle and apoptosis. *Eur. J. Cancer*, 35: 1886-1894, 1999.