

Selective Killing of Cancer Cells by Leaf Extract of Ashwagandha: Identification of a Tumor-Inhibitory Factor and the First Molecular Insights to Its Effect

Nashi Widodo,^{1,3} Kamaljit Kaur,¹ Bhupal G. Shrestha,¹ Yasuomi Takagi,² Tetsuro Ishii,³ Renu Wadhwa,¹ and Sunil C. Kaul¹

Abstract **Purpose:** Ashwagandha is regarded as a wonder shrub of India and is commonly used in Ayurvedic medicine and health tonics that claim its variety of health-promoting effects. Surprisingly, these claims are not well supported by adequate studies, and the molecular mechanisms of its action remain largely unexplored to date. We undertook a study to identify and characterize the antitumor activity of the leaf extract of ashwagandha. **Experimental Design:** Selective tumor-inhibitory activity of the leaf extract (i-Extract) was identified by *in vivo* tumor formation assays in nude mice and by *in vitro* growth assays of normal and human transformed cells. To investigate the cellular targets of i-Extract, we adopted a gene silencing approach using a selected small hairpin RNA library and found that p53 is required for the killing activity of i-Extract. **Results:** By molecular analysis of p53 function in normal and a variety of tumor cells, we found that it is selectively activated in tumor cells, causing either their growth arrest or apoptosis. By fractionation, purification, and structural analysis of the i-Extract constituents, we have identified its p53-activating tumor-inhibiting factor as withanone. **Conclusion:** We provide the first molecular evidence that the leaf extract of ashwagandha selectively kills tumor cells and, thus, is a natural source for safe anticancer medicine.

Ashwagandha (*Withania somnifera*, an evergreen shrub commonly found in the drier parts of the Indian subcontinent) is widely used in Indian natural medicine, Ayurveda. Extracts from different parts of ashwagandha have been claimed to promote physical and mental health due to its effects, ranging from antistress, antiinflammatory, antioxidant, antipyretic, analgesic, antiarthritic, antidepressant, anticoagulant, immunomodulatory, adaptogenic, cardioprotective, rejuvenating, and regenerating properties (1–13). Few reports have characterized the activities of the root extract of ashwagandha and include an induction of nitric oxide synthase-inducible protein expression (4, 14), down-regulation of p34cdc2 expression (15), and its antioxidant, free radical-scavenging, and detoxifying properties (16–19). However, the mechanistic aspects of its effects, including tumor suppression and isolation of active components, have largely

remained unexplored. Hence, the use of ashwagandha has not been developed to a systemic medicine.

Although ashwagandha roots are most commonly used in Indian Ayurvedic medicine, we undertook a study to examine the effects of its leaf extract because of the easy accessibility and abundant availability. We earlier reported that the leaf extract obtained by a series of extractions (20) has an antimutagenic effect (21). In the present study, we examined the effect of the leaf extract on human normal and cancer cells and found that it selectively kills tumor cells. Fractionation of the tumor-inhibitory extract (i-Extract) and characterization of its constituents led to the identification of a tumor-inhibitory factor (i-Factor). Nuclear magnetic resonance (NMR) spectra revealed its identity as withanone. By employing small hairpin RNA (shRNA) library and molecular analysis, we report for the first time that the selective killing of tumor cells by i-Extract and i-Factor involves an activation of the wild-type p53 function.

Authors' Affiliations: ¹Research Institute for Cell Engineering, ²GENE Therapeutics, Inc., National Institute of Advanced Industrial Science and Technology, Central 4, 1-1-1 Higashi, Tsukuba; and ³Department of Molecular and Cellular Physiology, University of Tsukuba, Ibaraki, Japan
Received 4/18/06; revised 10/10/06; accepted 11/30/06.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Grant support: The study was partially supported by grants from the New Energy and Industrial Technology Development Organization (NEDO), Japan.

Requests for reprints: Renu Wadhwa, National Institute of Advanced Industrial Science and Technology, Central 4, 1-1-1 Higashi, Tsukuba, Ibaraki 305 8562, Japan. Phone: 81-29-861-9464; Fax: 81-29-861-2900; E-mail: renu-wadhwa@aist.go.jp.

©2007 American Association for Cancer Research.
doi:10.1158/1078-0432.CCR-06-0948

Materials and Methods

Preparation of leaf extract from ashwagandha from field-raised plants. Ashwagandha (*W. somnifera*) leaf extracts were prepared as described earlier (20–22). The leaves were air dried, ground to a fine powder, and subjected to extraction with methanol (60°C) in Soxhlet apparatus for 4 to 5 days. The methanol extracts were further extracted with hexane to remove chlorophyll and other pigments and then with diethyl ether that was evaporated to obtain the ether extract. Ether extract solubilized in DMSO was used for the present studies.

Nude mice assay. BALB/c nude mice (4 weeks old, female) were bought from Nihon Clea (Japan). Mice were fed on standard food pellet and water *ad libitum*, acclimatized to our laboratory condition at

a temperature of $24 \pm 2^\circ\text{C}$, relative humidity of 55% to 65%, and 12-h light/dark cycle, for 3 days. Fibrosarcoma (HT1080) cells (1×10^6 suspended in 0.5 mL of growth medium) were injected s.c. into the flank of nude mice (one site per mouse). i-Extract injections were commenced at three different times, i.e., (a) mixed with cells at the time of injection, (b) injection before the formation of tumor buds, and (c) injection when small tumor buds (5 mm) were formed. In each case, s.c. injection (0.3 mL of 24 $\mu\text{g}/\text{mL}$ i-Extract in the cell growth medium) was given. Tumor formation was monitored during the next 15 to 20 days, with local injection of i-Extract to the tumor site every third day. For oral feeding, i-Extract or its components were suspended in 2% sterile carboxymethyl cellulose (a vehicle) and injected into the digestive tract of mice using a flexible Teflon needle on alternate days.

Characterization of i-Extract by column chromatography. Ether extract of the leaves was subjected to reversed-phase high-performance liquid chromatography (HPLC) analysis on a C-18 column (5 mm, 150×4.6 mm internal diameter; Waters, Milford, MA or YMC, Kyoto, Japan) at 40°C or 50°C using 1% methanol/ H_2O (solution A) and methanol/ethanol/isopropanol (52.25:45.30:2.45; solution B) for elution. Elution was done with a gradient of 35% to 45% solution B in 25 min at a flow rate of 1 mL/min. The detection was done at 220 nm. Withaferin A, 12-deoxywithastramonolide, and withanolide D were used as standards for comparison.

Human cell culture and treatments. Normal diploid fibroblasts (TIG-1 and WI-38), osteogenic sarcoma (U2OS and Saos-2), breast carcinoma (MCF7, HS578T, and SK-BR3), fibrosarcoma (HT1080), colon carcinoma (HCT116), and lung carcinoma (PC14) cells were cultured in DMEM (Life Technologies, Gaithersburg, MD), supplemented with 10% fetal bovine serum in a humidified incubator (37°C and 5% CO_2). Cells (~ 50 -60% confluency) were treated with i-Extract (6-36 $\mu\text{g}/\text{mL}$) for time periods as indicated.

Growth assays. Equal number of cells, counted by Neubauer hemocytometer, was plated in six-well dishes for control and treatment wells. Cells were harvested every 24 h up to 96 h, counted, and plotted as growth curves. Viability was monitored by WST-based cell proliferation kit⁴ (Roche, Mannheim, Germany).

Preparation and use of shRNAs. shRNAs for the genes listed in Table 1 were cloned in a U6-driven expression vector as described earlier (23). Two target sites per gene were used; the sequences for each of the target site are listed in Table 1. Cells were plated in 96-well plates and were transfected at $\sim 70\%$ confluency with 50 ng of the plasmid DNA. At 24 h posttransfection, cells were selected in puromycin (2 $\mu\text{g}/\text{mL}$) supplemented medium for 48 to 72 h, expanded to 70% confluency, and were then treated with i-Extract (24 $\mu\text{g}/\text{mL}$). shRNAs that resulted in the survival of cells (in the presence of i-Extract) were selected and taken through the second round of transfections and confirmation. Cell viability was measured by staining with Crystal Violet and AlamarBlue assay (BioSource International, Camarillo, CA).

Western blotting. The protein sample (20 μg) separated on SDS-polyacrylamide gel was electroblotted onto a nitrocellulose membrane (BA85, Schleicher and Schuell, Dassel, Germany) using a semidry transfer blotter. Immunoassays were done with anti-p53 (DO-1; Santa Cruz Biotechnology, Santa Cruz, CA), anti-actin (Chemicon International, Temecula, CA) and anti-p21^{WAF-1} (C-19; Santa Cruz Biotechnology) antibodies. The immunocomplexes formed were visualized with horseradish peroxidase-conjugated anti-rabbit immunoglobulin G (IgG; ECL; Amersham Pharmacia Biotech, Piscataway, NJ).

Immunoprecipitation. Control and i-Extract/i-Factor-treated cells were lysed in Nonidet P-40 lysis buffer. About 400 μg protein was incubated with wild-type p53-specific antibody (Ab-5; Calbiochem, Darmstadt, Germany) at 4°C for 1 to 2 h. Immunocomplexes were separated by incubation with Protein-A/G Sepharose, and Western

blotting was done with anti-p53-specific antibody (FL-393; Santa Cruz Biotechnology).

Immunostaining. Immunostaining was done as described earlier (24, 25). Cells were stained with anti-p53 antibody (DO-1; Santa Cruz Biotechnology), a monoclonal anti-mortalin antibody (MA3-028, Affinity Bioreagents, Golden, CO), or a polyclonal anti-mortalin antibody. Immunostaining was visualized by secondary staining with goat anti-rabbit IgG (Alexa-594-conjugated) and anti-mouse IgG (Alexa 488-conjugated; Molecular Probes, Eugene, OR).

Reporter assays. Cells were transfected with the p53-responsive luciferase reporter plasmid, PG-13luc (kindly provided by Dr. Bert Vogelstein, Johns Hopkins Medical Institutions, Baltimore, MD) and pPur plasmid (for puromycin resistance) in a 1:30 ratio. Transfected cells were selected in puromycin (2-3 $\mu\text{g}/\text{mL}$)-supplemented medium for 48 to 96 h and were then treated with i-Extract (24 $\mu\text{g}/\text{mL}$) for 48 h. Cells were lysed, and luciferase activity was measured by Dual-Luciferase reporter assay system (Promega, Madison, WI). Luciferase activity was calculated per microgram of protein and presented as the percent relative activity normalized against untreated cells. Results presented are from three independent experiments.

Terminal nucleotidyl transferase-mediated nick end labeling assays. Terminal nucleotidyl transferase (TdT)-mediated nick end labeling (TUNEL) assay was done using DeadEnd Fluorometric TUNEL System (Promega). Cells were grown on coverslips and treated with i-Extract for 48 h, after which they were fixed with 4% formaldehyde solution in PBS at 4°C for 20 min. Cells were washed with PBS and permeabilized with 0.2% Triton X-100 solution in PBS for 5 min and were then incubated with 100 μL equilibration buffer for 10 min followed by the addition of 50 μL rTdT incubation buffer for tailing reaction (37°C for 60 min in a humidified chamber). The reaction was terminated by incubation in $2 \times \text{SSC}$ for 15 min at room temperature. Cells were washed with PBS, stained with propidium iodide (1 $\mu\text{g}/\text{mL}$ in PBS), followed by washing with water and examined under a fluorescence microscope.

NMR analysis of i-Factor. One-dimensional NMR:¹H NMR (CDCl_3 , 500 MHz): d 0.86 (s, 3H, H-18), 1.04 (d, $J = 7$ Hz, 3H, H-21), 1.18 (s, 3H, H-19), 1.26 to 1.40 (m, 3H, H-11, 15), 1.53 to 1.62 (m, 2H, H-9, H-12), 1.67 to 1.80 (m, 2H, H-8, H-12), 1.88 (s, 3H, H-27), 1.94 (s, 3H, H-28), 1.88 to 1.95 (m, 2H, H-16), 1.98 to 2.07 (m, 1H, H-14), 2.30 to 2.35 (m, 1H, H-20), 2.42 to 2.56 (m, 4H, 17-OH, H-4, 23), 2.67 to 2.71 (m, 1H, H-4), 2.80 to 2.84 (m, 1H, H-11), 3.05 (d, $J = 3.7$ Hz, 1H, H-6), 3.15 to 3.16 (m, 1H, 5-OH), 3.31 to 3.33 (m, 1H, H-7), 4.59 to 4.63 (m, 1H, H-22), 5.85 (dd, $J = 10.1, 2.6$ Hz, 1H, H-2), 6.58 to 6.61 (m, 1H, H-3).¹³C NMR (CDCl_3 , 125 MHz): d 9.5 (C-21), 12.3 (C-27), 14.7 (C-19), 15.1 (C-18), 20.5 (C-28), 21.6 (C-11), 22.9 (C-15), 32.5 (C-12), 32.8 (C-23), 35.2 (C-9), 36.0 (C-8), 36.7 (C-16), 36.8 (C-4), 42.9 (C-20), 45.9 (C-14), 48.7 (C-13), 51.0 (C-10), 56.3 (C-6), 57.2 (C-7), 73.2 (C-5), 78.7 (C-22), 84.6 (C-17), 121.4 (C-25), 129.0 (C-2), 139.7 (C-3), 150.4 (C-24), 167.1 (C-26), 203.1 (C-1).

Results and Discussion

To examine the tumor-inhibitory effect of the leaf extract (i-Extract), we did nude mice assays using highly malignant human fibrosarcoma, HT1080 (mutant p53 and telomerase positive) cells. Subcutaneous injection of 10^6 cells resulted in the formation of solid tumors within 10 to 15 days (Fig. 1A). Injections of i-Extract, commenced either along with the injection of cells or before the formation of tumor buds or when small tumor buds (5 mm) were formed, resulted in tumor suppression. Of note, mice fed with i-Extract showed significant tumor suppression (Fig. 1B), suggesting that the i-Extract has a strong tumor-suppressive activity *in vivo*. We fractionated the i-Extract into its components by reverse-phase HPLC and identified the presence of withaferin A, withanolide

⁴ Cell Proliferation Reagent WST-1, a colorimetric assay for the quantification of cell proliferation and cell viability, based on the cleavage of the tetrazolium salt WST-1 by mitochondrial dihydrogenases in viable cells.

Table 1. List of genes and their target sequences of shRNA plasmids used to identify cellular targets of i-Extract

Serial number	Accession number	Gene name	Target sequence
1	NM_004936	CDKN2B	AATAGAGGTTGAACTTCAA
2	NM_004517	ILK	GATCTAAATTTGACGTGAT
3	NM_004517	ILK	GCGGGGATTTCAATGGAGA
4	NM_004064	p27 Kip1	AAGGGAGGGTTTATGTAGA
5	NM_004064	p27 Kip1	GAGGCAGTTGATATTTTCAT
6	NM_002611	PDK2	GGTCCAATTTCTAAGGTGGA
7	NM_002611	PDK2	AGGTCAACTTGCTTCTCGA
8	NM_002610	PDK1	GCTGGAGCTTAGAAGTCTA
9	NM_002610	PDK1	AATGGGAATTGAATTTTCAT
10	NM_002498	NEK3	GCTCCAGTATGGAGGCTA
11	NM_002498	NEK3	GTTTGGGAATTGAATGTAT
12	NM_002497	NEK2	GCTGGAATGTTAACTATA
13	NM_002497	NEK2	GATAGAAGTTGAGAAGCAA
14	NM_002376	MARK3	GAGGTAAGATTTAAATTTA
15	NM_002376	MARK3	AGTAAGGCCTCGATCTCTA
16	NM_002314	LIMK1	GCTAGAGGATTTATGGTGA
17	NM_002314	LIMK1	ACGAGAGGTTCAATTTTCAT
18	NM_001800	CDKN2D	GGTAGAGTATTTAAGGGTA
19	NM_001800	CDKN2D	GGGAGGAATACTTTCTCTT
20	NM_001799	CDK7	GGGTAAAGTGTTATGGGAA
21	NM_001799	CDK7	GGATATAAATTTAATGTTA
22	NM_001798	CDK2	GCTATAATGTTTAGAGAGA
23	NM_001798	CDK2	GGGTGAATAATTATGTTTA
24	NM_001262	CDKN2C	GTTCTGATGTAGTTGCA
25	NM_001262	CDKN2C	ATGTGAACGTTAATGTACA
26	NM_001261	CDK9	GGTCAAATGTGGACAGCTA
27	NM_001261	CDK9	GCTAGAGGGTGGCTTTGAA
28	NM_001260	CDK8	GCTTGAATATTTCAACGTTA
29	NM_001260	CDK8	GATAGAATATTTAGTGTA
30	NM_001259	CDK6	GATTAAGATTTGACCGCTT
31	NM_001259	CDK6	GGGAAGTCTATTTCTGAA
32	NM_001258	CDK3	GCGGTAAGTTCTATATCAA
33	NM_001258	CDK3	GGGAGGGATTGGAAGGGAT
34	NM_000077	CDKN2A	ACTAGAGAGGTTCTGGGAA
35	NM_000077	CDKN2A	ATTGCGGTGTCGCATGGTA
36	NM_000076	CDKN1C	GTGTGAAGCTTTGAGGGTA
37	NM_000076	CDKN1C	GTTCTAAAGTGTAAGGCTT
38	NM_000075	CDK4	GGGTGAGGGTAATCTGGAA
39	NM_000075	CDK4	AGGGTAGTCTTTGTCTTTA
40	NM_138793	SHAPY	GCGGGGGGTTGTAGGGAAA
41	NM_138793	SHAPY	GCGTGGAAATATGAGGGCAT
42	NM_000314	PTEN	GGTGGAGCTTGAATTTCTA
43	NM_000314	PTEN	GGGATAATATTGATGGTGT
44	NM_000546	p53	GCTAAAGAAGAAATCGCA
45	NM_000546	p53	GCTGGGAGGGTTAATATCT
46	NM_000077	p16INK4A	GAGTGTGAGAGTCTCTAAT
47	NM_000077	p16INK4A	GATAACTCTGATGTTTGAG
48	NM_015322	FEM	ATTTTCATGTATAAAGGAC
49	NM_015322	FEM	GTCATAAACTTGACACAGAC
50	NM_002156	HSP60	GGTACTAGCTCTTCATTTT
51	NM_002156	HSP60	GCTCGAGTCTTAATGCTTG

Table 1. List of genes and their target sequences of shRNA plasmids used to identify cellular targets of i-Extract (Cont'd)

Serial number	Accession number	Gene name	Target sequence
52	NM_003156	STIM1	GGGAAGACCTCAATTACCA
53	NM_003156	STIM1	GGCCAAGAAGACATTATA
54	NM_014622	AW551984	GAACAACTTCTATCTTTA
55	NM_014622	AW551984	AGTCAAACCTGTAAATAGA
56	NM_007215	POLG2	GAAGCAAACCTTACTACAA
57	NM_007215	POLG2	GGGAAAGGAGCCAATAGAA
58	NM_016248	PRKA-2	GTGAAGCTTTATCAAATGC
59	NM_016248	PRKA-2	ACAACATGGAAAGTCATG
60	NM_005923	MAP3K5	GGGACAGCTTTCAAATTA
61	NM_005923	MAP3K5	ATCTGTTTCTGTATAGAAT
62	NM_004672	MAP3K6	GTGGGTTGCTTAAGGTTTC
63	NM_004672	MAP3K6	AGGGATCGTTCTCACAGA
64	NM_145333	MAP3K7	GGGTGAATTTGGATGTTTA
65	NM_145333	MAP3K7	AGGTGTTGCTATTAATATA
66	NM_005204	MAP3K8	GTAAAGCACCTTCACTGGAG
67	NM_005204	MAP3K8	AGGAAGTGATTCATTATGA
68	NM_001880	ATF2	GGGTAGCTTTTACTCAC
69	NM_001880	ATF2	AGTTCTGGCTTAAGTGTAG
70	NM_002759	PRKR	AAGATAATTGTAATTTG
71	NM_002759	PRKR	ATTGGTCGCTAAATTTGCG
72	NM_005157	ABL1	GGTCTAGCTTTTACTCCTG
73	NM_005157	ABL1	GTTCAGCTCTGCTACCTAT
74	NM_001626	AKT2	ATCTCATGTTGGACAAAGG
75	NM_001626	AKT2	GGCTTCGATTATCTCAAT
76	NM_001166	BIRC2	AGGTAAGCTTTGAATGATA
77	NM_001166	BIRC2	GGATATGTCTTAATGAGA
78	NM_001165	BIRC3	ACTTTGACTTTCTATTCTAG
79	NM_001165	BIRC3	AGCTTTAAATGTATAGAAA
80	NM_001167	BIRC4	GTATCATACTGTAACCTGAA
81	NM_001167	BIRC4	GTGTTGGACTTTACTACAC
82	NM_001168	BIRC5	ATTGTTGATTCATAGAATG
83	NM_001168	BIRC5	GTAGGTTCCCTGTCTGTCA
84	NM_000633	BCL2 α	AGTGTGAATTTAACTTTA
85	NM_000633	BCL2 α	AAGTAAGGCTTTGAATGAT
86	NM_004935	CDK5	ATGGTGGCCTTGATCTTGA
87	NM_004935	CDK5	GGGTGGGGTTGGATGATGA
88	NM_023109	FGFR1	GGTGGAACTTACGCCCGG
89	NM_023109	FGFR1	ATTTAAACCTGATCACAGA
90	NM_000142	FGFR3	GTGTATATTTCTGTAATG
91	NM_005248	FGR	GGTCCGGCCTGTATCCTCG
92	NM_005248	FGR	GGGCCATCTATATCGTGA
93	NM_000875	IGF1R	ATACCATCCTGCATAATAC
94	NM_000875	IGF1R	AGTTATGGTTTAAATTTAA
95	NM_000208	INSR	GGTGAGATTTATATGATTA
96	NM_000208	INSR	GGAATGACTTTCATTCTGG

NOTE: shRNA (1, 12, 36, 44, 78, and 86) that resulted in the abrogation of the effect of i-Extract are indicated in bold.

A, and 12-deoxywithastramonolide (Fig. 2A). These components, however, did not show any antitumor activity either by local injections into the tumor sites or by oral feeding (Fig. 1B and data not shown). Upon HPLC analysis of different batches of i-Extract that were used for nude mice assays, we noticed that the content of its component that elutes at 22 min in HPLC correlated positively with its tumor suppressor activity (Fig. 2A). We purified this ingredient of interest from i-Extract and named it as i-Factor because of its tumor-inhibitory activity (Fig. 2B). We next examined the effect of i-Factor in *in vivo* and *in vitro*

tumor assays as described below. As shown in Fig. 2C, i-Factor showed tumor suppressor activity, similar to i-Extract, and was not toxic to mice. On the other hand, when mice were given with either withaferin A or 12-deoxywithastramonolide, they lost weight and looked sick (data not shown). Most interestingly, in *in vitro* assays, (a) both i-Extract and i-Factor selectively killed human tumor cells and (b) i-Factor, when added along with withaferin A, partially neutralized the toxicity of the latter in normal human cells (Fig. 2D). The antitumor activity of i-Extract was supported by treating a large variety of

human transformed (including MCF7, U2OS, Saos-2, SKBR3, HS578T, PC-14, HCT116, HeLa, HT1080) cells, whereas normal (MRC-5, TIG-1, and WI-38) cells were least affected (data not shown). Whereas withaferin A was toxic to both cancer and normal cells, withanolide A and 12-deoxywithastramonolide did not show any effect on either of the cell types (data not shown).

We next set out to characterize the molecular mechanism of i-Extract-induced growth arrest of cancer cells. To get an insight to the signaling pathways that are involved in i-Extract-induced death of cancer cells, we prepared 96 shRNA expression plasmids to target 48 genes (most of them were

targeted to kinases; Table 1) as described in Materials and Methods. Six shRNAs that had target sites for (a) cyclin-dependent kinase inhibitor 2B (CDKN2B), (b) never in mitosis arrest-related kinase 2, (c) cyclin-dependent kinase 8 (CDK8), (d) tumor suppressor protein (p53), (e) inhibitor of apoptosis protein 1/baculoviral inhibitor of apoptosis repeat-containing 3 (BIRC3), and (f) CDK5 resulted in the abrogation of cell death induced by i-Extract. In the present study, we analyzed the involvement of p53 in i-Extract- and i-Factor-induced selective death of cancer cells as described below.

Normal (TIG-1) and tumor-derived (U2OS with wild-type p53 protein) cells were treated with i-Extract or i-Factor. The




Fig. 1. Tumor suppression by i-Extract in nude mice assays. *A*, *in vivo* tumor suppression by i-Extract in nude mice. Control HT1080 cells formed a large tumor in 6 to 15 d. Local injections of i-Extract (0.3 mL of 24 μ g/mL) to tumor sites significantly inhibited their growth. Injections were repeated every third day. Columns, average tumor size (mm) in control and i-Extract – injected mice. *B*, oral feeding of i-Extract (100 mg/kg body weight), but not of withaferin A (10 mg/kg body weight), withanolide A (10 mg/kg body weight), or 12-deoxywithastramonolide (10 mg/kg body weight), resulted in tumor suppression. Columns, average tumor size (mm) in control and treated mice.





Fig. 2. Fractionation of i-Extract by reverse-phase HPLC. *A*, the identified components included withaferin A, withanolide A, and 12-deoxywithastramonolide. In addition, a component that eluted at 22 min was identified in all the i-Extract preparations, and its amount correlated with the tumor suppressor activity (TSA) of i-Extract, shown by strong (+++), moderate (++), and low (-). *B*, purification of peak 22 component. *C*, tumor suppression by i-Extract (0.3 mL of 24 μ g/mL) and i-Factor (0.3 mL of 10 μ g/mL), but not by withaferin A (0.3 mL of 10 μ g/mL). Columns, average tumor size (mm) in control and treated mice. *D*, selective killing of tumor cells by i-Extract (24 μ g/mL) and i-Factor (10 μ g/mL). Whereas withaferin A (0.1 μ mol/L) was toxic to normal cells also, withanolide A (1 μ mol/L) and 12-deoxywithastramonolide (1 μ mol/L) did not show any activity. i-Factor (10 μ g/mL) reduced the toxicity of withaferin A in normal cells.

level of p53 protein and its transcriptional activation function (p21^{WAF-1} level) increased in U2OS, but not in normal (TIG-1) cells (Fig. 3A). Interestingly, i-Factor-treated cells showed a moderate decrease in p53 level and its transcriptional activation function only in normal cells (Fig. 3A and data not shown). In contrast, withaferin A led to an induction of the p53 function in normal cells, and i-Factor partially neutralized its effect (Fig. 3A). The activity of wild-type p53 is regulated by multiple factors, including its interactions with binding partners. In transformed cells, mortalin, a hsp70 family member, interacts and inactivates p53 by sequestering it in the cytoplasm (24–28). Pancytoplasmic subcellular distribution of mortalin has been shown as a marker for normal cells; tumor cells show perinuclear staining pattern (29). Interestingly, a shift in perinuclear mortalin staining pattern, typical of tumor cells, to pancyto-

plasmic pattern was observed in i-Extract-treated tumor cells. Tumor cells that were induced to undergo senescence-like growth arrest in response to treatment with a rhodacyanine dye (MKT-077) or with bromodeoxyuridine also showed a shift in perinuclear staining to the pancytoplasmic type (24, 30). Taken together, these data suggested that the i-Extract abrogated mortalin-p53 interactions and induced senescence-like growth arrest in cancer cells through an activation of wild-type p53 function. Similar to i-Extract, i-Factor induced a shift in the staining pattern of mortalin and nuclear translocation of p53 in cancer cells (Fig. 3B). The induction of senescence was also supported by positive senescence-associated β -gal staining in i-Extract- and i-Factor-treated cells (data not shown).

We next used tumor-derived cell lines [breast carcinomas HS578T (V157F), SK-BR3 (H175V), and MDA-MB-436

Fig. 3. i-Extract and i-Factor activate wild-type tumor suppressor p53 in cancer cells. **A**, Western blot analysis showing the activation of p53 and its downstream effector p21^{WAF-1} by i-Extract (24 μ g/mL) and i-Factor (10 μ g/mL) in osteosarcoma (U2OS), but not in normal human skin fibroblasts (TIG-1). Withaferin A (0.1 μ mol/L) activated the wild-type p53 function both in cancer and normal cells. Actin was used as a loading control. **B**, i-Extract caused a shift in the staining pattern of mortalin (red) in cancer (U2OS) cells. Consistent with the shift of mortalin staining pattern (from perinuclear to pancytoplasmic), p53 was translocated into the nucleus of cancer cells (green nuclear staining). **C** and **D**, cell viability assay of normal and tumor-derived human cells grown in normal, i-Extract, or i-Factor-supplemented culture medium. All the cancer cells were growth arrested; normal cells remained unaffected. **E**, i-Extract restored the wild-type p53 characteristics and function to the mutant p53 as detected by a decrease in the level of p53 and increased level of p21^{WAF-1}, MDM2, and Bax. **F**, immunoprecipitation of wild-type p53 in i-Extract-treated mutant p53-possessing cells. **G**, similar to i-Extract, i-Factor led to a decrease in the level of p53 protein in mutant p53 cells. **H**, detection of wild-type p53-specific reporter activity in i-Extract-treated cells. p53-dependent luciferase activity in i-Extract/i-Factor-treated cells is plotted as percent relative activity to that of the untreated control cells (U2OS, wild-type; HT1080, wild-type and mutant p53; and PC14, mutant p53). **I**, i-Factor triggered apoptosis as detected by TUNEL staining in mutant p53 containing (SKBR3) cells; no apoptosis was detected in HeLa and U2OS cells.






Fig. 4. One-dimensional NMR spectra of i-Factor and a structure of withanone. One-dimensional NMR of i-Factor was done in CDCl_3 . *A*, ^1H NMR spectrum. *B*, ^{13}C NMR spectrum. *C*, withanone.

(H273V); lung carcinoma PC14 (Q248V); and fibrosarcoma HT1080 (mutant p53)] that harbor full-length p53 protein with point mutations resulting in one amino acid change (as indicated in parenthesis) in the core domain that recognizes p53 DNA-binding sites. Growth of all these cells was also inhibited

by i-Extract and i-Factor (Fig. 3C and D), very similar to the ones that contain wild-type but inactivated p53. We analyzed the function of p53 in these tumor cells with mutant p53. Noticeably, i-Extract restored the wild-type p53 function, at least in part, in these cells. In response to i-Extract treatment, the level

of p53 decreased (typical of wild-type p53 degradation) in three of the four (PC14, HS578T, and HT1080) cell lines. This was accompanied by the enhanced expression of, at least, two of the three downstream effectors of wild-type p53 (p21^{WAF-1}, MDM2, and Bax; Fig. 3E). Although SKBR3 cells did not show a decrease in p53 level subsequent to the i-Extract treatment, the increase in MDM2 and Bax supported the presence of wild-type activity. It was further confirmed by immunoprecipitation with the wild-type p53-specific antibody (Fig. 3F). On the similar line, Bax was not detected in HT1080 and HS578T cells. These data indicate the cell line specific response, operative cellular factors, and pathways that warrant further molecular analysis. Immunoprecipitation with wild-type p53-specific antibody revealed the existence of wild-type p53 protein only in i-Extract-treated PC14, HS578T, SK-BR3, and HT1080 cells (Fig. 3F). Of note, the induction of wild-type p53 in HT1080 was not as significant as in the other three cell lines (Fig. 3E). Consistent with this, the p21^{WAF-1} level in HT1080 cells showed only a small increase (Fig. 3D). Similar to i-Extract, i-Factor-treated tumor cells also showed a decrease in the level of p53 (Fig. 3G) and the wild-type p53-dependent reporter activity (Fig. 3H). We next examined the induction of apoptosis by i-Extract in tumor cells with variable (wild-type: functional or nonfunctional; and mutant type) p53 status. We found that the i-Extract induced apoptosis in tumor cells with mutant p53 (HS578T, SK-BR3, and PC14). The tumor cells (U2OS, MCF7, and HeLa) with wild-type p53 (functional or nonfunctional) exhibited growth arrest (Fig. 3I). These data were similar to another study that reported the induction of apoptosis by wild-type p53 protein expression in HT29A4 (possess mutant p53) cells (31). The data showed that i-Extract restored the wild-type function (transcriptional activation) of the mutant p53. Such an ability to endow wild-type p53 function (reactivation of the growth arrest or the apoptotic pathway)

to mutated p53 protein is regarded to be the most beneficial approach for cancer therapeutics (32, 33). Similar reactivation of p53 was achieved by low-molecular-weight peptides, including PRIMA-1 (34–37) and by global suppressor motif approach (38). Although the molecular mechanism of induction of p53 function by i-Extract and i-Factor warrants further studies, our data showed that they could serve as very useful natural tools for such reactivation of p53. We therefore determined the structure of i-Factor by NMR analysis. As shown in Fig. 4, ¹H NMR and ¹³C NMR data of isolated i-Factor were consistent with published values for withanone (39, 40). The assignment and its structure were also confirmed by 2D/NMR: ¹H-¹H correlated spectroscopy, heteronuclear multiple quantum coherence, heteronuclear multiple bond coherence, and homonuclear Hartman Hahn (data not shown). Electrospray-mass spectrometry of i-Factor showed *m/z* 493.2 [M + Na⁺] that was also consistent with a theoretical value of withanone (C₂₈H₃₆O₆: *M*, 470.58). Based on these data, we concluded that i-Factor is withanone and is a potential anticancer drug.

Identification of novel natural anticancer compounds is a highly demanding avenue of cancer therapeutics. In Indian traditional Ayurvedic medicine, ashwagandha is used to treat several illnesses, including tumors, inflammations, conjunctivitis, and tuberculosis. We have shown here, for the first time, that ashwagandha leaves have inhibitory activity selective for tumor cells that works, at least in part, through the activation of wild-type p53 activity. i-Extract and i-Factor, thus, are valuable sources for safe antitumor medicine.

Acknowledgments

We thank Nippon Shinyaku Co., Ltd. (Japan) for help in the partial characterization and purification of i-Extract and i-Factor.

References

- Scartezzini P, Speroni E. Review on some plants of Indian traditional medicine with antioxidant activity. *J Ethnopharmacol* 2000;71:23–43.
- Archana R, Namasivayam A. Antistressor effect of *Withania somnifera*. *J Ethnopharmacol* 1999;64:91–3.
- Davis L, Kuttan G. Effect of *Withania somnifera* on cell mediated immune responses in mice. *J Exp Clin Cancer Res* 2002;21:585–90.
- Davis L, Kuttan G. Effect of *Withania somnifera* on CTL activity. *J Exp Clin Cancer Res* 2002;21:115–8.
- Singh B, Chandan BK, Gupta DK. Adaptogenic activity of a novel withanolide-free aqueous fraction from the roots of *Withania somnifera* Dun. (part II). *Phytother Res* 2003;17:531–6.
- Bhattacharya SK, Muruganandam AV. Adaptogenic activity of *Withania somnifera*: an experimental study using a rat model of chronic stress. *Pharmacol Biochem Behav* 2003;75:547–55.
- Gupta YK, Sharma SS, Rai K, Katiyar CK. Reversal of paclitaxel induced neutropenia by *Withania somnifera* in mice. *Indian J Physiol Pharmacol* 2001;45:253–7.
- Prakash J, Gupta SK, Dinda AK. *Withania somnifera* root extract prevents DMBA-induced squamous cell carcinoma of skin in Swiss albino mice. *Nutr Cancer* 2002;42:91–7.
- Mishra LC, Singh BB, Dagenais S. Scientific basis for the therapeutic use of *Withania somnifera* (ashwagandha): a review. *Altern Med Rev* 2000;5:334–46.
- Dhuley JN. Adaptogenic and cardioprotective action of ashwagandha in rats and frogs. *J Ethnopharmacol* 2000;70:57–63.
- Bhattacharya SK, Bhattacharya A, Sairam K, Ghosal S. Anxiolytic-antidepressant activity of *Withania somnifera* glycowithanolides: an experimental study. *Phytomedicine* 2000;7:463–9.
- Tohda C, Kuboyama T, Komatsu K. Dendrite extension by methanol extract of ashwagandha (roots of *Withania somnifera*) in SK-N-SH cells. *Neuroreport* 2000;11:1981–5.
- Kuboyama T, Tohda C, Zhao J, Nakamura N, Hattori M, Komatsu K. Axon- or dendrite-predominant outgrowth induced by constituents from ashwagandha. *Neuroreport* 2002;13:1715–20.
- Iuvone T, Esposito G, Capasso F, Izzo AA. Induction of nitric oxide synthase expression by *Withania somnifera* in macrophages. *Life Sci* 2003;72:1617–25.
- Singh DD, Dey CS, Bhutani KK. Downregulation of p34cdc2 expression with aqueous fraction from *Withania somnifera* for a possible molecular mechanism of anti-tumor and other pharmacological effects. *Phytomedicine* 2001;8:492–4.
- Prakash J, Gupta SK, Kochupillai V, Singh N, Gupta YK, Joshi S. Chemopreventive activity of *Withania somnifera* in experimentally induced fibrosarcoma tumours in Swiss albino mice. *Phytother Res* 2001;15:240–4.
- Russo A, Izzo AA, Cardile V, Borrelli F, Vanella A. Indian medicinal plants as antiradicals and DNA cleavage protectors. *Phytomedicine* 2001;8:125–32.
- Panda S, Kar A. Evidence for free radical scavenging activity of ashwagandha root powder in mice. *Indian J Physiol Pharmacol* 1997;41:424–6.
- Davis L, Kuttan G. Effect of *Withania somnifera* on DMBA induced carcinogenesis. *J Ethnopharmacol* 2001;75:165–8.
- Lavie D, Kirson I, Glotter E. Constituents of *Withania somnifera* Dun. part X. The structure of withanolide D. *Isr J Chem* 1968;6:671–8.
- Rani G, Kaur K, Wadhwa R, Kaul SC, Nagpal A. Evaluation of the anti-genotoxicity of leaf extract of ashwagandha. *Food Chem Toxicol* 2005;43:95–8.
- Kaur K, Rani G, Widodo N, et al. Evaluation of the anti-proliferative and anti-oxidative activities of leaf extract from *in vivo* and *in vitro* raised ashwagandha. *Food Chem Toxicol* 2004;42:2015–20.
- Wadhwa R, Yaguchi T, Kaur K, et al. Use of a randomized hybrid ribozyme library for identification of genes involved in muscle differentiation. *J Biol Chem* 2004;279:51622–9.
- Wadhwa R, Sugihara T, Yoshida A, et al. Selective toxicity of MKT-077 to cancer cells is mediated by its binding to the hsp70 family protein mot-2 and reactivation of p53 function. *Cancer Res* 2000;60:6818–21.
- Wadhwa R, Takano S, Robert M, et al. Inactivation of tumor suppressor p53 by mot-2, a hsp70 family member. *J Biol Chem* 1998;273:29586–91.
- Mihara M, Erster S, Zaika A, et al. p53 has a direct apoptogenic role at the mitochondria. *Mol Cell* 2003;11:577–90.
- Dumont P, Leu JI, Della Pietra AC III, George DL, Murphy M. The codon 72 polymorphic variants of p53 have markedly different apoptotic potential. *Nat Genet* 2003;33:357–65.

28. Wadhwa R, Yaguchi T, Hasan MK, Mitsui Y, Reddel RR, Kaul SC. Hsp70 family member, mot-2/mthsp70/GRP75, binds to the cytoplasmic sequestration domain of the p53 protein. *Exp Cell Res* 2002;274:246–53.
29. Wadhwa R, Kaul SC, Mitsui Y, Sugimoto Y. Differential subcellular distribution of mortalin in mortal and immortal mouse and human fibroblasts. *Exp Cell Res* 1993;207:442–8.
30. Michishita E, Nakabayashi K, Suzuki T, et al. 5-Bromo-deoxyuridine induces senescence-like phenomena in mammalian cells regardless of cell type or species. *J Biochem* 1999;126:1052–9.
31. Barberi-Heyob M, Veldre PO, Merlin JL, et al. Wild-type p53 gene transfer into mutated p53 HT29 cells improves sensitivity to photodynamic therapy via induction of apoptosis. *Int J Oncol* 2004;24:951–8.
32. Lain S, Lane D. Improving cancer therapy by non-genotoxic activation of p53. *Eur J Cancer* 2003;39:1053–60.
33. Lane D. Curing cancer with p53. *N Engl J Med* 2004;350:2711–2.
34. Bykov VJ, Selivanova G, Wiman KG. Small molecules that reactivate mutant p53. *Eur J Cancer* 2003;39:1828–34.
35. Bykov VJ, Wiman KG. Novel cancer therapy by reactivation of the p53 apoptosis pathway. *Ann Med* 2003;35:458–65.
36. Bykov VJ, Issaeva N, Shilov A, et al. Restoration of the tumor suppressor function to mutant p53 by a low-molecular-weight compound. *Nat Med* 2002;8:282–8.
37. Bykov VJ, Issaeva N, Selivanova G, Wiman KG. Mutant p53-dependent growth suppression distinguishes PRIMA-1 from known anticancer drugs: a statistical analysis of information in the National Cancer Institute database. *Carcinogenesis* 2002;23:2011–8.
38. Baroni TE, Wang T, Qian H, et al. A global suppressor motif for p53 cancer mutants. *Proc Natl Acad Sci U S A* 2004;101:4930–5.
39. Kirson I, Glotter E, Lavie D, Abraham HR. Constituents of *Withania somnifera* Dunal. XII. The withanolides of an Indian chemotype. *J Chem Soc C Org Chem* 1971;11:2032–44.
40. Gottlieb HE, Kirson I. ¹³C NMR spectroscopy of the withanolides and other highly oxygenated C₂₈ steroids. *Org Magn Reson* 2005;16:20–5.

Clinical Cancer Research

Selective Killing of Cancer Cells by Leaf Extract of Ashwagandha: Identification of a Tumor-Inhibitory Factor and the First Molecular Insights to Its Effect

Nashi Widodo, Kamaljit Kaur, Bhupal G. Shrestha, et al.

Clin Cancer Res 2007;13:2298-2306.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/13/7/2298>

Cited articles This article cites 40 articles, 4 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/13/7/2298.full#ref-list-1>

Citing articles This article has been cited by 3 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/13/7/2298.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/13/7/2298>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.