

Histone γ H2AX and Poly(ADP-Ribose) as Clinical Pharmacodynamic Biomarkers

Christophe E. Redon¹, Asako J. Nakamura¹, Yong-Wei Zhang¹, Jiuping (Jay) Ji², William M. Bonner¹, Robert J. Kinders³, Ralph E. Parchment³, James H. Doroshow^{1,4}, and Yves Pommier¹

Abstract

Tumor cells are often deficient in DNA damage response (DDR) pathways, and anticancer therapies are commonly based on genotoxic treatments using radiation and/or drugs that damage DNA directly or interfere with DNA metabolism, leading to the formation of DNA double-strand breaks (DSB), and ultimately to cell death. Because DSBs induce the phosphorylation of histone H2AX (γ H2AX) in the chromatin flanking the break site, an antibody directed against γ H2AX can be employed to measure DNA damage levels before and after patient treatment. Poly(ADP-ribose) polymerases (PARP1 and PARP2) are also activated by DNA damage, and PARP inhibitors show promising activity in cancers with defective homologous recombination (HR) pathways for DSB repair. Ongoing clinical trials are testing combinations of PARP inhibitors with DNA damaging agents. Poly(ADP-ribosylation), abbreviated as PAR, can be measured in clinical samples and used to determine the efficiency of PARP inhibitors. This review summarizes the roles of γ H2AX and PAR in the DDR, and their use as biomarkers to monitor drug response and guide clinical trials, especially phase 0 clinical trials. We also discuss the choices of relevant samples for γ H2AX and PAR analyses. *Clin Cancer Res*; 16(18); 4532–42. ©2010 AACR.

The field of cancer treatment has a critical need to accelerate the validation of candidate drugs, while ultimately reducing costs. One approach promising to improve the efficiency and speed of clinical trials uses biological markers to measure pharmacodynamic parameters in samples taken from the cancer patients themselves during drug protocols (1). Because many anticancer agents target DNA and the DNA damage response (DDR) pathways, biomarkers are based on DNA damage endpoints. Biomarkers may also be useful in identifying individuals who are hypersensitive to radiotherapy in order to tailor treatments to minimize undesired side effects. The most desirable biomarkers would be those that are able to determine early in the course

of treatment whether a drug is reaching its target and acting as intended. The development of new biomarkers for clinical trials has been seen in the last few years (2), especially in the context of phase 0 trials (1). Here, we discuss two pharmacodynamic biomarkers, γ H2AX and PAR, currently being developed at the NIH Clinical Center (Bethesda, MD) by the NCI.

γ H2AX

H2AX belongs to the H2A family of histones, one of four histone families present in the nucleosomes that package eukaryotic DNA into chromatin (3). Upon double-strand break (DSB) formation, hundreds of H2AX molecules in the chromatin are rapidly phosphorylated by members of the phosphatidylinositol-3 kinase (PI3K) family (4, 5), forming a focus at the DSB site (see Fig. 1 for details). H2AX-deficient cells exhibit cell-cycle checkpoint deficiency, increased genomic instability, and sensitivity to genotoxic agents (6), observations corroborating findings of roles for γ H2AX in multiple processes necessary to return the broken chromatin to its original state (see Fig. 1 for details). After completion of DNA repair and chromatin remodeling, dephosphorylation of γ H2AX is carried out by several phosphatases including the p53-inducible phosphatase Wip1 (7, 8), protein phosphatases 6 (9), 4 (10), and 2A (11).

Since its discovery in 1998 at the NCI (5), γ H2AX has been a topic of basic research analyzing the DDR, and a topic of translational studies as a biosimulator to measure the genotoxic effects of drugs and/or radiation exposure

Authors' Affiliations: ¹Laboratory of Molecular Pharmacology, Center for Cancer Research, National Cancer Institute (NCI), ²National Clinical Target Validation Laboratory and ³Laboratory of Human Toxicology and Pharmacology, SAIC-Frederick, Inc., NCI-Frederick, Frederick, Maryland; and ⁴Division of Cancer Treatment and Diagnosis, NCI, Bethesda, Maryland

Note: This project has been funded in whole or in part with federal funds from the National Cancer Institute, National Institutes of Health, under Contract No. HHSN261200800001E. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

Corresponding Author: Yves G. Pommier, NIH/NCI, 37 Convent Drive, Building 37, Room 5068, Bethesda, MD 20892. Phone: 301-496-5944; Fax: 301-402-0752; E-mail: pommier@nih.gov.

doi: 10.1158/1078-0432.CCR-10-0523

©2010 American Association for Cancer Research.




Fig. 1. γ H2AX following DSB formation. H2AX is phosphorylated by members of the PI3K family; the member involved depends on the type of genotoxic stress (4). Although ATM and ATM- and Rad3-related (ATR) are primarily involved in H2AX phosphorylation following ionizing radiation and replication stress, respectively, DNA-PK and JNK-1 were shown to be responsible for γ H2AX formation during apoptosis (97, 98). γ H2AX foci are known to be involved in the recruitment and stabilization of DDR proteins including Mre11, Rad50, Nbs1 (the MRN complex), MDC1, 53BP1, BRCA1, ATM, and RNF8 (6, 99–101). DSB repair is done by the homologous recombination (HR) and NHEJ pathways. HR, driven by the *BRCA2*, *RAD51*, and *RAD52/54* genes, is more accurate because it uses a homologous DNA segment to act as a template for the damaged DNA region. Repair is also done by sister chromatid-dependent recombination repair via cohesin recruitment (102, 103). In contrast, NHEJ is faster, does not require a homologous DNA segment, and can operate in nonreplicating cells. However, it is error prone. The classical effectors of NHEJ are the end-binding proteins Ku70/80, DNA-PKcs, the nuclease Artemis, the scaffolding protein XRCC4, and ligase IV. Recently, a slow DNA-PK-independent NHEJ pathway involving PARP1, histone H1, XRCC1, and ligase III has been proposed (24, 25). The γ H2AX foci are also involved in chromatin alteration via recruitment of remodeling complexes and in signal transduction (accrued ATM activation, G2/M cell-cycle checkpoint). In addition, γ H2AX foci, through their recruitment of the cohesins and the MRN complexes, are involved in binding and tethering the broken DNA ends, and may help prevent the dissociation of the broken chromosome ends (104).

(4, 5). γ H2AX foci are found in almost all cell types after exposure to exogenous agents that directly induce DSBs, that is radiation therapy and classical chemotherapy with genotoxic drugs. Importantly, however, treatment with other drugs such as inhibitors of DNA topoisomerases and/or drugs interfering with DNA replication also generate a variety of DNA lesions that may result in the formation of DSBs (12), primarily in S-phase cells. It is important to note that γ H2AX foci, usually in small numbers, may be present in cells even in the absence of exogenous damage (4), because of DNA damage that can occur during many common cellular processes, including DNA replication, cellular senescence, and exposure to endogenous reactive oxygen species (ROS) DSBs may form during these processes or because of DNA repair defects, producing genomic instability and increasing cancer risk, as exemplified by the increased cancer incidence in individuals with genetic defects in DNA repair genes, such as *BRCA1*, *BRCA2*, the *FANC* genes, *ATM*, and *CHEK2*.

Poly(ADP-Ribose) Polymerases and Poly(ADP-Ribose) Polymers

Poly (ADP-ribose) polymerases (PARP) belong to a family of 17 structurally related mammalian enzymes (reviewed in refs. 13–16). Only three PARPs have demonstrable PARP activity: PARP1, PARP2 (PARP1/2), and tankyrase 1. Other

proteins containing all of the structural features of functional PARP have been identified (i.e., PARP3, PARP4), but their ability to form poly(ADP-ribose), abbreviated as PAR, chains remains to be shown. PARP1 is the most abundant PARP and is normally associated with chromatin. Binding of non-PARylated PARP1 to chromatin increases chromatin compaction (14). PARP1 activation and PARylation are associated with transcription activation, in some cases in association with DNA cleavage by topoisomerase II beta (17, 18). PARP1/2 are both strongly activated by DNA damage, including DNA single- and double-strand breaks. The N-terminal region of PARP1 contains zinc-binding domains that bind DNA (13–15). PARP1/2 complement each other. Single PARP1 or PARP2 knockout mice are viable, although they are hypersensitive to radiation and DNA damaging drugs, whereas the double PARP1/2 knockout is an early embryonic lethal (15).

PAR was discovered approximately 50 years ago by Chambon and coworkers (19). Its formation is catalyzed by PARP1/2 in several steps (20). PARP is activated upon DNA binding by intramolecular folding and dimerization. The PARP homodimer then catalyzes the transfer of an ADP-ribose moiety from NAD⁺ to a lysine (or glutamate) residue of an acceptor protein, followed by the sequential addition of multiple ADP-ribose units to the preceding ones, forming linear and branched PAR chains containing

up to 200 units. In response to DNA damage, PARP1 itself is the main PARylation acceptor (automodification), and more than 90% of PAR is found on PARP1 (13). PARylation is a reversible reaction. Poly(ADP-ribose) glycosylase (PARG) acts as an endo- and exo-glycosidase and hydrolyzes the glycosidic linkages between ADP-ribose units of PAR, producing free ADP-ribose, and ADP-ribosyl protein lyase hydrolyzes the remaining protein-proximal ADP-ribose monomers (14). PAR hydrolysis via endo- and exo-glycosidase activities would restore PARP1's ability to identify DNA single-strand breaks (SSB) and activate a new DNA damage response, if necessary. Another consequence of PAR hydrolysis would be the production of high cellular levels of ADP-ribose, which in turn, may be hydrolyzed to phospho-ribose and AMP, increasing the AMP to ATP ratio and inducing an autophagic state (16). PAR

functions as a docking polymer for a variety of chromatin and DNA repair proteins, including histones, XRCC1, Ku, DNA-dependent protein kinases (DNA-PKc), condensins, and DNA topoisomerases (13, 15). PARylation acts as an early DDR post-translational modification that recruits repair proteins to the DNA damage site, including scaffolding protein XRCC1, a factor involved in base excision repair, DNA ligase III, and DNA polymerase β (see Fig. 2 for details; refs. 21–23). PARP, along with XRCC1, has also been implicated in the alternative nonhomologous end-joining (NHEJ) pathway of DSB repair (24, 25), and in DSB repair during spermatogenesis (26). Because of PARP's involvement in SSB repair, its inhibition has been proposed to lead to DSB formation.

Because BRCA1 and BRCA2 are defective in many cancers and necessary for DSB repair, inhibition of PARP1 in such cancer cells results in enhanced and selective killing (Fig. 2; refs. 27, 28). PARP inhibitors provide a novel way to treat BRCA-deficient cells in combination with chemotherapy and radiation, while sparing normal cells. Therefore, PARP is a valuable target for pharmacologic strategies (16, 29), and PARP inhibitors are now used in an increasing number of clinical trials (30).





Fig. 2. Schematic of PARP1's role in SSB repair. Upon detection of the DNA SSB lesion, PARP1 is activated and, in turn, synthesizes PAR polymers attached to itself and other acceptor proteins at the DNA lesion site (histone H1, other core histones, TOP1, etc.). These accrued posttranslational modifications favor the recruitment of other factors involved in DNA repair, especially those involved in the base excision repair (XRCC1, Tdp1, Ligase III, Pol β). If left unrepaired, for example, with PARP inhibition (red arrows), SSBs can lead to DSBs and γ H2AX foci formation. DSB repair requires BRCA1/2, proteins that are deficient in many cancers, including breast and ovarian.

Biosampling for γ H2AX and PAR Measurements

Ideally, the efficiency of a cancer therapy based on PARP inhibitors would be to measure both γ H2AX formation and PARP activity (i.e., PAR levels) in tumor biopsies (31, 32). However, issues of tumor accessibility, patient discomfort, and risk of infection prevent the widespread acquisition of tumor biopsies. Even if tumor biopsies were routinely available, other confounding issues might prevent straightforward interpretations of any results obtained from them. One issue stems from tumor heterogeneity (33), owing to differing replicating fractions and to vascularization anomalies affecting oxygen, nutrient, and drug delivery. Therefore, γ H2AX and PAR amounts may differ among multiple biopsies of the same tumor. Perhaps comparability of tumor responses might be improved by characterizing the extent of hypoxia and/or replication status along with γ H2AX levels in the cells being analyzed. Another confounding factor involves decisions to repeatedly sample the same nodule with consequent influence on the biomarker, or to sample a different nodule with questions of biological variability. One unique solution for obtaining multiple tumor samples that may yield useful information involves the isolation of circulating tumor cells (CTC) from the patient's blood (34), which is much more feasible than obtaining multiple tumor biopsies (35).

Because of these confounding issues with tumor biopsies, procedures to assess other, more accessible, surrogate patient tissues by less invasive means are being developed. The advantages and inconveniences of different tumor and surrogate tissues for γ H2AX detection are summarized in Fig. 3. The most suitable surrogate sample should be

Fig. 3. Human biopsy analysis. A, biopsy sites and methods used to analyze γ H2AX levels and PARP activities. B, biosample accessibility, state of DNA metabolism, and expected γ H2AX response. Note that the most accessible tissues for analysis are not always the most appropriate for measuring a drug response. For example, lymphocyte and oral cells, two highly differentiated cell types, may exhibit poor responses to chemotherapeutic drugs targeting DNA replication. In contrast, tumor cells, although clearly appropriate, are often poorly accessible, particularly for repetitive sampling. γ H2AX formation is independent of the cell-cycle state, occurring in cancer cells as well as in lymphocytes and oral cells after irradiation (37, 40, 70).



	Biosampling	DNA metabolism	Expected γ H2AX response with drugs	Expected γ H2AX response after IR
Tumor cells	+/-	+++++	+++++	+++++
CTCs	+++	+++++	+++++	+++++
Lymphocytes	++++	-	+	+++++
Skin biopsies	++	++++	++++	+++++
Plucked hairs	+++++	++	+++	++++
Oral cells	+++++	-	+/-	++

selected on the basis of the type of treatment received by the patient (see Fig. 3 for details).

Circulating blood lymphocytes and leukocytes are terminally differentiated (postmitotic) cells that respond consistently to irradiation with robust γ H2AX signals proportional to the radiation doses (36, 37); however, their use for assessing the action of anticancer drugs that interfere with DNA replication is problematic. In contrast, PAR measurements in lymphocytes can be made, and reduction in PAR levels can readily be observed following PARP inhibitor treatment in most patients (32).

Buccal cells can be collected very simply by swabbing the patient's inner cheeks, but collection of numbers of living cells sufficient for testing can be problematic. In addition, buccal cells, a type of terminally differentiated stratified squamous epithelium, can present a challenge in procedures involving cell lysis or permeabilization when using microscopy (38). In addition, buccal cells seem to have very high background levels of DNA damage (38, 39). Nevertheless, significantly increased γ H2AX formation was detected in patients undergoing routine dental radiographic examination, indicating that the use of γ H2AX in oral cells could serve as sensitive indicators of low-dose radiation exposure (40).

Another minimally invasive procedure consists of collecting hairs plucked from the scalp or eyebrows. A sub-

stantial portion of the hair bulb may stay attached to the hair shaft and contain dividing and stem cells (41), which can be used to measure γ H2AX levels and PARP inhibition (42). However, considering that a patient's hair follicles are in various stages of growth, plucked hairs containing replicating cells (i.e., in anagen phase) should be favored for chemotherapy assessments.⁵

Finally, because skin contains proliferative cells, such a tissue would be valuable to determine DNA damage (i.e., γ H2AX levels) following treatment with drugs that interfere with DNA replication. However, like tumor biopsies, multiple sampling would result in patient discomfort and may lead to possible complications, such as infections.

A major issue is the relationship of the responses of γ H2AX and PAR in surrogate tissues versus the tumors, and the relationship between those responses and the treatment outcome. Although making these comparisons for solid tumors is challenging because of the difficulty of biopsy collection, blood malignancies may present a more accessible situation. γ H2AX and PAR levels could be directly measured in malignant cells collected in blood

⁵ C.E. Redon and W.M. Bonner, unpublished observations.

samples. However, the use of γ H2AX in surrogate tissues can help determine both a drug's genotoxicity and its pharmacokinetics *in vivo*. Such comparisons could be extended to multiple drug combinations.

Assays for γ H2AX and PAR Detection

Several immunologic techniques can be used to quantify γ H2AX and PAR (PARP activity is deduced from

changes in PAR levels). They may consist of counting antigen intensities in cells and tissues [i.e., microscopy and fluorescence-activated cell sorting (FACS)] or quantifying their overall levels [i.e., Western blotting and enzyme-linked immunosorbent assay (ELISA)]; Figs. 3 and 4; refs. 4, 32, 42–44]. In recent years, the Pharmacodynamic Assay Development and Implementation Section (PADIS; NCI), in collaboration with Abbott Laboratories and the National Clinical Target Validation Laboratory at the




Fig. 4. γ H2AX and PAR detection. A, γ H2AX detection in lymphocytes; B, tumor needle biopsies; and C, plucked hairs from patients undergoing chemotherapy. C, the white box in the lower panel marks the region of active γ H2AX formation in a typical plucked hair. Green, γ H2AX; red, DNA. D, pharmacodynamic assay developed at the NCI to measure PAR as a biomarker for PARP inhibition in both tumor biopsies and PBMCs.

Table 1. Anticancer drugs that produce γ H2AX

Drugs	Mechanism of induction	References
Bleomycin	Direct DSB, iron-mediated oxidative cleavage	70
Camptothecins and indenoisoquinolines	Indirect: conversion of SSB to DSB by replication	12, 71, 72
Doxorubicin, etoposide, mitoxantrone, irinotecan	Direct DSB by trapping topoisomerase II cleavage complexes. Also indirect: ROS formation	73–75
Cytarabine, gemcitabine, hydroxyurea	Indirect: replication fork collapse (chain termination; deoxyribonucleotide pool depletion)	76–78
Cisplatin, temozolomide, aminoflavone, trabectedin	Indirect: DNA alkylation	79–82
Imatinib (gleevec)	Indirect: apoptosis induced by Kit/PDGF tyrosine kinase inhibition	83
5-azacytidine SAHA (viconostat)	Indirect: epigenetic modifications	84, 85
PARP and DNA-PK inhibitors (see ref. 30 for PARPi)	Indirect: interference with SSB and DSB repair	27, 86, 87
SJG-136	Indirect: DNA alkylation	53
Tirapazamine	Indirect: ROS production in hypoxic cells	73, 88
TRAIL	Indirect: Death receptors-mediated activation of DNA-PK	89, 90
UCN-01 and AZD7762	Indirect: Checkpoint inhibitor potentiating IR- and replication-induced DNA damage by topoisomerase I inhibitors, cytarabine and gemcitabine.	77, 91, 92

NOTE: All the anticancer drugs listed can also induce delayed γ H2AX activation by apoptosis.
Abbreviations: PDGF, platelet-derived growth factor; IR, ionizing radiation.

NCI, has developed an ELISA for PAR levels in tissue samples. With this assay, the *in vivo* effect of a PARP inhibitor was measured (Fig. 4; ref. 45). Finally, a method using an electrochemoluminescent detection system (an assay derived from the Meso Scale Discovery Technology or MSD assay) was recently reported. This ELISA-based methodology allows γ H2AX measurements in tumors after irradiation (46).

Although each of these methods can be effective and gives important information, they are not currently usable for high-throughput screening and still involve extensive human labor. Recently, Garty and colleagues reported the development of a high-throughput screening system as a radiation biodosimetry tool for radiation triage, and using a γ H2AX immunofluorescence assay, called the RABIT (Rapid Automated Biodosimetry Tool), which would allow the screening of 6,500 samples a day (47).

Both γ H2AX and PAR assays are specific and very sensitive. A single DSB (corresponding to one γ H2AX focus) can be visualized by microscopy, and responses to radiation doses as low as 1.2 mGy and 100 mGy can be detected by microscopy and FACS, respectively (4). The ELISA-based measurements of PAR has a lower limit of detection < 6 pg/ml with a dynamic range of 7.8 to 1,000 pg/ml.⁶

⁶ J. Ji, personal communications.

γ H2AX in Clinical Oncology

γ H2AX in cancer chemotherapy

Many cancer therapies rely on agents that preferentially kill tumor cells by generating DNA damage (48), including DSBs (Table 1; reviewed in ref. 4). Methods to detect DSB levels directly in patient material cells could be invaluable for optimizing treatment, particularly with chemotherapeutic agents whose efficacies could vary among individuals with different genetic backgrounds. Because biosampling tumors is not always possible, alternate surrogate tissues have been developed depending on availability (i.e., hair sampling could be problematic in patients with alopecia due to prior treatments) or on the agent (i.e., terminally differentiated surrogate cells may not respond well when using a drug interfering with replication). For this reason, biosampling of several tissues would be preferable during drug development.

Clinical studies with different methods of γ H2AX analysis are described in Table 2. In a phase I clinical study for patients with refractory leukemia, flow cytometry revealed increased γ H2AX levels in circulating leukemia cells in 12 of 13 patients after treatment with a novel deoxyadenosine analog (clofarabine), combined with an alkylating agent (cyclophosphamide; ref. 49). The same group also reported that the peripheral blood mononuclear cells (PBMC) from 23 patients with myelodysplasia (MDS) and high-risk acute myelogenous leukemia (AML) exhibited increased γ H2AX levels after sequential administration of a DNA methyltransferase

inhibitor (5-azacytidine) and a histone deacetylase inhibitor (ectinostat; ref. 50). However, the latter data did not show any correlation between the patients' responses and γ H2AX induction. Another clinical study showed the ability of the minor groove binding agent SJG-136 to increase γ H2AX levels in both the patients' lymphocytes and tumor biopsies (51). Fong and colleagues described the use of plucked-eyebrow hairs to follow patients treated with a

PARP inhibitor. Their study showed a correlation between γ H2AX levels and PARP inhibition (42). Finally, a trial testing the combination of a farnesyltransferase inhibitor (tipifarnib) and a topoisomerase II inhibitor (etoposide) for individuals diagnosed with AML used a γ H2AX assay on AML marrow blasts (52). This drug combination showed genotoxicity by both the use of γ H2AX and increased subdiploid DNA content.

Table 2. Some published examples of clinical studies using γ H2AX detection *in vivo*

Application	Tissue	Study details	Method	References
A. γH2AX used in clinical trials				
Chemotherapy	PBMCs	Phase 1 study of clofarabine followed by cyclophosphamide for adults with refractory acute leukemias	FACS	49
Chemotherapy	PBMCs and/or tumor biopsies	A phase 1 study of SJG-136	Microscopy	51
Chemotherapy	AML marrow blasts	A phase 1 study of the combination of tipifarnib and etoposide for patients with AML	FACS	52
Chemotherapy	PBMCs	A phase 1 study of 5-azacytidine and entinostat for patients with MDS, chronic myelomonocytic leukemia, and AML	Immunoblotting	50
Chemotherapy	Plucked eyebrows	A phase 1 study of olaparib	Microscopy	42
B. γH2AX used for radiation biodosimetry				
Computed tomography	PBMCs	DNA damage measured after multidetector row CT	Microscopy	93
Radiotherapy	PBMCs	DNA damage measured in cancer patients after local radiotherapy to different sites of the body	Microscopy	54
Radiotherapy	Skin	DNA damage measured in prostate cancer patients undergoing radiotherapy with curative intent	Microscopy	53
X-ray examination	PBMCs	DNA damage measured after cardiac catheterization	Microscopy	94
X-ray examination	PBMCs	DNA damage measured after coronary CT angiographic procedure	Microscopy	95
X-ray examination	PBMCs	DNA damage measured after angiographic procedure	Microscopy	56
X-ray examination	PBMCs	DNA damage measured after percutaneous transluminal angioplasty	Microscopy	55
C. γH2AX used for diagnostics				
Diagnosis	Tumor biopsies	Diagnosis of metastatic renal cell carcinoma	Microscopy	64
Diagnosis	Tissue biopsies	Monitoring DNA damage in ulcerative colitis	Immunoblotting	96
Radiosensitivity diagnosis	T cells and lymphoblastoid cell lines and/or PBMCs	Confirmation of radiosensitive ataxia telangiectasia patients	FACS	58

NOTE: Studies are listed in three separate groups: A, γ H2AX used in clinical trials; B, γ H2AX used for radiation biodosimetry; C, γ H2AX used for diagnostics.
Abbreviation: CT, computed tomography.

γ H2AX use in radiotherapy

In contrast to chemotherapeutic agents, ionizing radiation can induce DSBs regardless of cell-cycle phase, thus, radiation-induced γ H2AX could, in principle, be detectable in all tissues or cells including G0 lymphocytes and oral cells. Counting γ H2AX foci has been used successfully for biodosimetry (36, 53–56). After exposure, γ H2AX foci levels in patient lymphocytes and skin biopsies exhibited a good correlation with the radiation doses.

Additionally, γ H2AX measurements might provide information useful in improving patient outcome during radiotherapy. Although treatment with radiation is commonly used, some patients develop severe, possibly lethal, side effects. Patients with DSB repair deficiencies, such as ataxia telangiectasia caused by mutation in the *ATM* gene, are highly sensitive to radiation exposure (57). Thus, γ H2AX measurements may allow the identification of hypersensitive individuals in order to improve the efficacy of radiotherapy and to avoid therapeutic accidents (58, 59).

It is important to note that therapeutic and accidental radiation exposures are rarely uniform and, therefore, can result in subpopulations of lymphocytes with different exposures, a phenomenon that should be taken into account for dose estimation. Moreover, because hypoxic microenvironments in solid tumors result in the decrease of the number of DSBs by radiotherapy (33), variations in γ H2AX levels could occur in different tumor samples for a same irradiation dose.

γ H2AX use in diagnostics

γ H2AX also has potential uses in medical diagnosis (Table 2). Because both precancerous and cancerous cells were found to exhibit increased genomic instability (60, 61), a biomarker able to identify such cells in patients would be a useful diagnostic tool. In fact, it was shown that γ H2AX can be used as a biomarker for cancer because tumor biopsies show increased γ H2AX levels (62, 63). γ H2AX measurements also established high levels of DNA damage in colon biopsies of patients with ulcerative colitis, an inflammation disease linked to higher risk for colorectal cancer. A recent study showed the dependability and accuracy of γ H2AX in the diagnosis of metastatic renal cell carcinoma (64).

PAR in Clinical Oncology

Several PARP inhibitors (at least eight: iniparib/BSI-201, olaparib/AZ2281, veliparib/ABT-888, AG014699, MK-4827, CEP-8933/CEP-9722, INO-1001, GPI 21016) have been developed and are now used alone or in combination with other drugs in several clinical trials (30). Because of the high homology between PARP1 and PARP2, most PARP inhibitors can target both enzymes, and future studies will probably look for specific inhibitors (65). The first report of a PARP clinical trial using the drug BSI-201 alone in a phase I study came out in 2008 (66). Since that time, this drug has been used in

more phase I trials in combination with other drugs, in phase II clinical studies, and is now entering phase III study (67, 68). Olaparib and veliparib/ABT-888 are also undergoing clinical development in several phase I and phase II trials. The NCI conducted the first clinical pharmacodynamic trial (phase 0) of ABT-888, an orally available small-molecule inhibitor of PARP, in patients with advanced malignancies. PAR levels in tumor biopsies and PBMCs were measured by using a validated ELISA (32). The data obtained could be used to guide the design of several phase I combination studies of veliparib, including ongoing trials with topotecan and cyclophosphamide, each of which include measurement of PAR as a pharmacodynamic endpoint.

In these studies, it is crucial to take measurements at the *in vivo* cellular level, both for the drug effect on PARP activity (i.e., PAR levels) and the consequences of this inhibition (i.e., DNA damage induction via γ H2AX detection). Recent years have seen the development of ELISA for PAR detection that can now replace the less reliable immunoassays (Fig. 4D; refs. 32, 69). Thus, PAR levels can be measured in both PBMCs (mostly lymphocytes) and tumor biopsies. These measurements show a significant correlation between the effects of the PARP inhibitor in PBMCs and the tumor samples (32), raising the possibility that blood samples could be used as tumor surrogates to follow PARP inhibition.

Conclusions

The availability of biomarkers of DNA damage offers the opportunity to evaluate clinical cancer samples and to determine their DDR status prior to and during therapy. The systematic use of γ H2AX and PAR in tumor samples, in the absence of treatment, may identify groups of patients with a particular prognosis. γ H2AX and PAR may also be useful in evaluating the activity of novel drugs in tumor and normal tissues in response to treatment, which could accelerate the drug triage process, for example aiding go or no-go decisions, or lead compound selection. Hopefully, incorporation of such biomarkers will eventually eliminate a substantial fraction of drugs that fail in phase II-III clinical trials.

The related question is if the role of γ H2AX and PAR could be developed beyond being a tool for clinical trials, to that of a marker with utility in cancer therapy. Following treatment, assessment of γ H2AX and PAR could allow identification of patients who had insufficient evidence of DNA damage, so that different drugs could be selected, or strategies to increase drug exposure, change schedule, or improve activity could be identified. It is likely that γ H2AX and PAR only represent the first generation of DDR biomarkers, and that more sensitive and convenient biomarkers will be developed in the near future. Ultimately, one of the major challenges remains in tumor sampling, and also in developing noninvasive detection procedures. In the meantime, it will be interesting to take advantage of analyses in CTCs.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Dr. Susan Bates, Christine Bryla, Robin Frye, Kathryn Compton, Merrill Goldsmith, Gareth Peters, and William Yutzy for their help in the biomarker studies.

Grant Support

This research was supported [in part] by the Developmental Therapeutics Program in the Division of Cancer Treatment and Diagnosis of the National Cancer Institute and by the Center for Cancer Research, Intramural Program, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.

Received 06/01/2010; revised 07/15/2010; accepted 07/29/2010; published OnlineFirst 09/07/2010.

References

- Kummar S, Kinders R, Rubinstein L, et al. Compressing drug development timelines in oncology using phase '0' trials. *Nat Rev Cancer* 2007;7:131–9.
- Frank R, Hargreaves R. Clinical biomarkers in drug discovery and development. *Nat Rev Drug Discov* 2003;2:566–80.
- Redon C, Pilch D, Rogakou E, Sedelnikova O, Newrock K, Bonner W. Histone H2A variants H2AX and H2AZ. *Curr Opin Genet Dev* 2002;12:162–9.
- Bonner WM, Redon CE, Dickey JS, et al. γ H2AX and cancer. *Nat Rev Cancer* 2008;8:957–67.
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* 1998;273:5858–68.
- Celeste A, Petersen S, Romanienko PJ, et al. Genomic instability in mice lacking histone H2AX. *Science* 2002;296:922–7.
- Cha H, Lowe JM, Li H, et al. Wip1 directly dephosphorylates γ -H2AX and attenuates the DNA damage response. *Cancer Res* 2010;70:4112–22.
- Moon SH, Nguyen TA, Darlington Y, Lu X, Donehower LA. Dephosphorylation of γ H2AX by WIP1: An important homeostatic regulatory event in DNA repair and cell cycle control. *Cell Cycle* 2010;9. Epub 2010 Jun 12.
- Douglas P, Zhong J, Ye R, Moorhead GB, Xu X, Lees-Miller SP. Protein phosphatase 6 interacts with the DNA-dependent protein kinase catalytic subunit and dephosphorylates γ -H2AX. *Mol Cell Biol* 2010;30:1368–81.
- Nakada S, Chen GI, Gingras AC, Durocher D. PP4 is a γ H2AX phosphatase required for recovery from the DNA damage checkpoint. *EMBO Rep* 2008;9:1019–26.
- Chowdhury D, Keogh MC, Ishii H, Peterson CL, Buratowski S, Lieberman J. γ -H2AX dephosphorylation by protein phosphatase 2A facilitates DNA double-strand break repair. *Mol Cell* 2005;20:801–9.
- Pommier Y. Topoisomerase I inhibitors: camptothecins and beyond. *Nat Rev Cancer* 2006;6:789–802.
- D'Amours D, Desnoyers S, D'Silva I, Poirier GG. Poly(ADP-ribosyl)ation reactions in the regulation of nuclear functions. *Biochem J* 1999;342:249–68.
- Kim MY, Zhang T, Kraus WL. Poly(ADP-ribosyl)ation by PARP-1: 'PAR-laying' NAD⁺ into a nuclear signal. *Genes Dev* 2005;19:1951–67.
- Schreiber V, Dantzer F, Ame JC, de Murcia G. Poly(ADP-ribose): novel functions for an old molecule. *Nat Rev Mol Cell Biol* 2006;7:517–28.
- Rouleau M, Patel A, Hendzel MJ, Kaufmann SH, Poirier GG. PARP inhibition: PARP1 and beyond. *Nat Rev Cancer* 2010;10:293–301.
- Ju BG, Lunyak VV, Perissi V, et al. A topoisomerase II β -mediated dsDNA break required for regulated transcription. *Science* 2006;312:1798–802.
- Lis JT, Kraus WL. Promoter cleavage: a topolI β and PARP-1 collaboration. *Cell* 2006;125:1225–7.
- Chambon P, Weill JD, Mandel P. Nicotinamide mononucleotide activation of new DNA-dependent polyadenylic acid synthesizing nuclear enzyme. *Biochem Biophys Res Commun* 1963;11:39–43.
- Altmeyer M, Messner S, Hassa PO, Fey M, Hottiger MO. Molecular mechanism of poly(ADP-ribosyl)ation by PARP1 and identification of lysine residues as ADP-ribose acceptor sites. *Nucleic Acids Res* 2009;37:3723–38.
- Masson M, Niedergang C, Schreiber V, Muller S, Menissier-de Murcia J, de Murcia G. XRCC1 is specifically associated with poly(ADP-ribose) polymerase and negatively regulates its activity following DNA damage. *Mol Cell Biol* 1998;18:3563–71.
- Caldecott KW. XRCC1 and DNA strand break repair. *DNA Repair (Amst)* 2003;2:955–69.
- El-Khamisy SF, Masutani M, Suzuki H, Caldecott KW. A requirement for PARP-1 for the assembly or stability of XRCC1 nuclear foci at sites of oxidative DNA damage. *Nucleic Acids Res* 2003;31:5526–33.
- Audebert M, Salles B, Calsou P. Involvement of poly(ADP-ribose) polymerase-1 and XRCC1/DNA ligase III in an alternative route for DNA double-strand breaks rejoining. *J Biol Chem* 2004;279:55117–26.
- Iliakis G. Backup pathways of NHEJ in cells of higher eukaryotes: cell cycle dependence. *Radiother Oncol* 2009;92:310–5.
- Ahmed EA, de Boer P, Philippens ME, Kal HB, de Rooij DG. Parp1-XRCC1 and the repair of DNA double strand breaks in mouse round spermatids. *Mutat Res* 2010;683:84–90.
- Bryant HE, Schultz N, Thomas HD, et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature* 2005;434:913–7.
- Farmer H, McCabe N, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005;434:917–21.
- Drew Y, Plummer R. PARP inhibitors in cancer therapy: two modes of attack on the cancer cell widening the clinical applications. *Drug Resist Updat* 2009;12:153–6.
- Annunziata CM, O'Shaughnessy J. Poly(ADP-Ribose) polymerase as a novel therapeutic target in cancer. *Clin Cancer Res* 2010;16:4517–26.
- Oka K, Tanaka T, Enoki T, et al. DNA damage signaling is activated during cancer progression in human colorectal carcinoma. *Cancer Biol Ther* 2010;9:246–52.
- Kummar S, Kinders R, Gutierrez ME, et al. Phase 0 clinical trial of the poly (ADP-ribose) polymerase inhibitor ABT-888 in patients with advanced malignancies. *J Clin Oncol* 2009;27:2705–11.
- Chan N, Bristow RG. "Contextual" synthetic lethality and/or loss of heterozygosity: tumor hypoxia and modification of DNA repair. *Clin Cancer Res* 2010;16:4553–60.
- Wang LH, Pfister TD, Parchment RE, et al. Monitoring drug-induced γ H2AX as a pharmacodynamic biomarker in individual circulating tumor cells. *Clin Cancer Res* 2010;16:1073–84.
- Maheswaran S, Haber DA. Circulating tumor cells: a window into cancer biology and metastasis. *Curr Opin Genet Dev* 2010;20:96–9.
- Lobrich M, Rief N, Kuhne M, et al. *In vivo* formation and repair of DNA double-strand breaks after computed tomography examinations. *Proc Natl Acad Sci U S A* 2005;102:8984–9.
- Redon CE, Dickey JS, Bonner WM, Sedelnikova OA. γ -H2AX as a biomarker of DNA damage induced by ionizing radiation in human peripheral blood lymphocytes and artificial skin. *Adv Space Res* 2009;43:1171–8.
- Szeto YT, Benzie IF, Collins AR, et al. A buccal cell model comet assay: development and evaluation for human biomonitoring and nutritional studies. *Mutat Res* 2005;578:371–81.

39. Rothkamm K, Horn S. γ -H2AX as protein biomarker for radiation exposure. *Ann Ist Super Sanita* 2009;45:265–71.
40. Yoon AJ, Shen J, Wu HC, et al. Expression of activated checkpoint kinase 2 and histone 2AX in exfoliative oral cells after exposure to ionizing radiation. *Radiat Res* 2009;171:771–5.
41. Gho CG, Braun JE, Tilli CM, Neumann HA, Ramaekers FC. Human follicular stem cells: their presence in plucked hair and follicular cell culture. *Br J Dermatol* 2004;150:860–8.
42. Fong PC, Boss DS, Yap TA, et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med* 2009;361:123–34.
43. Kunzmann A, Liu D, Annett K, et al. Flow-cytometric assessment of cellular poly(ADP-ribose)ylation capacity in peripheral blood lymphocytes. *Immun Ageing* 2006;3:8.
44. Yang SX, Kummar S, Steinberg SM, et al. Immunohistochemical detection of poly(ADP-ribose) polymerase inhibition by ABT-888 in patients with refractory solid tumors and lymphomas. *Cancer Biol Ther* 2009;8:2004–9.
45. Kinders RJ, Hollingshead M, Khin S, et al. Preclinical modeling of a phase 0 clinical trial: qualification of a pharmacodynamic assay of poly(ADP-ribose) polymerase in tumor biopsies of mouse xenografts. *Clin Cancer Res* 2008;14:6877–85.
46. Avondoglio D, Scott T, Kil WJ, Sproull M, Tofilon PJ, Camphausen K. High throughput evaluation of γ -H2AX. *Radiat Oncol* 2009;4:31.
47. Garty G, Chen Y, Salerno A, et al. The RABIT: a rapid automated biodosimetry tool for radiological triage. *Health Phys* 2010;98:209–17.
48. Plummer R. Perspective on the pipeline of drugs being developed with modulation of DNA damage as a target. *Clin Cancer Res* 2010;16:4527–31.
49. Karp JE, Ricklis RM, Balakrishnan K, et al. A phase 1 clinical-laboratory study of clofarabine followed by cyclophosphamide for adults with refractory acute leukemias. *Blood* 2007;110:1762–9.
50. Fandy TE, Herman JG, Kerns P, et al. Early epigenetic changes and DNA damage do not predict clinical response in an overlapping schedule of 5-azacytidine and entinostat in patients with myeloid malignancies. *Blood* 2009;114:2764–73.
51. Hochhauser D, Meyer T, Spanswick VJ, et al. Phase I study of sequence-selective minor groove DNA binding agent SJG-136 in patients with advanced solid tumors. *Clin Cancer Res* 2009;15:2140–7.
52. Karp JE, Flatten K, Feldman EJ, et al. Active oral regimen for elderly adults with newly diagnosed acute myelogenous leukemia: a pre-clinical and phase 1 trial of the farnesyltransferase inhibitor tipifarnib (R115777, Zarnestra) combined with etoposide. *Blood* 2009;113:4841–52.
53. Qvarnstrom OF, Simonsson M, Johansson KA, Nyman J, Turesson I. DNA double strand break quantification in skin biopsies. *Radiother Oncol* 2004;72:311–7.
54. Sak A, Grehl S, Erichsen P, et al. γ -H2AX foci formation in peripheral blood lymphocytes of tumor patients after local radiotherapy to different sites of the body: dependence on the dose-distribution, irradiated site and time from start of treatment. *Int J Radiat Biol* 2007;83:639–52.
55. Geisel D, Heverhagen JT, Kalinowski M, Wagner HJ. DNA double-strand breaks after percutaneous transluminal angioplasty. *Radiology* 2008;248:852–9.
56. Kuefner MA, Grudzinski S, Schwab SA, et al. DNA double-strand breaks and their repair in blood lymphocytes of patients undergoing angiographic procedures. *Invest Radiol* 2009;44:440–6.
57. Taylor AM, Byrd PJ. Molecular pathology of ataxia telangiectasia. *J Clin Pathol* 2005;58:1009–15.
58. Porcedda P, Turinetto V, Brusco A, et al. A rapid flow cytometry test based on histone H2AX phosphorylation for the sensitive and specific diagnosis of ataxia telangiectasia. *Cytometry A* 2008;73:508–16.
59. Porcedda P, Turinetto V, Orlando L, et al. Two-tier analysis of histone H2AX phosphorylation allows the identification of Ataxia Telangiectasia heterozygotes. *Radiother Oncol* 2009;92:133–7.
60. Bartkova J, Horejsi Z, Koed K, et al. DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature* 2005;434:864–70.
61. Gorgoulis VG, Vassiliou LV, Karakaidos P, et al. Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* 2005;434:907–13.
62. Sedelnikova OA, Bonner WM. γ H2AX in cancer cells: a potential biomarker for cancer diagnostics, prediction and recurrence. *Cell Cycle* 2006;5:2909–13.
63. Yu T, MacPhail SH, Banath JP, Klovov D, Olive PL. Endogenous expression of phosphorylated histone H2AX in tumors in relation to DNA double-strand breaks and genomic instability. *DNA Repair (Amst)* 2006;5:935–46.
64. Wasco MJ, Pu RT. Utility of antiphosphorylated H2AX antibody (γ -H2AX) in diagnosing metastatic renal cell carcinoma. *Appl Immunohistochem Mol Morphol* 2008;16:349–56.
65. Ishida J, Yamamoto H, Kido Y, et al. Discovery of potent and selective PARP-1 and PARP-2 inhibitors: SBDD analysis via a combination of X-ray structural study and homology modeling. *Bioorg Med Chem* 2006;14:1378–90.
66. Kopetz SMM, Mok I, et al. First in human phase I study of BSI-201, a small molecule inhibitor of poly ADP-ribose (PARP) in subjects with advanced solid tumors [abstract]. *J Clin Oncol* 2008;26:abstr 3577.
67. Mahany JJ, Lewis N, Heath EI, et al. A phase IB study evaluating BSI-201 in combination with chemotherapy in subjects with advanced solid tumors [abstract]. *J Clin Oncol* 2008;26:abstr 3579.
68. O'Shaughnessy J, Osborne C, Pippen J, et al. Efficacy of BSI-201, a poly(ADP-ribose) polymerase-I (PARP1) inhibitor, in combination with gemcitabine/carboplatin (G/C) in patients with metastatic triple-negative breast cancer (TNBC): results of a randomized phase II trial [abstract]. *J Clin Oncol* 2009;27:abstr 3.
69. Liu X, Palma J, Kinders R, et al. An enzyme-linked immunosorbent poly(ADP-ribose) polymerase biomarker assay for clinical trials of PARP inhibitors. *Anal Biochem* 2008;381:240–7.
70. Banath JP, Macphail SH, Olive PL. Radiation sensitivity, H2AX phosphorylation, and kinetics of repair of DNA strand breaks in irradiated cervical cancer cell lines. *Cancer Res* 2004;64:7144–9.
71. Furuta T, Takemura H, Liao ZY, et al. Phosphorylation of histone H2AX and activation of Mre11, Rad50, and Nbs1 in response to replication-dependent DNA double-strand breaks induced by mammalian DNA topoisomerase I cleavage complexes. *J Biol Chem* 2003;278:20303–12.
72. Antony S, Agama KK, Miao ZH, et al. Novel indenoisoquinolines NSC 725776 and NSC 724998 produce persistent topoisomerase I cleavage complexes and overcome multidrug resistance. *Cancer Res* 2007;67:10397–405.
73. Olive PL, Banath JP, Sinnott LT. Phosphorylated histone H2AX in spheroids, tumors, and tissues of mice exposed to etoposide and 3-amino-1,2,4-benzotriazine-1,3-dioxide. *Cancer Res* 2004;64:5363–9.
74. Kurz EU, Lees-Miller SP. DNA damage-induced activation of ATM and ATM-dependent signaling pathways. *DNA Repair (Amst)* 2004;3:889–900.
75. Rao VA, Agama K, Holbeck S, Pommier Y. Batracylin (NSC 320846), a dual inhibitor of DNA topoisomerases I and II induces histone γ -H2AX as a biomarker of DNA damage. *Cancer Res* 2007;67:9971–9.
76. Krynetskaia N, Xie H, Vucetic S, Obradovic Z, Krynetskiy E. High mobility group protein B1 is an activator of apoptotic response to antimetabolite drugs. *Mol Pharmacol* 2008;73:260–9.
77. Ewald B, Sampath D, Plunkett W. H2AX phosphorylation marks gemcitabine-induced stalled replication forks and their collapse upon S-phase checkpoint abrogation. *Mol Cancer Ther* 2007;6:1239–48.
78. Kurose A, Tanaka T, Huang X, Traganos F, Dai W, Darzynkiewicz Z. Effects of hydroxyurea and aphidicolin on phosphorylation of ataxia telangiectasia mutated on Ser 1981 and histone H2AX on Ser 139 in relation to cell cycle phase and induction of apoptosis. *Cytometry A* 2006;69:212–21.
79. Pabla N, Huang S, Mi QS, Daniel R, Dong Z. ATR-Chk2 signaling in p53 activation and DNA damage response during cisplatin-induced apoptosis. *J Biol Chem* 2008;283:6572–83.

80. Mirzoeva OK, Kawaguchi T, Pieper RO. The Mre11/Rad50/Nbs1 complex interacts with the mismatch repair system and contributes to temozolomide-induced G2 arrest and cytotoxicity. *Mol Cancer Ther* 2006;5:2757-66.
81. Meng LH, Kohlhagen G, Liao ZY, Antony S, Sausville E, Pommier Y. DNA-protein cross-links and replication-dependent histone H2AX phosphorylation induced by aminoflavone (NSC 686288), a novel anticancer agent active against human breast cancer cells. *Cancer Res* 2005;65:5337-43.
82. Guirouilh-Barbat J, Redon C, Pommier Y. Transcription-coupled DNA double-strand breaks are mediated via the nucleotide excision repair and the Mre11-Rad50-Nbs1 complex. *Mol Biol Cell* 2008;19:3969-81.
83. Liu Y, Tseng M, Perdreau SA, et al. Histone H2AX is a mediator of gastrointestinal stromal tumor cell apoptosis following treatment with imatinib mesylate. *Cancer Res* 2007;67:2685-92.
84. Kiziltepe T, Hideshima T, Catley L, et al. 5-Azacytidine, a DNA methyltransferase inhibitor, induces ATR-mediated DNA double-strand break responses, apoptosis, and synergistic cytotoxicity with doxorubicin and bortezomib against multiple myeloma cells. *Mol Cancer Ther* 2007;6:1718-27.
85. Conti C, Leo E, Eichler GS, et al. Inhibition of histone deacetylase in cancer cells slows down replication forks, activates dormant origins, and induces DNA damage. *Cancer Res* 2010;70:4470-80.
86. Albert JM, Cao C, Kim KW, et al. Inhibition of poly(ADP-ribose) polymerase enhances cell death and improves tumor growth delay in irradiated lung cancer models. *Clin Cancer Res* 2007;13:3033-42.
87. Zhao Y, Thomas HD, Batey MA, et al. Preclinical evaluation of a potent novel DNA-dependent protein kinase inhibitor NU7441. *Cancer Res* 2006;66:5354-62.
88. Evans JW, Chernikova SB, Kachnic LA, et al. Homologous recombination is the principal pathway for the repair of DNA damage induced by tirapazamine in mammalian cells. *Cancer Res* 2008;68:257-65.
89. Solier S, Pommier Y. The apoptotic ring: a novel entity with phosphorylated histones H2AX and H2B, activated DNA damage response kinases. *Cell Cycle* 2009;8:1853-9.
90. Solier S, Sordet O, Kohn KW, Pommier Y. Death receptor-induced activation of the Chk2- and histone H2AX-associated DNA damage response pathways. *Mol Cell Biol* 2009;29:68-82.
91. Furuta T, Hayward RL, Meng LH, et al. p21CDKN1A allows the repair of replication-mediated DNA double-strand breaks induced by topoisomerase I and is inactivated by the checkpoint kinase inhibitor 7-hydroxystaurosporine. *Oncogene* 2006;25:2839-49.
92. Mitchell JB, Choudhuri R, Fabre K, et al. *In vitro* and *in vivo* radiation sensitization of human tumor cells by a novel checkpoint kinase inhibitor, AZD7762. *Clin Cancer Res* 2010;16:2076-84.
93. Rothkamm K, Balroop S, Shekhdar J, Fernie P, Goh V. Leukocyte DNA damage after multi-detector row CT: a quantitative biomarker of low-level radiation exposure. *Radiology* 2007;242:244-51.
94. Beels L, Bacher K, De Wolf D, Werbrouck J, Thierens H. γ -H2AX foci as a biomarker for patient X-ray exposure in pediatric cardiac catheterization: are we underestimating radiation risks? *Circulation* 2009;120:1903-9.
95. Kuefner MA, Hinkmann FM, Alibek S, et al. Reduction of X-ray induced DNA double-strand breaks in blood lymphocytes during coronary CT angiography using high-pitch spiral data acquisition with prospective ECG-triggering. *Invest Radiol* 2010;45:182-7.
96. Risques RA, Lai LA, Brentnall TA, et al. Ulcerative colitis a disease of accelerated colon aging: evidence from telomere attrition and DNA damage. *Gastroenterology* 2008;135:410-8.
97. Lu C, Zhu F, Cho YY, et al. Cell apoptosis: requirement of H2AX in DNA ladder formation, but not for the activation of caspase-3. *Mol Cell* 2006;23:121-32.
98. Mukherjee B, Kessinger C, Kobayashi J, et al. DNA-PK phosphorylates histone H2AX during apoptotic DNA fragmentation in mammalian cells. *DNA Repair (Amst)* 2006;5:575-90.
99. Rossetto D, Truman AW, Kron SJ, Côté J. Epigenetic modification in double-strand break DNA damage signaling and repair. *Clin Cancer Res* 2010;16:4543-52.
100. Downs JA, Allard S, Jobin-Robitaille O, et al. Binding of chromatin-modifying activities to phosphorylated histone H2A at DNA damage sites. *Mol Cell* 2004;16:979-90.
101. van Attikum H, Gasser SM. ATP-dependent chromatin remodeling and DNA double-strand break repair. *Cell Cycle* 2005;4:1011-4.
102. Unal E, Arbel-Eden A, Sattler U, et al. DNA damage response pathway uses histone modification to assemble a double-strand break-specific cohesin domain. *Mol Cell* 2004;16:991-1002.
103. Xie A, Puget N, Shim I, et al. Control of sister chromatid recombination by histone H2AX. *Mol Cell* 2004;16:1017-25.
104. Bassing CH, Alt FW. H2AX may function as an anchor to hold broken chromosomal DNA ends in close proximity. *Cell Cycle* 2004;3:149-53.

Clinical Cancer Research

Histone γ H2AX and Poly(ADP-Ribose) as Clinical Pharmacodynamic Biomarkers

Christophe E. Redon, Asako J. Nakamura, Yong-Wei Zhang, et al.

Clin Cancer Res 2010;16:4532-4542. Published OnlineFirst September 7, 2010.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-10-0523](https://doi.org/10.1158/1078-0432.CCR-10-0523)

Cited articles This article cites 103 articles, 42 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/16/18/4532.full#ref-list-1>

Citing articles This article has been cited by 29 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/16/18/4532.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/16/18/4532>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.