

Imaging, Diagnosis, Prognosis

See related article by Salazar and Tabernero, p. 2019

Hypoxia-Driven Gene Expression Is an Independent Prognostic Factor in Stage II and III Colon Cancer Patients

Jeroen Dekervel¹, Daphne Hompes³, Hannah van Malenstein¹, Dusan Popovic², Xavier Sagaert⁴, Bart De Moor², Eric Van Cutsem⁵, André D'Hoore³, Chris Verslype^{1,5}, and Jos van Pelt¹

Abstract

Purpose: Hypoxia is considered a major microenvironmental factor influencing cancer behavior. Our aim was to develop a hypoxia-based gene score that could identify high and low risk within stage II and III colon cancer patients.

Experimental Design: Differential gene expression of CaCo-2 colon cancer cells cultured in chronic hypoxia versus normoxia was tested for correlation with prognostic variables in published microarray datasets. These datasets were further used to downsize and optimize a gene score, which was subsequently determined in paraffin-embedded material of 126 patients with colon cancer treated in our center.

Results: In the CaCo-2 cells, 923 genes with a 2-fold change and Limma corrected $P \leq 0.0001$ were found differentially expressed in hypoxia versus normoxia. We identified 21 genes with prognostic value and overlapping in three different training sets and ($n = 224$). With a fourth published dataset ($n = 177$), the six-gene Colon Cancer Hypoxia Score (CCHS) was developed. Patients with low CCHS showed a significant better disease-free survival at three years (77.3%) compared with high CCHS patients (46.4%; log-rank, $P = 0.006$). This was independently confirmed in an external patient cohort of 90 stage II patients (86.9% vs. 52.2%; $P = 0.001$).

Conclusions: Hypoxia-driven gene expression is associated with high recurrence rates in stage II and III colon cancer. A six-gene score was found to be of independent prognostic value in these patients. Our findings require further validation and incorporation in the current knowledge on molecular classification of colon cancer. *Clin Cancer Res*; 20(8); 2159–68. ©2014 AACR.

Introduction

In Western countries, colorectal cancer is the third most common cancer in both men and women and the second leading cause of cancer-related death, accounting for approximately 500,000 deaths annually (1). Tumor stage (American Joint Committee on Cancer; AJCC) at presentation is the main factor for therapeutic decisions and prognostic estimates. Patients with stage II disease generally undergo surgery only, whereas in stage III, adjuvant chemotherapy after resection of the affected bowel segment and attached lymph nodes is the preferred treatment option (2). However, data from historical trials show that 60% of

stage III patients do not recur following surgery only, whereas 20% of stage II patients do have recurrent disease after resection (3, 4). These findings have led to an extensive search for prognostic markers that could classify patients with colorectal cancer more accurately, reducing over- or undertreatment. The finding that stage II patients with a high degree of tumor microsatellite instability (MSI-H) are at very low risk of recurrence helps guiding treatment (5). Driven by the success of this approach in breast cancer (6), extensive research has led to the development of dozens of prognostic gene signatures for colon cancer (7). Although used by some oncologists, gene signatures are currently neither approved by the U.S. Food and Drug Administration nor adopted by any colorectal cancer treatment guideline (8).

In this study, the aim was to develop a robust prognostic gene score to improve the subclassification of stage II and III colon cancer patients using a mechanism-based approach previously described for hepatocellular carcinoma (9). It is now recognized that microenvironmental factors such as hypoxia play a role in tumor behavior. Exposed to low oxygen supplies, cancer cells become more aggressive, invasive, and resistant to therapy (10).

In vitro, we identified the differentially expressed genes under chronic hypoxia versus normoxia. Using bioinformatics on three published sets of expression data with

Authors' Affiliations: ¹Laboratory of Hepatology, Department of Clinical and Experimental Medicine, University Hospitals Leuven; ²Department of Electrical Engineering (ESAT), STADIUS–iMinds Future Health Department, KU Leuven; Departments of ³Abdominal Surgical Oncology; ⁴Imaging and Pathology, and ⁵Clinical Digestive Oncology, University Hospitals Leuven, Leuven, Belgium

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Corresponding Author: Jos van Pelt, Laboratory of Hepatology, KU Leuven, Herestraat 49, bus 703, 3000 Leuven, Belgium. Phone: 32-16-330694; Fax: 32-16-330701; E-mail: Jos.vanPelt@med.kuleuven.be

doi: 10.1158/1078-0432.CCR-13-2958

©2014 American Association for Cancer Research.

Translational Relevance

Hypoxia is a well-known microenvironmental factor determining tumor behavior. Despite this knowledge, its influence is never taken into account in clinical practice. We looked at gene expression changes in colon cancer cells exposed to chronic hypoxia. The hypoxia-induced expression pattern correlated with the clinical outcome of more than 200 patients with colon cancer in published microarray datasets, confirming the detrimental role of intratumoral hypoxia on prognosis. We further describe the development of a 6-gene Colon Cancer Hypoxia Score (CCHS), which was found prognostic in two additional patient cohorts. Its value was independent of other important variables such as disease stage, differentiation grade, and vascular invasion. Moreover, the CCHS performed excellent in both fresh-frozen and paraffin-embedded material.

corresponding clinical information, we could identify 21 genes, related to both hypoxia and prognosis. Further optimization of the model led to a Colon Cancer Hypoxia Score (CCHS) of six genes, which we evaluated in formalin-fixed, paraffin-embedded (FFPE) tissue of our own patient cohort as well as in an external dataset.

Materials and Methods

Cell culture and RNA isolation

Method details are described in the Supplementary Files. Briefly, as *in vitro* model, we used the human adenocarcinoma cell line CaCo-2 (HTB-37; American Type Culture Collection). Cells were grown in a humidified incubator (Sanyo MCO-18M O₂/CO₂ incubator; 5% CO₂ at 37°C) in MEM medium (Invitrogen) supplemented with 10% fetal calf serum, 2 mmol/L L-glutamine, 100 U/mL penicillin, 100 µg/mL streptomycin, and 1 mmol/L sodium pyruvate.

For the determination of chronic hypoxia conditions by real-time PCR (RT-PCR), cells were seeded in 25 cm² culture flasks (10⁶ cells/flask) that were placed in either 2% O₂ or 20% O₂ after 24-hour attachment. All culture conditions were performed in triplicate and cells were collected for RNA isolation at 0, 10, 24, 48, and 72 hours.

For microarray analysis, cells were seeded at 3 × 10⁶ in 75 cm² tissue culture flasks (*n* = 4) at 20% O₂ and were grown until 70% confluence. After reaching near-confluence, two flasks were placed in a humidified incubator with hypoxic conditions (2% O₂, 5% CO₂ at 37°C) for 72 hours, whereas two other flasks remained in normoxic conditions (20% O₂).

Cells were harvested after trypsin treatment and mixed with TRIzol (Invitrogen). RNA was isolated with the RNeasy Kit (Qiagen) according to the manufacturer's instructions.

RT-PCR

RNA was reverse transcribed into cDNA using SuperScript II reverse transcriptase and random hexamer primers (Invitrogen). The PCR reaction was carried out using TaqMan

Universal PCR Master Mixture and Assays-on-Demand Gene Expression products (Applied Biosystems). The primers used are listed in Supplementary Table S1. β-2 microglobulin was used as housekeeping gene. RT-PCR amplification and data analysis were performed using the A7500 Fast Real-Time PCR System (Applied Biosystems). The ΔΔC_t method was used to determine the relative gene expression levels.

Microarray

Samples were hybridized on dual color Agilent's Human Whole Genome Oligo Microarray (Cat# G4112F, Agilent) that contained 44 k 60-mer oligonucleotide probes representing around 41,000 well-characterized human transcripts. Sample processing, quality control, and microarray data analysis are described in detail in the Supplementary Materials and Methods. To identify the highly significant differentially expressed genes under chronic hypoxic conditions, probes with a 2-fold change and a corrected *P* value less than 0.0001 after multiple testing corrections were selected (11). Results were confirmed by performing RT-PCR on selected genes.

Correlation of the hypoxia gene expression pattern with prognosis in external datasets

Biomarker development and evaluation were done following the REMARK guidelines for tumor marker studies (12). To determine the clinical relevance of the hypoxia gene expression pattern, we compared our findings with published microarray datasets containing corresponding clinical information.

We considered four datasets of patients with colon carcinoma, published in Gene Expression Omnibus (GEO; Table 1), all based on the GeneChip Human Genome U133 Plus 2.0 array from Affymetrix. In Jorissen and colleagues, 63 microsatellite stable (MSS) and 11 unstable (MSI) colorectal cancer samples from the Royal Melbourne Hospital (Parkville, Australia) were investigated (13). In Aronow and colleague, recurrence status was available for 100 patients with colon cancer after exclusion of mucosa samples (14). A total of 232 patients with colon cancer and known clinical outcome were included in the study by Smith and colleagues (15). This set was divided into 55 samples from Vanderbilt Medical Center (VMC; Nashville, TN) and 177 samples from the Moffitt Cancer Center (MCC; Tampa, FL).

All datasets were independent of one another, except for the dataset of Aronow and colleagues, which shows partial overlap with MCC samples (own observations). For this reason, we used the datasets of Jorissen and Aronow and the VMC subset of Smith to determine the prognostic value of the hypoxia gene expression pattern with a global test by Goeman and colleagues (16).

Subsequently, overlap between the hypoxia gene expression pattern and the three datasets was used to downsize the number of genes (prognostic signature; see Fig. 1).

The MCC subset of Smith was used to optimize this signature by backward regression analysis using SPSS package 19 (IBM). A probability of *F* with a *P* value of 0.05 was set

Table 1. Clinical overview of published microarray datasets used for global performance testing of the hypoxia expression pattern and for development and optimization of the CCHS

Author Reference	Jorissen 13	Aronow 14	Smith 15	
			VMC database	MCC database
Dataset ID	GSE13294	GSE5206	GSE17537	GSE17536
Used for	Training signature	Training signature	Training signature	Developing score
Array type		Affymetrix Human Genome U133 Plus 2.0 Array		
Sample type	Fresh frozen	Fresh frozen	Fresh frozen	Fresh frozen
N samples	74	105	55	177
N colon cancer	74	100	55	177
Rectal cancer included	Yes	Yes	Yes	Yes
N other	—	5 normal mucosa	—	—
Age (median, y)	NA	66	62	66
Min.		26	23	26
Max.		92	94	92
Sex (M/F)	NA	—	—	—
Male	—	46 (46%)	26 (47%)	96 (54%)
Female	—	54 (54%)	29 (53%)	81 (46%)
AJCC stage	NA	—	—	—
I	—	15 (15%)	4 (7%)	24 (14%)
II	—	29 (29%)	15 (27%)	57 (32%)
III	—	33 (33%)	19 (35%)	57 (32%)
IV	—	20 (20%)	17 (31%)	39 (22%)
NA	—	3 (3%)	0	0
Differentiation	NA	—	—	—
Good	—	8 (8%)	1 (2%)	16 (9%)
Moderate	—	78 (78%)	32 (58%)	134 (76%)
Poor	—	10 (10%)	3 (5%)	27 (15%)
NA	—	4 (4%)	19 (35%)	0
Recurrence	NA	—	—	—
Yes	—	23 (23%)	19 (34%)	36 (24%)
No	—	77 (77%)	36 (66%)	109 (62%)
NA	—	—	—	32 (18%)
Median FU (mo.)	NA	NA	50.2	42.3
Min.	—	—	0.4	0.9
Max.	—	—	111	142.5
MSI	—	NA	NA	NA
MSS	63 (85%)	—	—	—
MSI	11 (15%)	—	—	—

Abbreviation: NA, not available.

as the threshold for entry and removal. This way, we selected those genes that contributed the most to the model and enriched them with a coefficient, which describes the relative contribution of the gene. The obtained score was named the "Colon Cancer Hypoxia Score."

Prospective evaluation of the Colon Cancer Hypoxia score

The Leuven cohort. A total of 162 patients with stage II or III colon cancer treated in our center between 2004 and 2006 were selected to test the performance of the CCHS. Patients with missing clinical data, patients with rectal

cancer, and those who did not survive at least 1 month after surgery were excluded. For this study, we used two parallel slides of FFPE material per patient. The first slide was hematoxylin and eosin stained, and the tumor tissue marked by a pathologist. Tumor was then dissected from nontumorous tissue on the second slide and subsequently deparaffinized. RNA was extracted using a modified RNeasy FFPE protocol (Supplementary Materials and Methods). Quantification, sizing, and quality control were performed with the Bioanalyzer platform (Agilent). Only samples with at least 50% of RNA fragments longer than 200 bases were selected. Samples were analyzed for expression of selected




Figure 1. Process to obtain the CCHS. Overlap between three published datasets and the hypoxia gene expression pattern in CaCo-2 cells was used to downsize the number of genes to 21. After adding *TP53*, a backward regression analysis was performed to obtain a six gene score.

genes by the nCounter technique (NanoString Technologies) which gives a direct read-out of transcripts without the use of enzymatic reactions (17). The used CodeSet is described in Supplementary Table S2.

The Amsterdam cohort. The AMC-AJCCII-90 dataset consists of gene expression data on fresh-frozen tumor tissue of 90 stage II patients treated in the Academic Medical Center (AMC) in Amsterdam between 1997 and 2006. The Affymetrix Human Genome U133 Plus 2.0 Array was used as microarray platform (GSE33113; ref. 18).

Statistical analysis

All data are presented as median with range. A $P < 0.05$ was considered significant. To determine an optimal cutoff value for our CCHS, a receiver operating characteristic curve was used. For the survival analysis, Kaplan–Meier survival curves and univariate and multivariate Cox proportional HR were used. All statistics were performed using SPSS package 19 (IBM).

Results

Dynamics in gene expression in CaCo-2

To investigate the dynamics of hypoxia-related gene expression in our *in vitro* model and to determine the optimal timing for microarray analysis, we performed

RT-PCR at different time points for a set of eight representative genes known to be responsive to hypoxia. Gene expression was tested at 0, 10, 24, 48, and 72 hours in hypoxia (2% O_2) versus normoxia. It should be noted that when CaCo-2 cells were kept beyond 72 hours, their cultures tended to become super-confluent with partly detachment of the cells resulting in an uncontrolled fluctuation of the gene expression levels (data not shown). Most of the mRNA expression levels investigated showed a dynamic curve over time at 2% O_2 with exception of *BCL2* that did not change significantly at any of the time points investigated (Supplementary Fig. S1). These observations support the assumption that the acute hypoxic state (up to 24 hours) has a different gene expression pattern compared with the more chronic state (72 hours). As such, we chose the latter condition for our microarray experiment.

Microarray: primary data analysis and development of a Colon Cancer Hypoxia Score

Using Agilent technology, in CaCo-2 cells cultured for 72 hours at either 20% oxygen or in hypoxic conditions at 2% oxygen, a total of 37,707 spots showed a representative signal of which 3,389 with a \log_2 fold-change >1 or <-1 and a Limma corrected p -value <0.05 . These microarray data are available at NCBI under number GSE31079.

For the development of our gene signature, we started with the top 923 differentially expressed genes (with a \log_2 fold-change >1 or <-1 and a Limma corrected $P < 0.0001$): the hypoxia gene expression pattern. Of those 923 genes, 704 were upregulated and 219 downregulated in hypoxic conditions. The top altered genes and significant KEGG pathways involved are listed in Supplementary Tables S3 and S4.

Subsequently, the Goeman global test showed that the overall expression pattern of these hypoxia genes was significantly related to the prognostic factor considered for each of the three training datasets ($P < 0.0001$ for Jorissen; $P = 0.00028$ for Aronow; and $P = 0.0107$ for Smith VMC).

Next, when only keeping the significant genes with a z -score, above 1,530 genes remained for the dataset of Jorissen, 360 genes for Aronow, and 338 genes for Smith VMC. Finally, genes for which the direction of altered expression did not correspond to the direction observed *in vivo* in at least two out of three datasets were removed. With this approach, we were able to downsize the number of genes to twenty-one, found to overlap between the three training datasets (Fig. 1). In this gene set of 21 genes, 17 genes were upregulated in hypoxic conditions and four were downregulated (Table 2).

CaCo-2 cells contain a *TP53* mutation resulting in undetectable p53 protein levels (19). As such, *TP53* was not considered early in the development of our gene signature despite being clinically of potential prognostic importance in colon cancer (20). For these reasons, *TP53* was added afterward to the *in vitro*-derived 21 hypoxia cancer genes.

Using a backward linear regression analysis with the 21 genes and *TP53* as independent variables and the AJCC disease stage of patients in a fourth dataset (Smith MCC) as

Table 2. Overview of the 21 gene signature with their relative direction *in vitro* under hypoxia and cellular processes involved

Official gene symbol	Direction <i>in vitro</i>	Gene name	Processes involved	Used for CCHS
BCCIP	▼	BRCA2 and CDKN1A-interacting protein isoform C	DNA metabolic/repair	✓
BNIP3L	▲	BCL2/adenovirus E1B 19 kDa protein-interacting protein 3-like	Apoptosis	✓
BTBD14A	▲	Nucleus accumbens-associated protein 2	Unknown	—
DDIT4	▲	DNA damage-inducible transcript 4 protein	Apoptosis/hypoxia	—
ERRFI1	▲	ERBB receptor feedback inhibitor 1	GTPase regulation	—
FAM46C	▼	Hypothetical protein LOC54855	Unknown	—
GADD45B	▲	Growth arrest and DNA damage-inducible protein GADD45 beta	Apoptosis	✓
HBP1	▲	HMG box-containing protein 1	Cell cycle	—
ING5	▼	inhibitor of growth protein 5	DNA metabolic/repair	—
INSIG2	▲	insulin-induced gene 2 protein	ER-nuclear signaling	✓
KLF7	▲	Kruppel-like factor 7	Cell morphogenesis	—
MPHOSPH6	▼	M-phase phosphoprotein 6	Cell cycle	✓
NNMT	▲	Nicotinamide N-methyltransferase	N-methyltransferase	—
P4HA1	▲	Prolyl 4-hydroxylase subunit α -1 isoform 1 precursor	Metabolic processes	—
P4HA2	▲	Prolyl 4-hydroxylase subunit α -2 isoform 1 precursor	Metabolic processes	—
PLAUR	▲	Urokinase plasminogen activator surface receptor isoform 3 precursor	Metabolic processes	—
RHO	▲	rho-related GTP-binding protein RhoU	Cell cycle	—
TMCC1	▲	Transmembrane and coiled-coil domains protein 1 isoform a	Transmembrane signaling	—
UBE2H	▲	Ubiquitin-conjugating enzyme E2 H isoform 1	Proteolysis	—
ULBP2	▲	NKG2D ligand 2 precursor	Transmembrane signaling	—
XIAP/BIRC4	▲	Baculoviral IAP repeat-containing protein 4	Apoptosis	—
TP53	NA	Tumor protein p53	Apoptosis/cell cycle	✓

NOTE: *TP53* was added to the signature and eventually incorporated in the CCHS.

dependent variable, six genes showed the highest contribution to the model. These genes with their corresponding coefficient form the CCHS (Fig. 1). The score consists of the expression value of *BCCIP*, *GADD45B*, *INSIG2*, *BNIP3L*, *MPHOSPH6*, and *TP53*. The latter three have an inverse effect on the score value. All genes are involved in processes related to cell cycle, apoptosis, and DNA repair.

Prospective evaluation of CCHS

Of the 162 stage II/III patients treated in our center between 2004 and 2006, 129 were selected for nCounter analysis after exclusion of clinical aberrant cases and quality control (Fig. 2). Gene expression level of the CCHS genes was successfully obtained in 126 samples (97.7%). The known clinical and pathologic features of these patients are summarized in Table 3. Estimated median disease-free survival (DFS) times were 66.2 months [95% confidence interval (CI), 60–71.5] and 46 months (95% CI, 37.5–54.5) for stage II and stage III patients, respectively.

Using an optimal cutoff (see Supplementary Fig. S2), patients with a low CCHS showed to have significant better DFS at 3 years compared with those with a high CCHS (77.3% vs. 46.4%, respectively; $P = 0.006$). These DFS rates remained unchanged at 5 years. Median DFS was 62.2




Figure 2. Inclusion and selection process of the stage II/III colon cancer patients in our center. Of the 162 patients with colon cancer included in this study, after clinical and technical selection, the expression levels of 126 patients were successfully determined.

Table 3. Clinical and pathologic features of the Leuven cohort as well as the AMC-AJCCII-90 dataset (16)

	Leuven cohort			AMC-AJCC-II
	All patients	Stage II	Stage III	
Patient number	126	71	55	90
Age (median, y)	71.2	70	72.9	73.4
Min. (y)	35.9	35.9	38.8	34.6
Max. (y)	92.9	92.9	86.2	95.1
Sex				
Male	67 (53%)	34 (48%)	33 (60%)	42 (47%)
Female	59 (47%)	37 (52%)	22 (40%)	48 (53%)
Localization				
Right-sided tumor	63 (50%)	39 (55%)	24 (43.5%)	48 (53%)
Left-sided tumor	63 (50%)	32 (45%)	31 (56.5%)	42 (47%)
T-stage				NA
T2	4 (3%)	0 (0%)	4 (7%)	—
T3	109 (86.5%)	67 (94%)	42 (76.5%)	—
T4	13 (10.5%)	4 (6%)	9 (16.4%)	—
N-stage				NA
N0	71 (56%)	71 (100%)	0 (0%)	—
N1	39 (31%)	0 (0%)	39 (71%)	—
N2	16 (13%)	0 (0%)	16 (29%)	—
AJCC stage				
II	71 (56%)	71 (100%)	0 (0%)	90 (100%)
III	55 (44%)	0 (0%)	55 (100%)	0 (0%)
Mucinous type				NA
Yes	36 (29%)	22 (31%)	41 (74, 5%)	—
No	90 (71%)	49 (69%)	14 (25, 5%)	—
Differentiation				NA
Good	14 (11%)	7 (10%)	7 (13%)	—
Moderate	63 (50%)	40 (56.5%)	23 (42%)	—
Poor	45 (36%)	20 (28%)	25 (45%)	—
Undifferentiated	1 (1%)	1 (1.5%)	0 (0%)	—
Not available	3 (2%)	3 (4%)	0 (0%)	—
Peritoneal invasion				NA
Yes	1 (1%)	1 (1%)	0 (0%)	—
No	125 (99%)	70 (99%)	55 (100%)	—
Extramural venous invasion				NA
Yes	36 (28.6%)	11 (15%)	25 (45%)	—
No	81 (64.3%)	59 (84%)	22 (40%)	—
NA	9 (7.1%)	1 (1%)	8 (15%)	—
Resection margin				NA
Positive	1 (1%)	0 (0%)	1 (2%)	—
Negative	124 (98%)	71 (100%)	53 (96%)	—
NA	1 (1%)	0 (0%)	1 (2%)	—
Bowel perforation				NA
Yes	5 (4%)	3 (4%)	2 (4%)	—
No	121 (96%)	68 (96%)	53 (96%)	—
Examined lymph nodes				NA
12 or more	83 (66%)	49 (69%)	34 (62%)	—
Less than 12	42 (33%)	22 (31%)	20 (36%)	—
NA	1 (1%)	0 (0%)	1 (2%)	—
Adjuvant chemotherapy				NA
Yes	63 (50%)	50 (70.4%)	42 (76.5%)	—
No	63 (50%)	21 (29.6%)	13 (23.5%)	—

(Continued on the following page)

Table 3. Clinical and pathologic features of the Leuven cohort as well as the AMC-AJCCII-90 dataset (16) (Cont'd)

	Leuven cohort			AMC-AJCC-II
	All patients	Stage II	Stage III	
Relapse				—
Yes	35 (28%)	12 (17%)	23 (42%)	19 (21%)
No	91 (72%)	59 (83%)	32 (58%)	71 (79%)
Median FU (mo)	45.42	49.14	23.1	39.3
Min. (mo)	1.3	1.3	1.5	1.6
Max. (mo)	76.4	76.4	73.9	120

Abbreviations: Median FU, median follow-up defined as time to relapse or last contact; NA, not available.

months (95% CI, 56.6–67.7) and 42.7 months (95% CI, 33.4–52) for CCHS low and CCHS high patients, respectively ($P = 0.006$; Fig. 3A). Patients with a favorable CCHS were more likely to be alive at 3 years (estimated overall survival 81.5% vs. 66.6%; $P = 0.044$) and at 5 years (71.9% vs. 59.1%; $P = 0.036$).

When disease AJCC stage was taken into account, survival analysis could identify three groups with significant difference in survival (Fig. 3B). Patients with stage II disease and unfavorable CCHS had almost similar disease behavior as stage III patients with a good (low) CCHS. Moreover, the CCHS could identify a subgroup of stage III patients with an estimated recurrence rate at 3 years of 75% and a median DFS of only 28.2 months (95% CI, 16–40.4).

The characteristics of the 90 patients of the Amsterdam cohort are listed in Table 3. Estimated DFS time for the whole cohort was 95.5 months (95% CI, 85.8–105.2). Similar to the Leuven patient cohort, the CCHS identified two subgroups of stage II patients with different DFS at 3 years (86.9% vs. 52.2%; $P < 0.001$) and 5 years (86.9% vs. 52.2%; $P < 0.001$). Median DFS was 102.5 months (95% CI, 93.1–111.9) versus 55.4 months (95% CI, 38.7–72; $P = 0.001$; Fig. 3C). HR for recurrence in CCHS high versus CCHS low patients was 3.970 (95% CI, 1.6–9.8; $P = 0.004$).

Subsequently, a univariate Cox regression analysis was performed on the Leuven cohort analyzing all known clinical and pathologic variables. Four variables with significant prognostic value were included in a multivariate analysis. The multivariate Cox regression model confirmed that the CCHS is a predictor for recurrence independent of disease stage, venous invasion, and the number of examined lymph nodes (Table 4).

Discussion

Colon cancer is associated with high morbidity and mortality, and prognostic subclassification of stage II and III patients remains an ongoing clinical challenge. We describe the development and testing of a mechanism-based prognostic gene score. Hypoxia as microenvironmental factor is well known for its influence on cancer behavior. If hypoxia does not induce cell death, it increases tumor

aggressiveness, invasiveness, and metastatic potential mainly through activation of the hypoxia-inducible factor (HIF) pathway (10). Despite this knowledge, hypoxia is seldom taken into account in the clinical setting, in part, due to the lack of good endogenous hypoxia markers (21). HIF overexpression has been proven associated with prognosis, but HIF itself is not considered to be a hypoxia marker exclusively due to the fact that it can be activated in various settings, including oncogene drive (22). Our method shows a correlation between the altered expressions of hundreds of genes under hypoxic conditions *in vitro*, with that of tumors in published microarray datasets with clinical data available. This confirms the role of intratumoral hypoxia in disease behavior as was previously shown in HCC (9).

Colon cancer is subject to extensive global scientific research and many gene signatures and scores have been developed in the past mostly based on hierarchical clustering. Some of these signatures have made it to commercially available tests. Although a promising approach to capture the complexity of cancer biology, few signatures show good power in external patient cohorts (7). We tried to address these limitations by using a mechanism-driven approach and involving different datasets in the development of our signature. This resulted in a workable amount of genes derived from data of more than 400 patient samples. Moreover, we hypothesized that the optimization process with a backward regression model has added power to the prognostic value of the genes.

p53 is a known important tumor suppressor in colon cancer and its pathway is modulated by hypoxia (23). This is further supported by our findings that, *in vitro* under hypoxic conditions, the p53 pathway is a highly significant altered KEGG pathway (Supplementary Table S4 and Supplementary Fig. S3). For this reason and those stated above, *TP53* was appended to the gene list in the knowledge that the backward regression analysis would eliminate this gene immediately in the absence of an added value to the model. The fact that it was withheld in the CCHS suggests that *TP53* gene expression is of prognostic value for patients with colon cancer.

Evaluation of the CCHS was done in FFPE material. For mRNA level measurements, this type of material often




Figure 3. A, Kaplan-Meier curve for DFS in the Leuven cohort ($n = 126$). B, AJCC disease stage and CCHS divide patients in three groups with significant differences in DFS. When the patients of AJCC with stage II disease and a high CCHS have comparable outcome to patients with stage III disease and a low CCHS. C, Kaplan-Meier curve for DFS in the Amsterdam cohort ($n = 90$).

presents a challenge as it is subjected to RNA degradation. After strict quality control (Fig. 2), the nCounter platform showed excellent performance and reliability for determining the expression levels of all six CCHS genes. Independent of other clinical and pathologic parameters, the CCHS was found to predict relapse in stage II and III colon cancer. In stage II patients, it identified a high-risk subgroup that

might benefit from adjuvant therapy. Furthermore, stage III patients with unfavorable CCHS suffer from high recurrence rates.

Recently, progress has been made in the field of molecular classification of colon cancer based on gene expression clustering, epigenetic markers, and therapy response. This approach led to the proposition of three to six colorectal

Table 4. Univariate and multivariate Cox regression analysis on all clinical and pathologic variables in the Leuven cohort

	Univariate Cox regression analysis					Multivariate Cox regression analysis				
	β coefficient	<i>P</i>	HR	95% CI		β coefficient	<i>P</i>	HR	95% CI	
				Lower	Upper				Lower	Upper
Age	0.03	0.070	1.03	1.00	1.07					
Location (left vs. right)	0.13	0.701	1.14	0.59	2.21					
Stage (III vs. II)	1.23	0.001	3.41	1.69	6.86	0.95	0.017	2.58	1.19	5.61
Peritoneal invasion (yes vs. no)	1.30	0.203	3.66	0.50	26.89					
Extramural venous invasion (yes vs. no)	1.60	<0.001	4.95	2.43	10.07	1.17	0.003	3.21	1.50	6.89
Resection margin (positive vs. negative)	1.47	0.149	4.36	0.59	32.24					
Bowel perforation (yes vs. no)	0.70	0.336	2.02	0.48	8.45					
Mucinous type (yes vs. no)	−0.27	0.506	0.76	0.35	1.68					
Differentiation		0.856								
Differentiation (moderate vs. good)	0.38	0.541	1.46	0.43	4.97					
Differentiation (poor vs. good)	0.66	0.296	1.94	0.56	6.77					
Adjuvant chemotherapy (yes vs. no)	0.19	0.587	1.20	0.62	2.35					
CCHS (high vs. low)	0.91	0.008	2.49	1.27	4.86	0.95	0.013	2.58	1.23	5.43
Lymph nodes examined (12 or more vs. less than 12)	−0.73	0.034	0.48	0.25	0.95	−0.40	0.283	0.67	0.33	1.39

NOTE: CCHS is an independent predictor of disease recurrence.

cancer subtypes (18, 24, 25). De Sousa and colleagues identified two classes which seem to represent the known subgroups of MSI/CIMP and chromosomal-unstable tumors, respectively (18). A third class, previously unidentified and related to serrated polyps, was associated with a poor prognosis. Tumors of this subgroup have stemness features, which is in line with others who have isolated a stem cell or epithelial-to-mesenchymal transition associated subgroup of patients with colon cancer with high recurrence rates (24–26). We see a striking overlap between the KEGG pathways involved in adapting to hypoxic conditions and those pathways attributed to the stem cell-like phenotype proposed by Marisa and colleagues (ref. 25; Supplementary Table S4). Further prospective validation in a larger patient cohort is needed to clarify whether a high CCHS is indicative for the stemness subtype. If so, the limited number of genes together with the performance in FFPE samples renders our CCHS a potential clinical tool.

This study has some limitations. Unfortunately, at the time of study onset, extensive clinical data were not available for some of the published datasets used for CCHS development. This forced us to use surrogate parameters such as MSI which could influence the power of the obtained score. During study design, after careful evaluation of the available information, considerable overlap between different datasets published in the GEO was noticed. For example, most samples of the dataset provided by Aronow and colleagues (GSE5206) have also been included in both GSE14333 and GSE17536 (own observations). This requires great caution when designing studies with this widely used published data. In our case, we

compared the individual patient data of the datasets and found that there was a partial overlap between the database by Aronow and Smith MCC. In our study design, the latter was only involved in optimization of the gene signature and this overlap should therefore not affect our conclusions.

When testing the performance of the CCHS, we excluded rectal cancer. Despite the obvious similarities between colon and rectal cancer, we wanted to exclude the differences in treatment modalities as a confounder. As such, our results cannot be extrapolated to these patients.

Finally, our study was designed to evaluate the CCHS as a prognostic marker. Further research is needed to determine whether the score has predictive abilities, that is, potency to identify patients likely to benefit from adjuvant treatment.

In conclusion, we discovered that the gene expression pattern of CaCo-2 cells cultured in hypoxia shows good correlation with clinical parameters of patients with colon cancer in published microarray datasets. This confirms the importance of intratumoral hypoxia in disease behavior. From this point, a six-gene CCHS was developed which proved to be an independent prognostic biomarker for relapse in stage II and III colon cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J. Dekervel, H.v. Malenstein, D. Popovic, X. Sagaert, E.V. Cutsem, C. Verslype, J.v. Pelt

Development of methodology: J. Dekervel, H.v. Malenstein, D. Popovic, E.V. Cutsem, C. Verslype, J.v. Pelt

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): J. Dekervel, D. Hompes, H.v. Malenstein, X. Sagaert, E.V. Cutsem, A. D'Hoore, C. Verslype, J.v. Pelt
Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J. Dekervel, D. Hompes, H.v. Malenstein, D. Popovic, E.V. Cutsem, A. D'Hoore, C. Verslype, J.v. Pelt
Writing, review, and/or revision of the manuscript: J. Dekervel, H.v. Malenstein, D. Popovic, X. Sagaert, B. De Moor, E.V. Cutsem, C. Verslype, J.v. Pelt
Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): D. Hompes, D. Popovic, A. D'Hoore, C. Verslype
Study supervision: J. Dekervel, B. De Moor, E.V. Cutsem, C. Verslype, J.v. Pelt

Acknowledgments

The authors thank P. Windmolders for her excellent technical assistance and Prof. F. Nevens for his support, Dr. A. Daemen (ESAT) for her contribution to the set-up of this study and the VIB SWITCH laboratory (Profs. J.

Schymkowitz and F. Rousseau) for their cooperation and discussion of the results. nCounter analysis was performed by VIB Nucleomics Core (Drs. R. van Eijnsden and J. Allemeersch; www.nucleomics.be).

Grant Support

This work was supported by GOA/10/09 MaNet, IWT: TBM Rectal Cancer, iMinds SBO, VLK Stichting E. van der Schueren: rectal cancer, Federal Government Cancer Plan 2012-2015 KPC-29-023 (prostate). J. Dekervel was supported by the Flemish League against Cancer (VLK). X. Sagaert is a Senior Clinical Investigator of the Fund for Scientific Research-Flanders (FWO-Vlaanderen).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received October 28, 2013; revised January 10, 2014; accepted January 18, 2014; published OnlineFirst January 31, 2014.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127:2893-917.
2. Schmoll HJ, Van Cutsem E, Stein A, Valentini V, Glimelius B, Haustermans K, et al. ESMO Consensus Guidelines for management of patients with colon and rectal cancer. A personalized approach to clinical decision making. *Ann Oncol* 2012;23:2479-516.
3. Moertel CG, Fleming TR, Macdonald JS, Haller DG, Laurie JA, Tangen CM, et al. Fluorouracil plus levamisole as effective adjuvant therapy after resection of stage III colon carcinoma: a final report. *Ann Intern Med* 1995;122:321-6.
4. Quasar Collaborative G, Gray R, Barnwell J, McConkey C, Hills RK, Williams NS, et al. Adjuvant chemotherapy versus observation in patients with colorectal cancer: a randomised study. *Lancet* 2007;370:2020-9.
5. Tejpar S, Bertagnolli M, Bosman F, Lenz HJ, Garraway L, Waldman F, et al. Prognostic and predictive biomarkers in resected colon cancer: current status and future perspectives for integrating genomics into biomarker discovery. *Oncologist* 2010;15:390-404.
6. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 2002;415:530-6.
7. Sanz-Pamplona R, Berenguer A, Cordero D, Riccadonna S, Sole X, Crous-Bou M, et al. Clinical value of prognosis gene expression signatures in colorectal cancer: a systematic review. *PLoS ONE* 2012;7:e48877.
8. Black ER, Falzon L, Aronson N. Gene Expression Profiling for Predicting Outcomes in Stage II Colon Cancer. Rockville MD; 2012.
9. van Malenstein H, Gevaert O, Libbrecht L, Daemen A, Allemeersch J, Nevens F, et al. A seven-gene set associated with chronic hypoxia of prognostic importance in hepatocellular carcinoma. *Clin Cancer Res* 2010;16:4278-88.
10. Semenza GL. Hypoxia-inducible factors: mediators of cancer progression and targets for cancer therapy. *Trends Pharmacol Sci* 2012;33:207-14.
11. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Statist Soc B* 1995;57:289-300.
12. McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM, et al. REporting recommendations for tumour MARKer prognostic studies (REMARK). *Br J Cancer* 2005;93:387-91.
13. Jorissen RN, Gibbs P, Christie M, Prakash S, Lipton L, Desai J, et al. Metastasis-associated gene expression changes predict poor outcomes in patients with dukes stage B and C colorectal cancer. *Clin Cancer Res* 2009;15:7642-51.
14. Kaiser S, Park YK, Franklin JL, Halberg RB, Yu M, Jessen WJ, et al. Transcriptional recapitulation and subversion of embryonic colon development by mouse colon tumor models and human colon cancer. *Genome Biol* 2007;8:R131.
15. Smith JJ, Deane NG, Wu F, Merchant NB, Zhang B, Jiang A, et al. Experimentally derived metastasis gene expression profile predicts recurrence and death in patients with colon cancer. *Gastroenterology* 2010;138:958-68.
16. Goeman JJ, van de Geer SA, de Kort F, van Houwelingen HC. A global test for groups of genes: testing association with a clinical outcome. *Bioinformatics* 2004;20:93-9.
17. Geiss GK, Bumgarner RE, Birditt B, Dahl T, Dowidar N, Dunaway DL, et al. Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol* 2008;26:317-25.
18. De Sousa EMF, Wang X, Jansen M, Fessler E, Trinh A, de Rooij LP, et al. Poor-prognosis colon cancer is defined by a molecularly distinct subtype and develops from serrated precursor lesions. *Nat Med* 2013;19:614-8.
19. Liu Y, Bodmer WF. Analysis of P53 mutations and their expression in 56 colorectal cancer cell lines. *Proc Natl Acad Sci U S A* 2006;103:976-81.
20. Sarasqueta AF, Forte G, Corver WE, de Miranda NF, Ruano D, van Eijk R, et al. Integral analysis of p53 and its value as prognostic factor in sporadic colon cancer. *BMC Cancer* 2013;13:277.
21. Moon EJ, Brizel DM, Chi JT, Dewhirst MW. The potential role of intrinsic hypoxia markers as prognostic variables in cancer. *Antioxid Redox Signal* 2007;9:1237-94.
22. Yoshimura H, Dhar DK, Kohno H, Kubota H, Fujii T, Ueda S, et al. Prognostic impact of hypoxia-inducible factors 1alpha and 2alpha in colorectal cancer patients: correlation with tumor angiogenesis and cyclooxygenase-2 expression. *Clin Cancer Res* 2004;10:8554-60.
23. Sermeus A, Michiels C. Reciprocal influence of the p53 and the hypoxic pathways. *Cell Death Dis* 2011;2:e164.
24. Sadanandam A, Lyssiotis CA, Homiczko K, Collisson EA, Gibb WJ, Wulschleger S, et al. A colorectal cancer classification system that associates cellular phenotype and responses to therapy. *Nat Med* 2013;19:619-25.
25. Marisa L, de Reynies A, Duval A, Selves J, Gaub MP, Vescovo L, et al. Gene expression classification of colon cancer into molecular subtypes: characterization, validation, and prognostic value. *PLoS Med* 2013;10:e1001453.
26. Budinska E, Popovici V, Tejpar S, D'Ario G, Lapique N, Sikora KO, et al. Gene expression patterns unveil a new level of molecular heterogeneity in colorectal cancer. *J Pathol* 2013;231:63-76.

Clinical Cancer Research

Hypoxia-Driven Gene Expression Is an Independent Prognostic Factor in Stage II and III Colon Cancer Patients

Jeroen Dekervel, Daphne Hompes, Hannah van Malenstein, et al.

Clin Cancer Res 2014;20:2159-2168. Published OnlineFirst January 31, 2014.

Updated version	Access the most recent version of this article at: doi: 10.1158/1078-0432.CCR-13-2958
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2014/02/04/1078-0432.CCR-13-2958.DC1

Cited articles	This article cites 25 articles, 7 of which you can access for free at: http://clincancerres.aacrjournals.org/content/20/8/2159.full.html#ref-list-1
Citing articles	This article has been cited by 4 HighWire-hosted articles. Access the articles at: /content/20/8/2159.full.html#related-urls

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org .