

Germline Variants and Advanced Colorectal Adenomas: Adenoma Prevention with Celecoxib Trial Genome-wide Association Study

Jiping Wang¹, Luis G. Carvajal-Carmona³, Jen-Hwa Chu², Ann G. Zauber⁴, APC Trial Collaborators, Michikai Kubo⁵, Koichi Matsuda⁵, Malcolm Dunlop⁶, Richard S. Houlston⁷, Oliver Sieber⁸, Lara Lipton⁸, Peter Gibbs⁸, Nicholas G. Martin⁹, Grant W. Montgomery⁹, Joanne Young¹⁰, Paul N. Baird¹¹, Mark J. Ratain¹², Yusuke Nakamura⁵, Scott T. Weiss², Ian Tomlinson³, and Monica M. Bertagnolli¹

Abstract

Purpose: Identification of single-nucleotide polymorphisms (SNP) associated with development of advanced colorectal adenomas.

Experimental Design: Discovery phase: 1,406 Caucasian patients (139 advanced adenoma cases and 1,267 controls) from the Adenoma Prevention with Celecoxib (APC) trial were included in a genome-wide association study (GWAS) to identify variants associated with postpolypectomy disease recurrence. Genome-wide significance was defined as false discovery rate less than 0.05, unadjusted $P = 7.4 \times 10^{-7}$. Validation phase: results were further evaluated using 4,175 familial colorectal adenoma cases and 5,036 controls from patients of European ancestry [COLOrectal Gene Identification consortium (CORGI), Scotland, Australia, and VQ58].

Results: Our study identified eight SNPs associated with advanced-adenoma risk in the APC trial (rs2837156, rs7278863, rs2837237, rs2837241, rs2837254, rs741864 at 21q22.2, and rs1381392 and rs17651822 at 3p24.1, at $P < 10^{-7}$ level with $OR > 2$). Five variants in strong pairwise linkage disequilibrium (rs7278863, rs2837237, rs741864, rs741864, and rs2837241; $r^2 = 0.8-1$) are in or near the coding region for the tight junction adhesion protein, IGSF5. An additional variant associated with advanced adenomas, rs1535989 [minor allele frequency, 0.11; OR, 2.09; 95% confidence interval (CI), 1.50–2.91], also predicted colorectal cancer development in a validation analysis ($P = 0.019$) using a series of adenoma cases or colorectal cancer (CORGI study) and 3 sets of colorectal cancer cases and controls (Scotland, VQ58, and Australia; $N = 9,211$).

Conclusions: Our results suggest that common polymorphisms contribute to the risk of developing advanced adenomas and might also contribute to the risk of developing colorectal cancer. The variant at rs1535989 may identify patients whose risk for neoplasia warrants increased colonoscopic surveillance. *Clin Cancer Res*; 1–8. ©2013 AACR.

Authors' Affiliations: ¹Department of Surgery, Division of Surgical Oncology, ²Center for Genomic Medicine, Channing Laboratory, Brigham and Women's Hospital, Boston, Massachusetts; ³Genome Center and Department of Biochemistry and Molecular Medicine, University of California, Davis, California; ⁴Department of Epidemiology and Statistics, Memorial Sloan-Kettering Cancer Center, New York; ⁵RIKEN Center for Genomic Medicine, Tokyo, Japan; ⁶Colon Cancer Genetics Group, Institute of Genetics and Molecular Medicine, University of Edinburgh and MRC Human Genetics Unit, Edinburgh; ⁷Section of Cancer Genetics, Institute of Cancer Research, Sutton, United Kingdom; ⁸Ludwig Colon Cancer Initiative Laboratory, Ludwig Institute for Cancer Research, Royal Melbourne Hospital, Parkville; ⁹Genetic and Molecular Epidemiology Laboratories, and ¹⁰Familial Cancer Laboratory, Queensland Institute of Medical Research, Brisbane; ¹¹Centre for Eye Research Australia, University of Melbourne, Royal Victorian Eye and Ear Hospital, East Melbourne, Australia; and ¹²Department of Medicine, University of Chicago, Chicago, Illinois

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Corresponding Author: Monica M. Bertagnolli, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115. Phone: 617-732-8910; Fax: 617-582-6177; E-mail: mbertagnolli@partners.org

doi: 10.1158/1078-0432.CCR-13-0550

©2013 American Association for Cancer Research.

Introduction

Colorectal cancer is a common malignancy, with prevalence in developed nations of 40 to 50 cases per 100,000 individuals (1). Approximately one-third of those diagnosed with colorectal cancer will die of their disease due to diagnosis at a stage not curable by locoregional therapy. Most colorectal cancer cases arise from premalignant adenomas that require years or even decades to progress to invasive disease. Colonoscopy to identify and remove precursor adenomas has been recommended for more than 25 years for patients at high colorectal cancer risk, and recently completed long-term analyses of screened cohorts confirmed the utility of adenoma removal for preventing deaths due to colorectal cancer (2).

Our goal is to understand the biology of colorectal cancer to develop effective prevention and therapy, and also to characterize individual risk in a manner that will identify patients most likely to benefit from colonoscopy to detect

Translational Relevance

Identification of patients at highest risk of colorectal cancer is essential for providing optimal disease screening and prevention. This study uncovers germline susceptibility loci that indicate risk of disease and potential for improved understanding of disease biology.

and remove premalignant adenomas. Identification of germline variants conveying an increased risk of colorectal cancer could be used to promote adherence to colonoscopy and polypectomy for patients at highest risk, improving utilization and cost-benefit of this life-saving procedure. In addition, more accurate characterization of high-risk individuals would facilitate participation in prevention clinical trials. Finally, therapy to treat or prevent colorectal cancer would be advanced if the biologic consequences of germline susceptibility variants were further characterized to uncover the molecular basis of colorectal cancer.

The adenoma–carcinoma sequence in the colorectum represents a disease spectrum. Adenomas with a low risk of cancer development are small (<0.6 cm diameter) and lack histologic features associated with progression, such as the presence of villous features or high-grade dysplasia. The identification of an advanced adenoma (size ≥ 1 cm; villous or tubulovillous histology; high-grade dysplasia) indicates that a patient has a higher risk of future adenoma and colorectal cancer development (3). Advanced adenomas, therefore, are the most important lesions to target for colorectal cancer prevention. In this study, we used a large cohort of patients with adenoma from a prospective randomized clinical trial to identify single-nucleotide polymorphisms (SNP) associated with increased risk of developing advanced adenomas. These variants were then further tested using large genotyped cohorts of patients and controls with advanced adenomas and colorectal cancer. In doing so, we identified variants associated with both advanced premalignant lesions and colorectal cancer.

Materials and Methods

Study design and populations

The overall study design is illustrated in Fig. 1.

Discovery phase. Of note, 1,406 evaluable Caucasian patients were identified from the Adenoma Prevention with Celecoxib (APC) trial, a randomized, placebo-controlled study to test whether celecoxib reduced the occurrence of endoscopically detected colorectal adenomas. The endpoint advanced adenoma was defined as any adenoma with size 1 cm or more, villous/tubulovillous histology, or high-grade dysplasia. During the prospective follow-up period, 139 participants developed advanced adenomas identified during a scheduled colonoscopy screening examination. Detailed information about the trial design and primary outcomes was reported elsewhere (4).

Validation phase. The advanced adenoma susceptibility that SNPs identified from APC trial were further evaluated using genome-wide association study (GWAS) data from the following four nonoverlapping colorectal cancer case-control series of European ancestry (5).

1. CORGI: 931 familial colorectal adenoma or colorectal cancer cases and 929 cancer-free controls of White British origin ascertained through the COloRectal Gene Identification (CORGI) consortium. All cases had at least one first-degree relative with colorectal tumors and no mutations in the known highly penetrant colorectal cancer genes. Controls were spouses or partners of the cases and had no personal history of colorectal cancer (6).
2. Scotland: 1,003 early-onset Scottish colorectal cancer cases (<55 years) and 979 cancer-free Scottish population controls. Known Mendelian syndromes were excluded. Controls were matched by age (± 5 years), gender, and area of residence (6).
3. VQ58: 1,800 British Stage II/III patients with colorectal cancer from the VICTOR ($N = 923$) and QUASAR2 (<http://www.octo-oxford.org.uk/alltrials/trials/q2.html>; $N = 877$) clinical trials, together with publicly available data from 2,690 population controls from the Wellcome Trust Case-Control Consortium (WTCCC) 1958 Birth Cohort (7).




Figure 1. Study design. *Including subjects with familial adenoma from CORGI. AA, advanced adenoma; CRC, colorectal cancer.

4. Australia: 441 colorectal cancer cases treated in the Royal Melbourne, Western and St Francis Xavier Cabrini Hospitals in Melbourne and 438 population controls from Brisbane Twin Nevus and Genes in Myopia studies, matched to the cases using principal component analysis (6).


Thus, 4,175 familial colorectal adenoma or colorectal cancer cases and 5,036 controls were included in the validation analysis. Human Subjects Committee approval to collect and genotype whole blood samples was obtained by Brigham and Women's Hospital and the RIKEN Center for Genomic Medicine.

Genotyping and quality control

DNA was isolated from blood samples using standard methods and quantified with picogreen. For the APC cohort, genotyping was performed by the RIKEN Center

for Genomic Medicine using the Illumina Human610-Quad BeadChip platform (Illumina). A White parent-child Centre d'Etude du Polymorphisme Humain (CEPH) trio from the HapMap was used to check for Mendelian transmission of alleles. χ^2 test based on genotype frequencies at each SNP was used to test for deviations from Hardy-Weinberg equilibrium (HWE). Any SNP with HWE $P < 0.001$ was excluded. Two cases and two controls were randomly chosen as duplicates for quality control of genotype concordance. A total of 28 subjects and 1,792 markers were excluded for quality control reasons, including duplicates, those that showed identity-by-descent more than 12.5% or were gender mismatched, samples with less than 98%, and markers with less than 99% call rate or heterozygous haploids. The final Manhattan plot and QQ plot indicated the satisfactory quality control process (Supplementary Figs. S1 and S2).

Figure 2. LD analysis of SNPs on 21q22.2 and 7q32.3 by pairwise r^2 . The 6 IGSF-5 related SNPs are within very tight linkage disequilibrium (LD) region.



For the additional susceptibility evaluation cohorts, samples were genotyped on Illumina Infinium SNP arrays, ranging from the Hap300 (for VQ58) to the Hap1M (for Australia). Details about genotyping and quality control for these studies have been provided previously (5). Ethics Committees approved these five studies and samples were collected in accordance with the tenets of the Declaration of Helsinki.

Among the top 19 SNPs identified from the APC trial, 12 SNPs had genotype data available from the CORGI, Scotland, VQ58, or Australia GWAS. Nine of these SNPs were typed in all four studies (rs1381392, rs17651822, rs17781398, rs16909065, rs9582985, rs2837156, rs2837241, and rs741864) and three were typed in three (rs13085889, rs1424593, and rs2837237; Supplementary Table S1).

Statistical analysis

To assess the strength of association between genotype and advanced-adenoma risk, a per allele unconditional logistic regression model was used to estimate ORs and their corresponding 95% confidence intervals (CI). For the APC trial, genotype–phenotype interactions were evaluated for sex, age at trial entry (\leq age 60 years vs. $>$ age 60 years), and family history (first-degree relatives with colorectal cancer). Genotype–environment interactions were evaluated for aspirin use at baseline and treatment with celecoxib. The Breslow–Day test was used to test the homogeneity of ORs. PLINK (<http://pngu.mgh.harvard.edu/~purcell/plink/>), STATA, and SAS were used to conduct all the

analysis. Genome-wide significance was defined as false discovery rate less than 0.05, which corresponds to an unadjusted $P = 7.4 \times 10^{-7}$ in this analysis (8, 9).

Results

Of note, 1,406 evaluable Caucasian patients were genotyped from the APC trial (4). Eight SNPs were identified by association with on-study development of advanced adenomas at a genome-wide level of significance: rs2837156, rs7278863, rs2837237, rs2837241, rs2837254, rs741864, all at 21q22.2, and rs1381392 and rs17651822, both at 3p24.1 (Table 1). The associations between the six SNPs in the 21q22.2 region and advanced adenoma development were all highly significant (unadjusted $P = 10^{-8}$ – 10^{-9}) with ORs per allele ranging from 2.22 to 2.55. All six SNPs in the 21q22.2 region were located near the coding region for the adherens junction protein, *IGSF5*, and five of these SNPs (rs7278863, rs2837237, rs741864, rs741864, and rs2837241) were in strong linkage disequilibrium ($r^2 = 0.8$ – 1 ; Fig. 1). For the 3p24.1 signal, the OR for genotype rs1381392 was 2.01 (95% CI, 1.52–2.65; unadjusted $P = 7.4 \times 10^{-07}$), and that for rs17651822 was 2.16 (95% CI, 1.61–2.91; unadjusted $P = 2.1 \times 10^{-7}$).

Eleven SNPs (rs11886781 at 2p24.2, rs13085889 at 3q22.2, rs1424593, rs1364512, and rs7778725 at 7q32.3, rs16909065 and rs16909036 at 9q33.2, rs17654765, rs1535989, and rs9582985 at 13q33.2) were associated with moderate (\sim 2-fold) ORs for advanced adenoma detection, but the associations did not reach genome-wide significance ($P \leq 10^{-6}$). Of these 11 SNPs,

Table 1. APC trial advanced adenoma susceptibility loci^a

Chromosome region	SNP	Position (bp)	Alleles	MAF	P	OR	Gene
2p24.2	rs11886781	18154780	A C	0.08	9.7E-06	2.25 (1.56–3.25)	KCNS3
3q22.2	rs13085889	135843760	A C	0.29	8.8E-06	1.77 (1.37–2.29)	EPHB1, KY
3p24.1	rs1381392	28724318	A G	0.18	7.4E-07	2.01 (1.52–2.65)	
3p24.1	rs17651822	28695130	A G	0.14	2.1E-07	2.16 (1.61–2.91)	
7p14.3	rs17781398	30807966	A G	0.10	9.0E-06	0.19 (0.08–0.43)	FAM188b
7q32.3	rs1424593	131605541	C A	0.50	9.1E-06	0.56 (0.44–0.73)	PLXNA4
7q32.3	rs1364512	131602384	C A	0.49	8.6E-06	0.56 (0.42–0.71)	PLXNA4
7q32.3	rs7778725	131614936	G A	0.49	4.0E-06	0.55 (0.42–0.71)	PLXNA4
9q33.2	rs16909065	121597606	A G	0.05	3.6E-06	2.59 (1.71–3.93)	
9q33.2	rs16909036	121587049	G A	0.05	3.7E-06	2.59 (1.71–3.93)	
13q33.2	rs1535989	104820723	G A	0.11	8.9E-06	2.09 (1.50–2.91)	
13q33.2	rs17654765	104828038	A G	0.10	4.7E-06	2.14 (1.53–2.98)	
13q33.2	rs9582985	104829133	C A	0.11	9.3E-06	2.05 (1.48–2.83)	
21q22.2	rs2837156	40048557	G A	0.12	3.2E-07	2.22 (1.62–3.03)	IGSF5
21q22.2	rs7278863	40087578	A G	0.10	1.4E-08	2.48 (1.80–3.42)	IGSF5
21q22.2	rs2837237	40119727	G A	0.12	3.6E-09	2.48 (1.82–3.38)	
21q22.2	rs2837241	40130476	A C	0.12	3.7E-09	2.48 (1.82–3.38)	
21q22.2	rs2837254	40143171	A G	0.11	2.9E-09	2.55 (1.86–3.51)	
21q22.2	rs741864	40129665	A G	0.11	1.1E-08	2.48 (1.80–3.41)	

^aThe total number of subjects is 1,406, of which 139 developed advanced adenomas.

Table 2. Meta-analysis using adenoma or colorectal cancer as a composite outcome

SNP	N	P	P(R)	OR	OR(R)	Q	I
rs13085889	4	0.1048	0.1048	1.0575	1.0575	0.5438	0
rs1381392	4	0.4636	0.4636	1.0295	1.0295	0.4186	0
rs1424593	3	0.405	0.405	1.027	1.027	0.4233	0
rs1535989	4	0.012	0.012	1.1304	1.1304	0.7925	0
rs16909065	4	0.1273	0.1273	0.9054	0.9054	0.498	0
rs17651822	4	0.689	0.689	1.0176	1.0176	0.8499	0
rs17781398	4	0.7972	0.7972	1.016	1.016	0.9487	0
rs2837156	4	0.8017	0.8017	0.9881	0.9881	0.4326	0
rs2837210	4	0.8706	0.9766	1.0083	0.9983	0.3114	16.05
rs2837237	3	0.3464	0.3464	0.9379	0.9379	0.413	0
rs2837241	4	0.938	0.8585	0.9963	0.9907	0.3395	10.69
rs741864	4	0.6248	0.6248	0.971	0.971	0.6068	0
rs9582985	4	0.05468	0.05468	1.1188	1.1188	0.9416	0

six mapped to gene coding regions: rs11886781 to *KCNS3*, rs17781398 to *FAM188b*, rs13085889 to *EPHB1* and *KY*, and rs1424593, rs1364512, and rs7778725 all to *PLXNA4* (Table 1 and Fig. 1).

There are no comparable adenoma chemoprevention cohorts currently available for validation of the APC GWAS results. We therefore further examined APC trial results using GWAS data from four nonoverlapping colorectal cancer case-control series of European ancestry, one of which (CORGI) also included advanced adenoma cases (5). Among the 19 advanced-adenoma risk SNPs with a nominal significance level of $P \leq 10^{-6}$, 12 were genotyped in at least three of the available four colorectal cancer GWA studies (Supplementary Table S1). Allelic frequencies of each variant and the corresponding associations with colorectal cancer phenotype were accessed in each of the four case-control samples. The results of the meta-analysis for overall associations with colorectal cancer risk are reported in Table 2 and Supplementary Fig. S3.

One of the 19 SNPs identified in the APC trial, rs1535989, was replicated in the independent colorectal cancer cohorts, with an OR for colorectal cancer development of 1.12 (95% CI, 1.019–1.23; $P = 0.019$). There was no evidence of inter-study heterogeneity ($P_{\text{het}} = 0.71$; $I^2 = 0.0\%$). An additional exploratory meta-analysis was performed, combining all five studies and using either advanced adenoma or colorectal cancer as the outcome (Table 2). SNP rs9582985 originally identified in the APC cohort showed marginally significant association with outcome (OR, 1.11; $P = 0.055$).

Clinical data from the APC trial were used to further characterize rs1535989 by examining the association of this variant with other susceptibility factors for advanced colorectal neoplasia, including age, sex, aspirin use at baseline, family history of colorectal cancer, and on-study treatment with celecoxib. SNP and environmental factors interaction terms were included in the model. SNP rs1535989 showed statistically significant interactions with subjects' age ($P = 0.0016$), sex ($P = 0.0057$), and aspirin use at baseline

($P = 0.02$). The associations with advanced neoplasia were stronger in older individuals (>60; OR, 3.20; 95% CI, 2.10–4.87), males (OR, 2.74; 95% CI, 1.89–3.97), and those using aspirin at baseline (OR, 3.63; 95% CI, 2.06–6.40; Table 3). There were no statistically significant interactions with colorectal cancer family history or on-study treatment with celecoxib.

Discussion

Among the approximately 145,000 colorectal cancer cases diagnosed per year in the United States, only 5% represent autosomal dominant predisposition syndromes, with the majority of these involving either hereditary non-polyposis colon cancer (HNPCC) or familial adenomatous polyposis (FAP). An additional 20% to 25% of colorectal cancer cases show a familial association without precise genetic characterization, and the majority of colorectal cancers occur in individuals without a family history of the disease. Self-reported family history does not accurately

Table 3. Genotype-phenotype/environment interactions for SNP rs1535989

Phenotype		OR	Interaction (P)
Age	≤60	0.91 (0.42–1.76)	0.0016
	>60	3.20 (2.10–4.87)	
Sex	Female	0.65 (0.25–1.68)	0.0057
	Male	2.74 (1.89–3.97)	
Family history	No	2.23 (1.51–3.28)	0.53
	Yes	1.74 (0.88–3.43)	
Prior aspirin use	No	1.58 (1.03–2.42)	0.02
	Yes	3.63 (2.06–6.40)	
Treatment	Placebo	1.63 (1.03–2.58)	NS
	200 mg	2.51 (1.27–4.95)	
	400 mg	2.43 (1.16–5.07)	

assess the inherited risk of advanced adenomas because patients' knowledge of their family history of colorectal adenomas is often unknown or incomplete (10). Recent GWAS from members of this collaboration have identified 18 colorectal cancer susceptibility variants with minor allele frequencies ranging from 0.07 to 0.48 that each convey a small degree of risk modification (OR per allele, 0.87–1.35; refs. 11–15). The results presented here expand these data to address inherited susceptibility for developing advanced adenomas that represent targets for colorectal cancer prevention. In addition to the studies whose data were used here, there have been a number of other GWAS with colorectal cancer or colorectal adenomas as the primary phenotype (16–20). These have yielded a substantial number of possible susceptibility variants, most conveying modestly altered risk. A recent case-control meta-analysis from 14 studies identified SNPs on 2q32.3 (rs11903757), 1q25.3 (rs10911251), 12p13.32 (rs3217810), and 12q24.21 (rs59336) that represented ORs ranging from 0.84 to 1.15 (16).

The APC trial was designed to determine whether the selective cyclooxygenase-2 inhibitor, celecoxib, prevented adenomas in patients at high risk for colorectal cancer. Eligibility criteria required that participants have at least one prior adenoma >6 mm in size, or multiple adenomas. During the 5 years of endoscopic surveillance, 21.3% of APC trial participants randomized to placebo developed recurrent advanced adenomas (21). This rate was decreased to 12.5% in patients receiving celecoxib 200 mg twice daily ($P < 0.0001$), however, concerns over cardiovascular toxicity currently prohibit the use of celecoxib for routine colorectal cancer chemoprevention (22). Results presented here showed that advanced adenomas were twice as likely to occur in APC trial participants with variant rs1535989, and that this increased risk was not affected by celecoxib treatment. For males or older individuals, the risk was more than 3-fold higher than that for females or participants younger than 60. The observed interaction between baseline aspirin use and advanced-adenoma risk is particularly interesting. Aspirin use reduces the incidence of colorectal adenomas and colorectal cancer, and subjects enrolled in the APC trial who used aspirin at baseline were those who developed adenomas despite aspirin use. These individuals may therefore have constituted a higher risk subset because they were relatively resistant to aspirin chemoprevention. The analyses conducted here showed that APC trial participants who both developed adenomas while taking aspirin and had variant rs1535989 demonstrated a 3.63-fold increase in advanced-adenoma risk during surveillance. If this association can be confirmed in other studies, and other variants of similar effect are found, then genotyping will represent a useful method to target high-risk patients for preventive treatments including more frequent colonoscopic screening.

Additional results from this GWAS suggest areas for further research about the molecular basis of colorectal neoplasia. Variants at rs2837156, rs7278863, rs2837237, rs2837241, rs2837254, and rs741864 are in close associa-

tion at 21q22.2 (Fig. 2). Two of these SNPs, rs2837156 and rs7278863, are within the coding region of *IGSF5*, a gene encoding a transmembrane protein whose murine homolog, *JAM4*, binds to the tumor suppressor *MAGI-1* at intestinal epithelial tight junctions (23). To explore the potential effect of *IGSF5* on the prognosis of patients with colorectal cancer, microarray expression data from the GEO data set (GSE14333) were retrieved for 229 Dukes A, B, and C patients (24). The gene expression profile was performed with Affymetrix u133p2 platform. Our preliminary analysis of these data indicated that the overexpression of *IGSF5* is associated with significantly worse relapse-free survival (unadjusted $P = 0.000004$; bonferroni adjusted $P = 0.00085$; Supplementary Fig. S4). In addition, rs1424593, rs1364512, and rs7778725 all involve *PLXNA4*, a member of the plexin family located on chromosome 7. Plexins are transmembrane, secreted, and GPI-anchored semaphorins that modulate the adhesive and migratory properties of malignant cells. The protein product of *PLXNA4* forms stable complexes with *FGFR1* and *VEGFR-2* tyrosine kinase receptors and enhances both VEGF-induced *VEGFR-2* phosphorylation and β FGF-induced cell proliferation (25). Finally, *EPHB1* encodes a ligand that binds to an Eph receptor tyrosine kinase to mediate bidirectional signaling required for intestinal epithelial homeostasis (26). EphB-ephrin B interactions regulate cell adhesion, migration, and positioning, and play an important role in colorectal tumor progression (27).

The limitation of our current study resides in the following two aspects. The limited number of advanced adenoma cases in the APC trial restricted the power to identify more advanced adenoma susceptibility SNPs. In addition, the colorectal cancer/adenoma cases with a strong family history of colorectal cancer were not excluded from replication datasets. This might under/overestimate the association between identified SNPs and sporadic colorectal-cancer risks.

In summary, this study identified 19 SNPs associated with advanced-adenoma risk at a level of $P \leq 10^{-6}$. Of these, 12 SNPs were tested in a meta-analysis using independent datasets to evaluate their association with colorectal cancer development, and rs1535989 was also associated with increased risk of both advanced adenomas and colorectal cancer. In addition, eight of the variants identified in the APC trial mapped to coding regions of genes previously implicated in colorectal cancer progression, and warrant further study to confirm their role in modifying tissue-specific biologic function.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J. Wang, L.G. Carvajal-Carmona, A.G. Zauber, APC Trial Collaborators, M. Kubo, M.J. Ratain, I. Tomlinson, M.M. Bertagnolli

Development of methodology: J. Wang, A.G. Zauber, APC Trial Collaborators, M.M. Bertagnolli

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): L.G. Carvajal-Carmona, APC Trial Collaborators, M. Kubo, K. Matsuda, M. Dunlop, R.S. Houlston, O. Sieber, L. Lipton, P. Gibbs,

N.G. Martin, G.W. Montgomery, J. Young, P.N. Baird, Y. Nakamura, S.T. Weiss, I. Tomlinson, M.M. Bertagnolli

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J. Wang, L.G. Carvajal-Carmona, J.-H. Chu, A.G. Zaubler, APC Trial Collaborators, M. Dunlop

Writing, review, and/or revision of the manuscript: J. Wang, L.G. Carvajal-Carmona, J.-H. Chu, A.G. Zaubler, APC Trial Collaborators, M. Dunlop, R.S. Houlston, O. Sieber, P. Gibbs, G.W. Montgomery, P.N. Baird, M.J. Ratain, S.T. Weiss, I. Tomlinson, M.M. Bertagnolli

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): J. Wang, APC Trial Collaborators, M. Kubo, N.G. Martin, G.W. Montgomery, M.J. Ratain, Y. Nakamura, M.M. Bertagnolli

Study supervision: J. Wang, M. Kubo, G.W. Montgomery, S.T. Weiss, M.M. Bertagnolli

Acknowledgments

The authors thank all the individuals who participated in the various studies. This study made use of genotyping data on the 1958 Birth Cohort and NBS samples, kindly made available by the Investigators of those studies and the Wellcome Trust Case-Control Consortium 2; a full list of the investigators who contributed to the generation of the data is available from <http://www.wtccc.org.uk>.

The CORGI acknowledgments and investigators: The CORGI study comprised: Huw Thomas, Family Cancer Clinic, St Mark's Hospital and Imperial College, London; Eamonn Maher, Dept. of Clinical Genetics, University of Birmingham, United Kingdom; Gareth Evans, Dept. of Clinical Genetics, University of Manchester, United Kingdom; Lisa Walker and Dorothy Halliday, Oxford Regional Genetics Service, Churchill Hospital, Oxford, United Kingdom; Anneke Lucassen, Wessex Regional Genetics Service, Princess Anne Hospital, Southampton, United Kingdom; Joan Paterson, Anglia Regional Genetics Service, Addenbrooke's Hospital, Cambridge, United Kingdom; Shirley Hodgson and Tessa Homfray, South-West Thames Regional Genetics Service, St George's Hospital, Tooting, London, United Kingdom; Lucy Side, North-East Thames Regional Genetics Service, Great Ormond Street Hospital, London, United Kingdom; Louise Izatt, South-East Thames Regional Genetics Service, Guy's Hospital, London, United Kingdom; Alan Donaldson and Susan Tomkins, South-West Regional Genetics Service, Bristol, United Kingdom; Patrick Morrison, Northern Ireland Regional Genetics Service, City Hospital, Belfast, United Kingdom; Carole Brewer, South-West Regional Genetics Service, Royal Devon and Exeter Hospital, Exeter, United Kingdom; Alex Henderson, Northern Regional Genetics Service, International Centre for Life, Newcastle, United Kingdom; Rosemarie Davidson and Victoria Munday, West of Scotland Regional Genetics Service, Yorkhill Hospital, Glasgow, United Kingdom; Jaqueline Cook, Sheffield Regional Genetics Service, Children's Hospital, Sheffield, United Kingdom; Neva Haites, North of Scotland Regional Genetics Service, Foresterhill Hospital, Aberdeen, United Kingdom; Timothy Bishop and Eamonn Sheridan, Yorkshire Regional Genetics Service, St James's Hospital, Leeds, United Kingdom; Andrew Green, Republic of Ireland Genetics Service, Our Lady's Hospital for Sick Children, Dublin, Republic of Ireland; Christopher Marks, Sue Carpenter and Mary Broughton, The Royal Surrey County Hospital, Egerton Road, Guildford, Surrey; Lynn Greenhalge, Department of Clinical Genetics, Royal Liverpool Children's Hospital, Eaton Road, Alder Hay, Liverpool; and Mohnish Suri, Department of Clinical Genetics, City Hospital, Hucknall Road, Nottingham.

APC acknowledgments and investigators: The following persons participated in the APC Study: Steering Committee: M.M. Bertagnolli, E.T. Hawk, C. J. Eagle; Statistical Team: A.G. Zaubler, K.M. Kim, D. Corle, R. Rosenstein, J. Tang, T. Hess, A. Wilton; Medical Monitors: W. Anderson, L. Doody; Central Pathology Review: M. Redston; Project Directors: G.M. Woloj, D. Bagheri, A. Crawford, M. Schietrum, V. Ladouceur; Data and Safety Monitoring Board: S. Rosen (chair), L. Friedman, R. Makuch, R. Phillips, P. Taylor; Principal Investigators, United States: S. Auerbach (California Professional Research, Newport Beach), C.F. Barish (Wake Research Associates, Raleigh, NC), T. Barringer (Carolinas Medical Center, Charlotte, NC), R.W. Bennetts (Northwest Gastroenterology Clinic, Portland, OR), M. Blitstein (Associates in Gastroenterology and Liver Disease, Lake Forest, IL), J. Bruggen (Wake Forest University Baptist Medical Center, Winston Salem, NC), P. Carricaburu (Veterans Affairs Hospital, Sheridan, WY), D. Chung (Massachusetts General Hospital, Boston, MA), F. Colizzo (Pentucket Medical Associates, Haverhill, MA), R. Curtis (Newton-Wellesley Hospital, Newton, MA), T. Dewar (Harris Methodist Hospital Fort Worth, Ft. Worth, TX), R. DuBois (Vanderbilt University Medical Center, Nashville, TN), T. Feinstat (Gastroenterology Consultants of Sacramento, Roseville, CA), T.R. Foley (Regional Gastroenterology Associates of Lancaster, Lancaster, PA), D. Gabbazadeh (Huntington Research Group, Huntington Station, NY), J. Geenen (Wisconsin Center for Advanced

Research, Milwaukee, WI), F. Giardiello (Johns Hopkins Hospital, Baltimore, MD), A. Goetsch (nTouch Research, Huntsville, AL), M. Goldberg (Regional Gastroenterology Associates of Lancaster, Evanston, IL), J.L. Goldstein (University of Illinois at Chicago, Chicago, IL), W. Harlan III (Asheville Gastroenterology Associates, Asheville, NC), R. Hogan (Gastrointestinal Associates, Jackson, MS), M. Kamionkowski (Gastroenterology Associates of Cleveland, Mayfield Heights, OH), M. Kelfer (Fallon Clinic, West Boylston, MA), B. Kerzner (Health Trends Research, Baltimore, MD), K. Kim (University of Chicago Medical Center, Chicago, IL), I. Klimberg (Gastroenterology Associates of Ocala, Ocala, FL), G. Koval (West Hills Gastroenterology Associates, Portland, OR), C. Krone (Advanced Clinical Therapeutics, Tucson, AZ), S. Krumholz (Waterside Clinical Research, West Palm Beach, FL), M.W. Layton (South Puget Sound Clinical Research Center, Olympia, WA), C. Lightdale (Columbia-Presbyterian Medical Center, New York, NY), P.J. Limburg (May Clinic, Rochester, MN), C. Lind (Vanderbilt University Medical Center, Nashville, TN), D. Lipkis (Institute for Health Care Assessment, San Diego, CA), M. Lloyd (Idaho Gastroenterology, Meridian, ID), D. Maccini (Spokane Digestive Disease Center, Spokane, WA), F. MacMillan Sr (Pentucket Medical Associates, Haverhill, MA), R. Madoff (University of Minnesota, Minneapolis, MN), A. Malik (Advanced Clinical Research, North Providence, RI), A. Markowitz (Memorial Sloan-Kettering Cancer Center, New York, NY), R. Marks (Alabama Digestive Research Center, Alabaster, AL), C. J. McDougall (Manhattan Associates, New York, NY), P. Miner (Oklahoma Foundation for Digestive Research, Oklahoma City, OK), M. Murphy (Southern Digestive and Liver Disease Institute, Savannah, GA), A. Namais (Gastrointestinal Physicians, Salem, MA), N. Nickl (University of Kentucky Medical Center, Lexington, KY), M. Pochapin (Jay Monahan Center for Gastrointestinal Health, New York, NY), R.E. Pruitt (Nashville Medical Research Institute, Nashville, TN), J. Puolos (Cumberland Research Associates, Fayetteville, NC), D.S. Riff (AGMG Clinical Research, Anaheim, CA), R. Roman (South Denver Gastroenterology, Englewood, CO), L. Rubin (New Jersey Physicians, Passaic, NJ), D. Ruff (Healthcare Discoveries, San Antonio, TX), M. Safdi (Consultants for Clinical Research, Cincinnati, OH), J. Saltzman (Brigham and Women's Hospital, Boston, MA), B. Salzberg (Atlanta Gastroenterology Associates, Atlanta, GA), J. A. Sattler (Western Clinical Research, Torrance, CA), P. Schleinitz (Americas Doctors Research, Medford, OR), J. Schwartz (Northwest Gastroenterologists, Arlington Heights, IL), M. Schwartz (Jupiter Research Association, Jupiter, FL), M. Silpa (Gastroenterology Associates of the East Bay Medical Group, Berkeley, CA), D. Silvers (Drug Research Services, Metairie, LA), D. Smoot (Howard University Cancer Center, Washington, DC), S. Sontag (Veterans Affairs Medical Center, Hines, IL), R.J. Sorrell (Gastroenterology Specialties, Lincoln, NE), D. Stanton (Community Clinical Trials, Orange, CA), J. Surgeon (Americas Doctors Research, Shawnee Mission, KS), J.P. Tracey (Hawthorne Medical Associates, North Dartmouth, MA), T. Werth (Charlotte Gastroenterology and Hepatology, Charlotte, NC), C.M. Wilcox (University of Alabama at Birmingham, Birmingham, AL), R. Wohlman (Northwest Gastroenterology Associates, Bellevue, WA), S. Woods (Gastroenterology Associates of Fairfield County, Bridgeport, CT); United Kingdom: J. Burn (South Cleveland Hospital, Middlesbrough); Australia: H. Ee (Sir Charles Gairdner Hospital, Nedlands, WA), M. Korman (Monash Medical Center, Clayton, Victoria), A. Lee (Concord Repatriation and General Hospital, Concord, NSW), B. Leggett (Royal Brisbane Hospital, Herston, Queensland), F. Macrae (Royal Melbourne Hospital, Melbourne, Victoria), L. Mollison (Freemantle Hospital, Freemantle, WA), N. Yeomans (Western Hospital, Footscray, Victoria), G. Young (Flinders Medical Center, Bedford, SA); Canada: G. Aumais (Hospital Maisonneuve-Rosemont, Montreal), R. Bailey (Hys Medical Center, Edmonton, Alberta), C. Bernstein (Winnipeg Health Sciences Centre, Winnipeg, Manitoba), L. Cohen (Sunnybrook and Women's Hospital, Toronto), C. Dallaire, R. Dube (Centre Hospitalier Universitaire de Quebec, Quebec), D. Morgan (McMaster University, Hamilton, Ontario), T. Sylwestrowicz (St. Paul's Hospital, Saskatoon, Saskatoon), G. Van Rosendaal (University of Calgary, Alberta), S.J. Van Zant (Queen Elizabeth II Health Sciences Centre, Halifax, NS).

Colon Cancer Family Registry acknowledgments and investigators: David Conti and Graham Casey, University of Southern California, Los Angeles; Polly Newcomb, Fred Hutchinson Cancer Research Center, Seattle; John Hopper and Mark Jenkins, University of Melbourne, Melbourne, Australia; Steven Gallinger, Cancer Care Ontario, Toronto, Ontario, Canada; David J. Duggan, Translational Genomics Research Institute, Phoenix, Arizona.

Grant Support

The APC study was funded by the U.S. National Cancer Institute (CA-N01-95015 and HHSN261201000082C) and by Pfizer. Cancer Research UK funded the CORGI, Scotland, and NSCCG studies. L.G. Carvajal-Carmona is supported by the EU FP7 CHIBCHA project. I. Tomlinson received support from the Oxford NIHR Comprehensive Biomedical Research Centre. Core infrastructure support to the Wellcome Trust Centre for Human Genetics, Oxford was provided by grant 090532/Z/09/Z. The United Kingdom

National Cancer Research Network supported the NSCCG. M. Dunlop received from the Medical Research Council (G0000657-53203), CORE and Scottish Executive Chief Scientist's Office (K/OPR/2/2/D333, CZB/4/449). The Colon Cancer Family Registry was supported by the National Cancer Institute, NIH under Request for Application #CA-95-011, and through cooperative agreements with the Australian Colorectal Cancer Family Registry (UO1 CA097735), the USC Familial Colorectal Neoplasia Collaborative Group (UO1 CA074799), Mayo Clinic Cooperative Family Registry for Colon Cancer Studies (UO1 CA074800), Ontario Registry for Studies of Familial Colorectal Cancer (UO1 CA074783), Seattle Colorectal Cancer Family Registry (UO1 CA074794), and The University of Hawaii Colorectal Cancer Family Registry (UO1 CA074806). P.N. Baird is supported by Australian National Health and Medical Research Council Senior Research Fellowship #1028444. CERA receives Operational Infrastructure Support from the Victorian Government, Australia.

This study was supported by NCI N01-95015 and HHSN26120100082C (to M.M. Bertagnolli, A.G. Zauber, and J. Wang), U01GM61393 (to M.J.

Raitan), U01GM61390, and the Biobank Japan Project funded by the Japanese Ministry of Education, Culture, Sports, Science and Technology. This work is part of the NIH Pharmacogenomics Research Network-RIKEN Center for Genomic Medicine Global Alliance. Pfizer, Inc. partially funded the APC Trial. L.G. Carvajal-Carmona and I. Tomlinson received support from Cancer Research UK and the Oxford Comprehensive Biomedical Research Centre, and the European Union (FP7 CHIBCHA Consortium). The Wellcome Trust Centre for Human Genetics is supported by a Wellcome Trust Core Grant 090532/Z/09/Z.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received March 2, 2013; revised September 21, 2013; accepted September 24, 2013; published OnlineFirst October 1, 2013.

References

- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012;62:10-29.
- Zauber AG, Winawer SJ, O'Brien MJ, Lansdorp-Vogelaar I, van Ballegoijen M, Hankey BF, et al. Colonoscopic polypectomy and long-term prevention of colorectal cancer deaths. *New Engl J Med* 2012;366:687-96.
- Saini SD, Kim HM, Schoenfeld P. Incidence of advanced adenomas at surveillance colonoscopy in patients with a personal history of colon adenomas: a meta-analysis and systematic review. *Gastrointest Endosc* 2006;64:614-26.
- Bertagnolli MM, Eagle CJ, Zauber AG, Redston M, Solomon SD, Kim K, et al. Celecoxib for the prevention of sporadic colorectal adenomas. *N Engl J Med* 2006;355:873-84.
- Carvajal-Carmona LG, Cazier J-B, Jones AM, Howarth K, Broderick P, Pittman A, et al. Fine-mapping of colorectal cancer susceptibility loci at 8q23.3, 16q22.1 and 19q13.11: refinement of association signals and use of *in silico* analysis to suggest functional variation and unexpected candidate target genes. *Hum Mol Genet* 2011;20:2879-88.
- Tomlinson IP, Carvajal-Carmona LG, Dobbins SE, Tenesa A, Jones AM, Howarth K, et al. Multiple common susceptibility variants near BMP pathway loci GREM1, BMP4, and BMP2 explain part of the missing heritability of colorectal cancer. *PLoS Genet* 2011;7:e1002105.
- Wellcome TCCC. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661-78.
- Benjamini YH, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B* 1995;57:12.
- Benjamini YH, Hochberg Y. The adaptive control of the false discovery rate in multiple hypotheses testing. *J Behav Educ Statist* 2000;25:14.
- Elias PS, Romagnuolo J, Hoffman B. Poor patient knowledge regarding family history of colon polyps: implications for the feasibility of stratified screening recommendations. *Gastrointest Endosc* 2012;75:598-603.
- Houlston RS, Cheadle J, Dobbins SE, Tenesa A, Jones AM, Howarth K, et al. Meta-analysis of three genome-wide association studies identifies susceptibility loci for colorectal cancer at 1q41, 3q26.2, 12q13.13 and 20q13.33. *Nat Genet* 2010;42:973-7.
- Houlston RS, Webb E, Broderick P, Pittman AM, Di Bernardo MC, Lubbe S, et al. Meta-analysis of genome-wide association data identifies four new susceptibility loci for colorectal cancer. *Nat Genet* 2008;40:1426-35.
- Tomlinson I, Webb E, Carvajal-Carmona L, Broderick P, Kemp Z, Spain S, et al. A genome-wide association scan of tag SNPs identifies a susceptibility variant for colorectal cancer at 8q24.21. *Nat Genet* 2007;39:984-8.
- Tomlinson IP, Webb E, Carvajal-Carmona L, Broderick P, Howarth K, Pittman AM, et al. A genome-wide association study identifies colorectal cancer susceptibility loci on chromosomes 10p14 and 8q23.3. *Nat Genet* 2008;40:623-30.
- Broderick P, Carvajal-Carmona L, Pittman AM, Webb E, Howarth K, Rowan A, et al. A genome-wide association study shows that common alleles of SMAD7 influence colorectal cancer risk. *Nat Genet* 2007;39:1315-7.
- Peters U, Jiao S, Schumacher FR, Hutter CM, Aragaki AK, Baron JA, et al. Identification of genetic susceptibility loci for colorectal tumors in a genome-wide meta-analysis. *Gastroenterology* 2013;144:799-807.
- Zanke BW, Greenwood CM, Rangrej J, Kustra R, Tenesa A, Farrington SM, et al. Genome-wide association scan identifies a colorectal cancer susceptibility locus on chromosome 8q24. *Nat Genet* 2007;39:989-94.
- Tenesa A, Farrington SM, Prendergast JG, Porteous ME, Walker M, Haq N, et al. Genome-wide association scan identifies a colorectal cancer susceptibility locus on 11q23 and replicates risk loci at 8q24 and 18q21. *Nat Genet* 2008;40:631-7.
- Jaeger E, Webb E, Howarth K, Carvajal-Carmona L, Rowan A, Broderick P, et al. Common genetic variants at the CRAC1 (HMPS) locus on chromosome 15q13.3 influence colorectal cancer risk. *Nat Genet* 2008;40:26-8.
- Dunlop MG, Dobbins SE, Farrington SM, Jones AM, Palles C, Whiffin N, et al. Common variation near CDKN1A, POLD3 and SHROOM2 influences colorectal cancer risk. *Nat Genet* 2012;44:770-6.
- Bertagnolli MM, Eagle CJ, Zauber AG, Redston M, Breazna A, Kim K, et al. Five-year efficacy and safety analysis of the adenoma prevention with celecoxib trial. *Cancer Prev Res* 2009;2:310-21.
- Solomon SD, McMurray JJ, Pfeffer MA, Wittes J, Fowler R, Finn P, et al. Cardiovascular risk associated with celecoxib in a clinical trial for colorectal adenoma prevention. *N Engl J Med* 2005;352:1071-80.
- Hirabayashi S, Tajima M, Yao I, Nishimura W, Mori H, Hata Y. JAM4, a junctional cell adhesion molecule interacting with a tight junction protein, MAGI-1. *Mol Cell Biol* 2003;23:4267-82.
- Jorissen RN, Gibbs P, Christie M, Prakash S, Lipton L, Desai J, et al. Metastasis-associated gene expression changes predict poor outcomes in patients with dukes stage B and C colorectal cancer. *Clin Cancer Res* 2009;15:7642-51.
- Kigel B, Rabinowicz N, Varshavsky A, Kessler O, Neufeld G. Plexin-A4 promotes tumor progression and tumor angiogenesis by enhancement of VEGF and bFGF signaling. *Blood* 2011;118:4285-96.
- Battle E, Bacani J, Begthel H, Jonkheer S, Gregorieff A, van de Born M, et al. EphB receptor activity suppresses colorectal cancer progression. *Nature* 2005;435:1126-30.
- Cortina C, Palomo-Ponce S, Iglesias M, Fernandez-Masip JL, Vivanco A, Whissell G, et al. EphB-ephrin-B interactions suppress colorectal cancer progression by compartmentalizing tumor cells. *Nat Genet* 2007;39:1376-83.

Clinical Cancer Research

Germline Variants and Advanced Colorectal Adenomas: Adenoma Prevention with Celecoxib Trial Genome-wide Association Study

Jiping Wang, Luis G. Carvajal-Carmona, Jen-Hwa Chu, et al.

Clin Cancer Res Published OnlineFirst October 1, 2013.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-13-0550](https://doi.org/10.1158/1078-0432.CCR-13-0550)

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.