

Phase I Trial of ¹³¹I-huA33 in Patients with Advanced Colorectal Carcinoma

Geoffrey Chong,¹ Fook Thean Lee,¹ Wendie Hopkins,¹ Niall Tebbutt,¹ Jonathan S. Cebon,¹ Angela J. Mountain,¹ Bridget Chappell,² Anthony Papenfuss,² Paul Schleyer,¹ Paul U,² Roger Murphy,¹ Veronika Wirth,¹ Fiona E. Smyth,¹ Nicole Potasz,¹ Aurora Poon,² Ian D. Davis,¹ Tim Saunder,² Graeme J. O'Keefe,² Antony W. Burgess,¹ Eric W. Hoffman,³ Lloyd J. Old,³ and Andrew M. Scott^{1,2}

Abstract Purpose: Humanized monoclonal antibody A33 (huA33) targets the A33 antigen which is expressed on 95% of colorectal cancers. A previous study has shown excellent tumor-targeting of iodine-131 labeled huA33 (¹³¹I-huA33). Therefore, we did a phase I dose escalation trial of ¹³¹I-huA33 radioimmunotherapy.

Experimental Designs: Fifteen patients with pretreated metastatic colorectal carcinoma each received two i.v. doses of ¹³¹I-huA33. The first was an outpatient trace-labeled "scout" dose for biodistribution assessment, followed by a second "therapy" dose. Three patients were treated at 20, 30, and 40 mCi/m² dose levels, and six patients at 50 mCi/m² to define the maximum tolerated dose.

Results: Hematologic toxicity was ¹³¹I dose-dependent, with one episode of grade 4 neutropenia and two episodes of grade 3 thrombocytopenia observed at 50 mCi/m². The maximum tolerated dose was determined to be 40 mCi/m². There were no acute infusion-related adverse events, and gastrointestinal toxicity was not observed despite uptake of ¹³¹I-huA33 in bowel. Seven patients developed pruritus or rash, which was not related to ¹³¹I dose. There was excellent tumor-targeting of ¹³¹I-huA33 shown in all patients. The serum T1/2β of ¹³¹I-huA33 was (mean ± SD) 135.2 ± 46.9 hours. The mean absorbed tumor dose was 6.49 ± 2.47 Gy/GBq. Four patients developed human anti-human antibodies. At restaging, 4 patients had stable disease, whereas 11 patients had progressive disease.

Conclusion: Radioimmunotherapy using ¹³¹I-huA33 shows promise in targeting colorectal tumors, and is deliverable at a maximum tolerated dose of 40 mCi/m². Further studies of ¹³¹I-huA33 in combination with chemotherapy are planned.

Radioimmunotherapy is coming of age as a treatment modality for cancer. Patients with hematologic malignancies have been the first to benefit from treatment with radiolabeled monoclonal antibodies (mAb; refs. 1, 2). Yet, there is much potential for the application of radioimmunotherapy to epithelial malignancies as well. Colorectal cancer is a significant public health problem and is the leading cancer by incidence in Australia and the U.S. (3). Unfortunately, ~50%

of all patients develop metastatic disease, for which curative treatment is not yet available (4).

The A33 antigen is a 43 kDa membrane bound glycoprotein present on the basolateral surfaces of normal colon and small bowel epithelial cells in mice and humans (5–7). It is also homogeneously expressed in >95% of human colorectal cancers and ~50% of gastric cancers (5). Approximately 50% of pancreatic cancers show heterogeneous expression. The A33 antigen is not detectably shed into the extracellular space or the bloodstream, in contrast to other tumor antigens such as carcinoembryonic antigen.

The A33 antigen has been sequenced and comprises three structural domains: an extracellular region containing two immunoglobulin-like domains, a hydrophobic transmembrane domain, and a highly polar intracellular tail (6–9). It is therefore regarded as part of a subfamily within the immunoglobulin superfamily, together with the transmembrane proteins CTX (corticthymocyte marker in *Xenopus*), CTM/CTH (mouse and human homologues of CTX) and CAR (Coxsackie and adenovirus receptor; ref. 7). Its structure is consistent with a putative role as a cell adhesion molecule or a novel cell surface receptor.

Previous radioimmunotherapy studies using iodinated murine A33 mAb (muA33) have shown the therapeutic potential of the A33 antigen-antibody system in patients with metastatic

Authors' Affiliations: ¹Ludwig Institute for Cancer Research, ²Departments of Nuclear Medicine and Centre for Positron Emission Tomography, Austin Hospital, Melbourne, Australia and ³Ludwig Institute for Cancer Research, New York, New York

Received 11/14/04; revised 1/28/05; accepted 1/31/05.

Grant support: National Health and Medical Research Council Program grant nos. 280912 and 98/1905.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Andrew M. Scott, Tumour Targeting Program, Ludwig Institute for Cancer Research, Level 1, Harold Stokes Building, Austin Hospital, 143-165 Studley Road, Heidelberg, Victoria 3084, Australia. Phone: 61-39496-5876; Fax: 61-39496-5892; E-mail: andrew.scott@ludwig.edu.au.

©2005 American Association for Cancer Research.

colorectal cancer (10–12). A phase I/II study of treatment doses of ¹³¹I-muA33 enrolled 23 patients with advanced unresectable colon cancer. Reversible hematologic toxicity was dose-limiting and the maximum tolerated dose was determined to be 75 mCi/m². No significant bowel toxicity was observed. Minor evidence of tumor response was seen, with three patients demonstrating mixed responses to treatment. Another phase I trial examined the effects of escalating doses of ¹²⁵I-muA33 in patients with advanced colon cancer (12). Twenty-one patients were enrolled and treated with ¹²⁵I-muA33 at doses ranging from 50 to 350 mCi/m². Dose-limiting toxicity was not observed and maximum tolerated dose was therefore not reached. One patient had a mixed tumor response and 12 patients had stable disease. This series of muA33 radioimmunotherapy trials showed safety, tolerability, and specific tumor-targeting in patients with advanced colorectal cancer. Hints of antitumor efficacy were also seen, justifying further clinical studies; however, the development of human anti-mouse antibodies in all patients prevented repeat treatment. In order to reduce immunogenicity, a CDR grafted, humanized mAb (huA33) was therefore developed (13). Clinical trials using multiple dose schedules of huA33 have shown safety and possible efficacy of huA33 mAb alone and combined with chemotherapy (14, 15).

As a prelude to ¹³¹I-huA33 radioimmunotherapy, a phase I pharmacokinetic and tumor-targeting study accrued patients with colorectal carcinoma who were scheduled for surgical resection of one or more tumors. One week prior to surgery, patients were infused with ¹³¹I and ¹²⁵I labeled huA33. Surgical tumor biopsies were analyzed for ¹³¹I-huA33 uptake and microdistribution. Favorable biodistribution and tumor-targeting data were obtained,⁴ thus a phase I trial using radioimmunotherapy doses of ¹³¹I-huA33 was conducted and is reported here.

Patients and Methods

Patients. Patients with metastatic colorectal cancer who had received at least one line of chemotherapy for metastatic disease were eligible for this trial if they fulfilled all of the following inclusion criteria: histologically proven colorectal cancer; measurable or evaluable disease, Karnofsky performance status $\geq 70\%$; expected survival ≥ 4 months; serum creatinine < 0.15 mmol/L; serum bilirubin < 34 μ mol/L; neutrophil count $> 1.5 \times 10^9$ /L; platelet count $> 150 \times 10^9$ /L; prothrombin time $< 1.3 \times$ control; age ≥ 18 years; ability to give written informed consent. Patients who fulfilled any of the following exclusion criteria were ineligible for study treatment: active central nervous system metastases; exposure to chemotherapy, radiotherapy or immunotherapy within 4 weeks of study entry; not fully recovered from surgery; metastatic disease involving $> 50\%$ of liver mass; positive huA33 human anti-human antibodies (HAHA) titer in the context of previous huA33 treatment; serious concurrent illness; pregnancy or lactation; concurrent treatment with systemic corticosteroids/immunosuppressants. All patients gave witnessed written informed consent to participate in this study. The protocol was approved by the Human Research Ethics Committee of the Austin Hospital, Melbourne, Australia.

⁴ Scott AM, Lee FT, Jones R, Hopkins W, MacGregor D, Cebon JS, Hannah A, Chong G, U P, Papenfuss A, Rigopoulos A, Sturrock S, Murphy R, Wirth V, Murone C, Smyth FE, Knight S, Welt S, Ritter G, Richards E, Nice EC, Burgess AW, Old LJ. Phase I biopsy-based study of humanised monoclonal antibody A33 in patients with colorectal carcinoma. Clin Can Res, in press.

A phase I dose escalation design was used where only the therapy ¹³¹I dose was escalated between dose levels; the huA33 protein dose was held constant at 10 mg/m². Three patients were initially treated per ¹³¹I dose level. Toxicity was defined as adverse events which were possibly, probably, or definitely related to the study agent, i.e., ¹³¹I-huA33. Dose-limiting toxicity was defined as grade 3 nonhematologic toxicity, or grade 4 hematologic toxicity as defined by the National Cancer Institute Common Toxicity Criteria version 2.0, 1999. If dose-limiting toxicity was experienced by any patient at a particular dose level, three more patients were to be accrued to that dose level. The maximum tolerated dose was defined as the ¹³¹I dose below that where two or more patients out of six experienced dose-limiting toxicities.

Within 2 weeks of study treatment, pretreatment assessments were done to confirm eligibility. Included were complete medical history and physical examination, documentation of measurable or evaluable tumor by computerized tomography, serum biochemistry, and hematology. A baseline serum sample for HAHA determination was taken in all patients. Assessments of cardiorespiratory function were also done as a baseline for monitoring of possible delayed radiotoxicity.

All patients commenced treatment with an i.v. trace-labeled "scout" dose of ¹³¹I-huA33 in order to assess tumor-targeting, dosimetry, and biodistribution. All patients received 5 mg (fixed dose) huA33 conjugated to 5 to 8 mCi (200–320 MBq) ¹³¹I, using an established labeling technique as previously described (16). Each patient was directly observed for evidence of anaphylactoid reactions during the entire 1-hour infusion period and for 1 hour after completion of the infusion. Vital signs were measured once during and immediately following the infusion, and 2, 3, and 4 hours after the completion of infusion. All patients received saturated solution of potassium iodide prior to the scout ¹³¹I-huA33 infusion and continued for 3 weeks in total at 10 drops orally thrice daily.

Each patient received an i.v. therapy dose of ¹³¹I-huA33 7 ± 2 days following the scout infusion. Patients were hospitalized in a self-contained radiation-shielded room for this dose. All patients received 10 mg/m² huA33 conjugated to 20, 30, 40, or 50 mCi/m² ¹³¹I depending on assigned dose level. Automatic monitoring of blood pressure, pulse, and oxygen saturation was done during this time using a programmable patient monitoring device (Agilent Technologies, Andover, MA). Patients remained confined to this room until emitted radiation at 2 m fell to < 25 μ Sv/hour or $< 25\%$ of the initial reading. Geiger counter measurements were done daily while patients were hospitalized. All radiation safety precautions were fully compliant with Australian Radiation Protection and Nuclear Safety Agency guidelines and were supervised by the Radiation Safety Officer of the Austin Hospital.

For 6 weeks following the therapy dose, patients were reviewed weekly in order to conduct assessments of adverse events and to determine their relationship, if any, to the study agent. Blood was drawn at each weekly visit for biochemistry, hematology, and HAHA measurement. Clinical and investigational adverse events were recorded in case report forms and were coded according to the Common Toxicity Criteria.

Six weeks after administration of the therapy ¹³¹I-huA33 dose, an end of study evaluation for each patient was done. This included clinical assessment of adverse events, as well as clinical and radiological tumor reassessment. Tumor response or progression was defined according to WHO criteria.

In order to detect possible delayed toxicity from ¹³¹I-huA33 therapy, all patients were followed, where possible, at 3, 6, and 12 months following completion of the initial study period. Follow up assessments included clinical assessment, biochemical assessment of renal and thyroid function, chest X-rays, respiratory function tests, and gated cardiac blood pool scans.

Pharmacokinetics. Blood samples for analysis of pharmacokinetics of ¹³¹I-huA33 (scout infusion) were drawn immediately prior to the infusion (day 0), 5 minutes after completion of ¹³¹I-huA33 infusion, and 1, 2, 4, and 20 ± 2 hours (days 2 or 3, 4 or 5, and prior to the therapy infusion). The serum obtained was aliquoted in duplicate and counted in a gamma scintillation counter (Packard Instruments, Melbourne,

Australia). Triplicate standards prepared from the injected material were counted for ^{131}I at each time point with serum samples to enable calculations to be corrected for the isotope's physical decay. The results of the serum were expressed as the percentage of injected dose per liter (%ID/L) and converted to micrograms per milliliter based on actual protein dose infused. Population pharmacokinetics used normalized %ID/L (expressing subsequent %ID/L values as a percentage of the initial value for each patient) in order to compare clearance patterns between different patients.

Pharmacokinetic calculations were done for serum data using a curve-fitting program (WinNonlin, Pharsight Co., Mountain View, CA). A two-compartment model was used to calculate pharmacokinetic results. The following variables were calculated from the model: C_{\max} (maximum serum concentration); AUC (area under the serum concentration curve extrapolated to infinite time); CL (total serum clearance); $T_{1/2\alpha}$ and $T_{1/2\beta}$ (half lives of the initial and terminal phases of disposition); and V_1 (central compartment distribution).

Biodistribution of ^{131}I -huA33. Whole-body gamma camera scans were done 1 to 4 hours post-scout ^{131}I -huA33 infusion, and then on days 1, 2 or 3, and 4 or 5. A standard of known activity was included in the field of view to allow quantitation. Single-photon emission computed tomography imaging of areas of known tumor was done on day 4 or 5. Analysis of biodistribution images was done following the scout ^{131}I -huA33 dose to ensure that no abnormal distribution of ^{131}I -huA33 had occurred prior to administration of the therapy ^{131}I -huA33 dose.

Whole-body clearance and dosimetry of ^{131}I -huA33. Dosimetric analysis was done on the series of gamma camera whole-body planar images acquired in all patients following scout infusion. Whole body, lungs, liver, spleen, kidney, and thyroid regions of interest were defined for each time point on both anterior and posterior images. Organ radioactivity content was estimated from the geometric mean of anterior and posterior regions of interest counts. The counts for each organ were corrected for background using regions of interest drawn adjacent to each organ. Correction for attenuation of individual organs was estimated using an analytic technique as described by Liu et al. (17). Resultant counts were converted to activity using a camera sensitivity factor calculated from a gamma camera standard of known activity which was scanned at the same time.

A time-activity curve was generated from the derived activity of each imaging time point for each organ. The time-activity curves were decay-corrected to time of injection and fitted with a single component exponential clearance expression. These fits were used to determine biological clearance half-life, T_{bio} , and initial fraction of injected activity, A_0 , for the whole body and various normal organs. The AUC was analytically determined and expressed as the residence time of the organ and remainder tissue (18). Residence times for the lungs, liver, spleen, kidney, thyroid, and remainder tissue were entered into the MIRDOSE3 computer software program (19) to calculate the estimated radiation absorbed dose to these specific organs per unit of injected activity (mGy/MBq). This value was multiplied by the total activity delivered to the patient to determine the absorbed radiation dose for each organ.

For tumor dosimetry calculations, a region of interest was drawn around all of or part of the tumor and a time-activity curve derived using similar methodology to normal organ dose. Tumor volumes were calculated from computerized tomography scans. Due to prolonged retention of ^{131}I -huA33 in tumor, additional imaging was done up to 28 days following therapy infusion in 5 of 15 patients to more accurately determine tumor uptake and dose. Tumor mean absorbed dose was calculated as follows:

$$\bar{D} = \bar{A}S$$

where \bar{D} , mean absorbed dose to target; \bar{A} , cumulated activity in target; S , absorbed dose in target per unit cumulated activity.

For ^{131}I , S values were obtained from predefined tables corresponding to the calculated tumor mass [MIRDOSE3, Nodule Module; ref. (19)].

To extrapolate therapy absorbed dose to tumor, the scout absorbed dose was normalized by the injected activity and then multiplied by the therapy injected activity.

Red marrow dosimetry calculations. Calculation of red marrow dose for ^{131}I -huA33 was done using a patient-specific marrow dosimetry methodology (20). (A) Serum clearance calculations of ^{131}I -huA33 were used to calculate serum cumulated activity concentration, and marrow cumulated activity concentration ($\mu\text{Ci} - \text{h/gm}$). (B) Whole body cumulated activity concentration ($\mu\text{Ci} - \text{h/gm}$) was calculated as previously described. The calculation of whole-body clearance for scout doses for red marrow dose was based on imaging data. As there were fewer imaging time points following the therapy infusion, red marrow-absorbed doses for therapy infusions were calculated using the whole body clearance determined from handheld dose rate monitor readings (UIMO LB 123, EG&G Berthold, Germany).

Red marrow absorbed dose (rad/mCi) was then calculated with the formula (ref. 20):

$$\frac{\text{Red marrow absorbed dose (rad/mCi)}}{\text{injected dose (mCi)}} = (0.313 \times A + 0.456 \times B)$$

Biacore HAHA analyses. Blood samples for HAHA assessment were taken prior to each ^{131}I -huA33 infusion, then at weekly intervals until the end-of-study visit (6 weeks after therapy dose). Antibody responses against humanized antibodies (HAHA) induced after treatment of patients with huA33 mAb were analyzed by surface plasmon resonance technology using a Biacore 2000 instrument as previously described (21–23). Patient serum was considered HAHA-positive if the RU value at a serum dilution of 1:100 exceeded a cutoff value, defined as the mean interpatient baseline RU value +3 SD of pretreatment sera at a serum dilution of 1:100, provided that such an increase was absent in the isotope-matched control antibody channel (24).

Results

Fifteen patients were entered and treated according to the study protocol. Table 1 shows patient age, gender, assigned dose level, Karnofsky performance status and site of primary tumor. The median age was 63 (range, 39-77). The male/female ratio was 11:4. Median performance status was 90 (range, 80-100). Sites of metastatic disease ranged from liver, lung, retroperitoneal/mediastinal lymph nodes, pelvic, and peritoneum (Table 2). All patients had received previous therapy for metastatic colorectal cancer (Table 2). The median number of lines of previous chemotherapy was two (range, 1-5). Three patients had been previously exposed to humanized antibodies as part of clinical trials, but there was no evidence of HAHA to huA33 prior to study entry.

Toxicity. Overall, ^{131}I -huA33 was well-tolerated at dose levels up to the maximum tolerated dose. No acute infusion-related toxicities were observed. Hematologic toxicity was observed at all dose levels and increased with increasing dose level. Table 3 shows the maximum grade of hematologic toxicity by Common Toxicity Criteria experienced by each patient. Because of concerns about hematologic toxicity, in particular, thrombocytopenia, the 50 mCi/m² dose level was expanded to six patients. There was one episode of hematologic dose-limiting toxicity observed, which was grade 4 neutropenia in patient 14. This patient also experienced grade 3 thrombocytopenia and the longest time to complete hematologic recovery (10 weeks). No episodes of febrile neutropenia or thrombocytopenia requiring platelet transfusion were observed at any dose level.

Unexpected, but clinically significant, cutaneous toxicity was observed in a total of seven patients. The frequency of cutaneous events was similar at all ¹³¹I dose levels. Two patients (nos. 7 and 14) experienced grade 3 extensive pruritic, erythematous, urticaria-like rashes requiring treatment with systemic corticosteroids. There was no evidence that the rash was associated with ¹³¹I-dose; it was most likely due to A33 protein. In the other patients, symptoms related to rashes were treated with oral antihistamines or no specific treatment. There was no clear relationship between development of rash and/or pruritis and the development of HAHA. Table 4 shows the occurrence of cutaneous events, the presence of HAHA, and ¹³¹I-huA33 dose levels. Skin biopsies from patient 7 were taken 3 weeks post-therapy ¹³¹I-huA33 infusion, and histologic features were consistent with a nonspecific spongiform dermatitis and further immunostains for immune complexes/complement were negative.

No clinically significant gastrointestinal toxicity was observed, either in the initial study follow-up or in the long-term follow-up phases. Patient 13 experienced a "serum sickness"-like clinical syndrome in association with Biacore evidence of HAHA formation. Symptoms of myalgias, chills, and fatigue were prominent and commenced 1 week following the ¹³¹I-huA33 therapy dose. Biodistribution following the scout infusion was normal and excellent tumor-targeting was evident.

Maximum tolerated dose. The maximum tolerated dose was determined to be 40 mCi/m² of ¹³¹I. This was based on the occurrence of two episodes of dose-limiting toxicity (grade 4 neutropenia and grade 3 liver function test abnormalities) across six patients treated at the 50 mCi/m² dose level.

Evaluation of delayed toxicity. Eight patients were available for at least one follow-up assessment. The other seven patients were not available because of death due to tumor progression, other treatment, or withdrawal of consent for further follow-up. One patient developed biochemical hypo-

thyroidism on long-term follow-up in the context of previous treatment with trace ¹³¹I-labeled mAb as part of a previous clinical trial. The patient remained clinically euthyroid and oral thyroid hormone replacement was commenced with subsequent improvement in biochemical thyroid function. No significant deterioration in cardiac, respiratory, renal, or liver function attributable to ¹³¹I-huA33 was detected on long-term follow-up on available patients.

Pharmacokinetics. Pharmacokinetic analysis of scout infusion data showed consistent results across all dose levels, with T1/2 α (mean \pm SD) 23.0 \pm 12.5 hours and T1/2 β 135.2 \pm 46.9 hours. The calculated (mean \pm SD) C_{max} was 1.40 \pm 0.23 mg/L, CL was 0.035 \pm 0.010 L/hour, AUC was 144.4 \pm 41.8 hour \times mg/L, and V1 3.48 \pm 0.64 L.

Biodistribution and dosimetry of ¹³¹I-huA33: Biodistribution. All patients had normal ¹³¹I-huA33 biodistribution following the scout infusion, with initial images demonstrating ¹³¹I-huA33 blood pool distribution, followed by tumor uptake and bowel localization (Fig. 1). Colon activity gradually resolved with blood clearance of ¹³¹I-huA33. Excellent uptake of ¹³¹I-huA33 in tumor was seen in all patients, with persisting high uptake visualized up to 4 weeks after the ¹³¹I-huA33 therapy infusion (Fig. 2). Post-therapy images showed essentially identical biodistribution and tumor uptake of ¹³¹I-huA33 compared with scout infusions in all patients.

Whole body clearance. Whole body clearance [biologic half-life (T1/2)] of ¹³¹I-huA33 scout dose was calculated to be 227.52 \pm 46.15 hours (mean \pm SD).

Whole body and normal organ dosimetry. The results for individual organs and whole-body doses were similar between patients at the four dose levels. Target organs with the highest doses were lungs, liver, spleen, kidney, and thyroid, consistent with the blood pool appearance on gamma camera images and minor thyroid uptake from free ¹³¹I (Fig. 1). Mean \pm SD values were for kidney (0.23 \pm 0.07 cGy/MBq), liver (0.17 \pm 0.06 cGy/MBq), lungs (0.13 \pm

Table 1. Baseline patient demographics

Patient no.	Dose level (mCi/m ²)	Therapy ¹³¹ I dose administered		Age at study entry (years)	Sex	Karnofsky performance status	Site of primary tumor
		(mCi)	(GBq)				
1	20	32.7	1.21	65	M	90	rectum
2	20	34.8	1.29	65	M	100	colon
3	20	37.3	1.38	63	M	90	colon
4	30	55.4	2.05	50	M	90	rectum
5	30	68.0	2.52	63	M	80	colon
6	30	63.4	2.35	39	F	90	rectum
7	40	77.0	2.85	53	M	90	colon
8	40	72.2	2.67	68	M	100	rectum
9	40	69.0	2.55	73	M	100	rectum
10	50	75.5	2.80	67	F	90	colon
11	50	94.7	3.50	56	M	100	rectum
12	50	85.5	3.16	77	M	90	colon
13	50	81.6	3.02	62	F	90	colon
14	50	70.8	2.62	54	F	100	colon
15	50	84.1	3.11	74	M	100	colon

Table 2. Patient treatment history and response to ^{131}I -huA33

Patient no.	Adjuvant radiotherapy	Chemotherapy for metastatic disease	Other treatment for metastatic disease	Sites of tumor at study entry	Tumor response to ^{131}I -huA33
1	yes	5-FU, I, O, R	RFA of liver metastases, huA33	liver, lung, LN	PD
2	no	5-FU	—	liver	SD
3	no	IHA 5-FU, 5-FU	resection of liver metastases, alcohol injection, huA33	liver, lung, LN, sacrum	PD
4	no	O/5-FU, I, R	—	liver, LN, presacral	PD
5	no	5-FU, I	—	liver, lung, LN	PD
6	yes	5-FU, I	—	liver, lung, LN	SD
7	no	5-FU, I	resection of pelvic and left iliac fossa metastases, pelvic radiotherapy	liver, lung, pelvic, LN	SD
8	yes	5-FU, I	resection of liver metastases	liver, lung, LN	PD
9	yes	O/5-FU	Sibrotuzumab	liver	SD
10	no	5-FU, I	right lung lower lobectomy	liver, lung	PD
11	yes	IHA 5-FU, I/5-FU	—	liver, lung	PD
12	no	I	—	liver, abdominal wall	PD
13	no	O/5-FU	—	liver, lung	PD
14	no	O/5-FU	—	liver	PD
15	no	I,O	—	lung, LN, ascites	PD

Abbreviations: I, irinotecan; O, oxaliplatin; R, raltitrexed; IHA, intrahepatic artery; RFA, radiofrequency ablation; sibrotuzumab, clinical trial monoclonal antibody; LN, lymph nodes; PD, progressive disease; SD, stable disease.

0.04 cGy/MBq), spleen (0.26 ± 0.06 cGy/MBq), and thyroid (0.39 ± 0.16 cGy/MBq). The whole-body dose (effective dose equivalent) was calculated to be 0.80 ± 0.11 mSv/MBq (mean \pm SD).

Red marrow absorbed dose. The total red marrow radiation absorbed doses was similar between scout and therapy doses across all patients (0.056 ± 0.012 versus 0.054 ± 0.012 cGy/MBq). Total red marrow absorbed dose increased in proportion to therapy dose levels, with the mean \pm SD values at the 20 mCi/m² dose level (69.00 ± 5.05 cGy), 30 mCi/m² dose level (96.68 ± 12.19 cGy), 40 mCi/m² dose level

(122.46 ± 13.12 cGy), and 50 mCi/m² dose level (153.49 ± 46.80 cGy).

Tumor dosimetry. The mean specific absorbed tumor dose across all dose levels was 6.49 ± 2.47 Gy/GBq (mean \pm SD). Total tumor dose increased with dose level, and was 19.26 ± 11.68 Gy at the 40 mCi/m² dose level (range 12.04-32.73 Gy), and was 21.19 ± 8.17 Gy at the 50 mCi/m² dose level (range 11.73-31.81 Gy).

Immunogenicity. There was Biacore evidence of HAHA formation in four patients (nos. 4, 5, 7, and 13). Only one patient had clinical manifestations probably due to HAHA

Table 3. Study agent – related toxicity (maximum Common Toxicity Criteria grade)

Patient no.	Dose level (mCi/m ²)	Hemoglobin	Leukocytes	Neutrophils	Platelets
1	20	0	0	0	1
2	20	0	1	1	0
3	20	0	0	0	1
4	30	0	0	0	1
5	30	0	0	0	2
6	30	0	0	0	1
7	40	0	2	2	1
8	40	0	2	2	0
9	40	0	1	2	2
10	50	0	0	0	1
11	50	0	1	1	2
12	50	0	2	2	3
13	50	0	0	0	0
14	50	0	4	4	3
15	50	0	3	2	2

Table 4. Occurrence of cutaneous events, presence of HAHA, and ^{131}I -huA33 dose level

Patient no.	Dose level (mCi/m ²)	Rash (Common Toxicity Criteria grade)	Pruritus (Common Toxicity Criteria grade)	HAHA
1	20	0	1	no
2	20	0	0	no
3	20	2	2	no
4	30	0	0	yes
5	30	0	2	yes
6	30	0	0	no
7	40	3	2	yes
8	40	0	1	no
9	40	0	0	no
10	50	0	0	no
11	50	0	0	no
12	50	2	2	no
13	50	0	0	yes
14	50	3	3	no
15	50	0	0	no

NOTE: All listed cutaneous events were considered to be related to the study agent.

(see Toxicity). Onset of Biacore-detected HAHA was generally 2 to 3 weeks following the scout ^{131}I -huA33 dose, with peak values within a week of onset. The exception was patient 13, whose HAHA was Biacore-detectable 1 week after the scout dose, peaking 1 week later. Biacore responses gradually declined after peaking, although in almost all affected patients, HAHA was still detectable 7 weeks after the scout infusion.

Tumor response. On WHO computerized tomography-based criteria, 4 patients had stable disease at the end of study visit (6 weeks after ^{131}I -huA33 therapy dose) and 11 patients had progressive disease (Table 1). Patients with more bulky disease tended to develop progressive disease while in the study, compared with patients with relatively low volume tumors who were more likely to have stable disease 6 weeks after treatment. In most cases, patients had clinical symptoms or signs of tumor progression in addition to computerized tomography evidence.

Discussion

This phase I radioimmunotherapy study reinforces our current understanding of the A33-huA33 mAb system and improves our ability to apply this to patients with advanced colorectal cancer. Like muA33, huA33 has been shown to be well-tolerated when given by i.v. infusion. Tumor-targeting and biodistribution with ^{131}I -huA33 is comparable to radio-labeled muA33. HuA33 has a longer serum half-life compared with muA33 with a $T_{1/2\beta}$ of 135 versus 38.5 hours (10). The incidence of HAHA is clearly less than the frequency of human anti-mouse antibody generated with the murine mAb.

The temporal pattern and magnitude of hematologic toxicity observed was consistent with previous radioimmunotherapy trials, including studies with ^{131}I -muA33 (11). In our study, two episodes of dose-limiting toxicity were observed at 50 mCi/m² dose level: one episode of grade 4

neutropenia was clearly due to ^{131}I -huA33, and it was not possible to exclude an acute radiation-induced hepatitis as the cause of the grade 3 liver function test abnormalities in an additional patient. The maximum tolerated dose in the ^{131}I -muA33 study was 75 mCi/m², which is somewhat higher than the maximum tolerated dose of 40 mCi/m² observed for ^{131}I -huA33. This can be readily explained by the longer half-life of the humanized mAb labeled with a moderately long-lived radioisotope, and the resultant red marrow dose.

Non-infusion-related cutaneous toxicity has not been shown in any of the preceding muA33 or huA33 clinical studies. Thus, the 47% incidence of rash and/or pruritus in this huA33 radioimmunotherapy study was unexpected. The etiology of these symptoms remains unclear; however, they were almost certainly related to the huA33 protein rather than the radioisotope. The lack of correlation between incidence and severity of cutaneous reactions and ^{131}I dose level supports this hypothesis. One might expect the mechanism to relate to HAHA formation, however, there was no correlation between skin toxicity and HAHA based on Biacore data.

Clinically significant gastrointestinal toxicity was not observed in any of the muA33 or huA33 radioimmunotherapy trials. Despite the presence of A33 antigen on normal colonic epithelium, and showed targeting of this in patients based on autoradiography and gamma imaging, clinical sequelae have not been observed. There are two potential reasons for this: the most likely explanation is that the radiation dose given was too low to cause radiation colitis. The other plausible explanation (which may act in tandem with the first) is that the normal physiologic turnover of colonic epithelial cells was protective against effects of radiation-induced apoptosis. This is consistent with the calculation of the average colon elimination (k_{30}) half-life of ^{131}I -huA33 to be 32.4 ± 8.1 hours from a previous biopsy-based huA33 trial.⁴ The selection of ^{131}I rather than other β emitters (e.g., ^{90}Y or ^{177}Lu) labeled to huA33 was based

in part on the bowel uptake of huA33, and our concern regarding potential bowel toxicity such as has been reported with other ^{90}Y -labeled mAbs that also target an antigen expressed in normal bowel (25). The choice of ^{131}I has been vindicated by the absence of clinically significant bowel toxicity in this trial. The lack of bowel toxicity, combined with the excellent uptake in tumor of ^{131}I -huA33, indicates that radioimmunotherapy with huA33 is a highly attractive strategy for therapy of colorectal cancer.

Hypothyroidism is a known potential complication of ^{131}I -based radioimmunotherapy (26). The standard preventive procedure of oral iodine administration has been shown to reduce thyroid uptake of ^{131}I to <1% of the injected activity in patients receiving radioimmunotherapy with ^{131}I -labeled mAbs (27). Interestingly, the only patient to become biochemically hypothyroid only received 5.2 Gy to the thyroid gland—well below the mean experienced by the study population. It is possible that preexisting mild thyroid dysfunction at study entry progressed despite seemingly adequate oral iodine blockade.

In this study, four patients had stable disease at restaging in response to ^{131}I -huA33, all of whom had previously received two or fewer agents for advanced colon cancer. Eleven patients

had progressive disease at restaging, of whom eight patients had previously received two or fewer agents for advanced colon cancer. Although this might suggest an inverse relationship between the number of prior therapies and response to ^{131}I -huA33, the numbers are too small to draw any definitive conclusions.

Other ^{131}I labeled humanized mAbs have been tested in trials and reported in humans with colorectal cancer. hMN-14 is a humanized mAb targeting carcinoembryonic antigen (28) and a phase II radioimmunotherapy trial using ^{131}I -hMN-14 has been recently reported (29). A dose of 60 mCi/m² was studied in 30 patients with small volume metastatic colorectal cancer. Moderate and reversible myelosuppression was the dominant toxicity, although one patient developed grade 4 thrombocytopenia. The pattern of myelotoxicity reported in this ^{131}I -hMN-14 trial was very consistent with that observed in our ^{131}I -huA33 radioimmunotherapy study. Normal organ and tumor dosimetry for ^{131}I -hMN-14 has been shown to be similar to that for ^{131}I -huA33, considering the differences in mAb specificity and radioisotope dose (28, 29). The prolonged retention of ^{131}I -huA33 in tumor (at least 4 weeks) is unique, however, and is likely due to the cellular location of the A33 antigen in tumor cells and lack of trafficking of A33





Fig. 1. Anterior whole body gamma camera images following infusion of ^{131}I -huA33 in patient 4 (30 mCi/m² dose level) are shown: (A) day 0, (B) day 1, (C) day 2, and (D) day 5 post-scout infusion, and (E) day 6 post-therapy infusion. A standard for quantitation of ^{131}I -huA33 uptake is present adjacent to the left shoulder. A, Initial (day 0) images show blood pool appearance only, with a metastatic lesion in the liver demonstrating an initial hypovascular appearance. B, excellent targeting of the metastatic lesion in the liver by ^{131}I -huA33 is clearly seen (arrow), as early as day 1, and (C-D) increasing rapidly with time. D, some central necrosis in the tumor is also evident (arrow). Gradual bowel uptake (double arrow) of ^{131}I -huA33 is also seen. E, the post-therapy image shows identical biodistribution and tumor uptake of ^{131}I -huA33 compared to the scout dose. F, computerized tomography scan of the metastatic lesion in the liver in this patient.

Fig. 2. Prolonged tumor localization following therapy ^{131}I -huA33 infusion in patient 12 (50 mCi/m^2). Gamma camera images of the upper abdomen (A) anterior and (B) posterior (day 7 post-therapy infusion); (C) anterior (D) posterior (day 21 post-therapy infusion); (E) anterior and (F) posterior (day 28 post-therapy infusion) are shown. Excellent uptake of ^{131}I -huA33 in a metastatic lesion in the posterior right lobe of liver (arrow) is shown up to day 28 post-therapy infusion. Bowel activity (double arrow) reduces with time, in contrast to the prolonged tumor retention of ^{131}I -huA33.



antigen/antibody complex to intracellular lysosomes. Our estimates of tumor retention and dose are therefore likely to be underestimates of the true radiation dose to tumor with this therapeutic approach.

Radioimmunotherapy has yet to establish itself as a standard part of therapeutic strategy for epithelial malignancies. In this respect, its application is not nearly as advanced as radioimmunotherapy for hematologic malignancies. The role of ^{131}I -huA33 mAb is also not as yet clearly defined, despite clear evidence of antitumor efficacy in animal models, and the excellent tumor-targeting properties and stable disease in 4 of 15 patients shown in this trial. The dose to tumor at maximum tolerated dose in this trial ranged from 12 to 33 Gy, indicating that a single dose of ^{131}I -huA33 may not be sufficient to achieve tumoricidal effects alone. Improved dose to tumor could be achieved with higher LET β emitters (e.g., ^{90}Y or ^{177}Lu); however, the problem of potential bowel toxicity will require careful evaluation in future trials.

Combined modality therapy is a currently favored route of drug and therapeutic development. Preclinical data have shown

enhanced radiation sensitivity of tumor cells pretreated with cytotoxics such as paclitaxel (30). As a result, the combination of chemotherapy and radiotherapy has become standard treatment for a number of epithelial tumors over the last 10 years. The synergy of trastuzumab with cisplatin or paclitaxel observed in animal models of breast cancer has also translated into clinical practice, and Bevacizumab in combination with 5-fluorouracil (5-FU)/folinic acid has shown greater tumor response rates than 5-FU/folinic acid alone (31). There is also animal model data suggesting synergy between ^{131}I -huA33 and 5-FU for colorectal tumors (32), and a trial of ^{90}Y -anti-carcinoembryonic antigen chimeric T84.66 with 5-FU has shown the tolerability of this approach (33). Importantly, an orally bioavailable 5-FU prodrug, capecitabine, has been found to produce similar tumor response rates to i.v.-administered 5-FU, and also has been shown to improve response rates when given in conjunction with irradiation (34). The rationale for the combination of ^{131}I -huA33 radioimmunotherapy and capecitabine is therefore compelling, and we intend to evaluate this therapeutic approach in future trials.

References

- DeNardo GL, O'Donnell RT, Kroger LA, et al. Strategies for developing effective radioimmunotherapy for solid tumors. *Clin Cancer Res* 1999;5:3219s–23s.
- Scott AM, Welt S. Antibody-based immunological therapies. *Curr Opin Immunol* 1997;9:717–22.
- Greenlee RT, Hill-Harmon MB, Murray T, Thun M. Cancer statistics, 2001. *CA Cancer J Clin* 2001; 51:15–36.
- Grothey A SH. New chemotherapy approaches in colorectal cancer. *Curr Opin Oncol* 2001;13:275–86.
- Sakamoto J, Kojima H, Kato J, Hamashima H, Suzuki H. Organ-specific expression of the intestinal epithelium-related antigen A33, a cell surface target for antibody-based imaging and treatment in gastrointestinal cancer. *Cancer Chemother Pharmacol* 2000;46 Suppl:S27–32.
- Heath JK, White SJ, Johnstone CN, et al. The human A33 antigen is a transmembrane glycoprotein and a novel member of the immunoglobulin superfamily. *Proc Natl Acad Sci U S A* 1997;94:469–74.
- Johnstone CN, Tebbutt NC, Abud HE, et al. Characterization of mouse A33 antigen, a definitive marker for basolateral surfaces of intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* 2000;279: G500–10.
- Moritz RL, Ritter G, Catimel B, et al. Micro-sequencing strategies for the human A33 antigen, a novel surface glycoprotein of human gastrointestinal epithelium. *J Chromatogr A* 1998;798:91–101.
- Ritter G, Cohen LS, Nice EC, et al. Characterization of posttranslational modifications of human A33 antigen, a novel palmitoylated surface glycoprotein of human gastrointestinal epithelium. *Biochem Biophys Res Commun* 1997;236:682–6.
- Welt S, Divgi CR, Real FX, et al. Quantitative analysis of antibody localization in human metastatic colon cancer: a phase I study of monoclonal antibody A33. *J Clin Oncol* 1990;8:1894–906.
- Welt S, Divgi CR, Kemeny N, et al. Phase I/II study of iodine 131-labeled monoclonal antibody A33 in patients with advanced colon cancer. *J Clin Oncol* 1994; 12:1561–71.
- Welt S, Scott AM, Divgi CR, et al. Phase I/II study of iodine 125-labeled monoclonal antibody A33 in patients with advanced colon cancer. *J Clin Oncol* 1996; 14:1787–97.
- King DJ, Antoniw P, Owens RJ, et al. Preparation and preclinical evaluation of humanised A33 immunocjugates for radioimmunotherapy. *Br J Cancer* 1995; 72:1364–72.
- Welt S, Ritter G, Williams C Jr, et al. Phase I study of anti-colon cancer humanised antibody A33. *Clin Cancer Res* 2003;9:1338–46.
- Welt S, Ritter G, Williams C Jr, et al. Preliminary report of a phase I study of combination chemotherapy and huA33 immunotherapy. *Clin Cancer Res* 2003; 9:1347–53.
- Lee FT, Hall C, Rigopoulos A, et al. Immuno-PET of human colon xenograft-bearing BALB/c nude mice using 124I-CDR-grafted humanized A33 monoclonal antibody. *J Nucl Med* 2001;42:764–9.
- Liu A, Williams LE, Raubitschek AA. A CT assisted method for absolute quantitation of internal radioactivity. *Med Phys* 1996;23:1919–28.
- Loevinger R, Budinger TF, Watson EE. MIRD primer for absorbed dose calculations. New York: The Society of Nuclear Medicine; 1991.
- Stabin MG. MIRDose: personal computer software for internal dose assessment in nuclear medicine. *J Nucl Med* 1996;37:538–46.
- Shen S, DeNardo GL, Sgouros G, O'Donnell RT, DeNardo SJ. Practical determination of patient-specific marrow dose using radioactivity concentration in blood and body. *J Nucl Med* 1999;40: 2102–6.

21. Ritter G, Cohen LS, Williams C Jr, Richards EC, Old LJ, Welt S. Serological analysis of human anti-human antibody responses in colon cancer patients treated with repeated doses of humanized monoclonal antibody A33. *Cancer Res* 2001;61:6851–9.
22. Scott AM, Lee FT, Hopkins W, et al. Specific targeting, biodistribution, and lack of immunogenicity of chimeric anti-GD3 monoclonal antibody KM871 in patients with metastatic melanoma: results of a phase I trial. *J Clin Oncol* 2001;19:3976–87.
23. Scott AM, Wiseman G, Welt S, et al. A phase I dose-escalation study of sibrutumab in patients with advanced or metastatic fibroblast activation protein-positive cancer. *Clin Cancer Res* 2003;9:1639–47.
24. Scott AM, Geleick D, Rubira M, et al. Construction, production, and characterization of humanized anti-Lewis Y monoclonal antibody 3S193 for targeted immunotherapy of solid tumors. *Cancer Res* 2000;60:3254–61.
25. Knox SJ, Goris ML, Tempero M, et al. Phase II trial of yttrium-90-DOTA-biotin pretargeted by NR-LU-10 antibody/streptavidin in patients with metastatic colon cancer. *Clin Cancer Res* 2000;6:406–14.
26. Press OW, Eary JF, Badger CC, et al. Treatment of refractory non-Hodgkin's lymphoma with radiolabeled MB-1 (anti-CD37) antibody. *J Clin Oncol* 1989;7:1027–38.
27. Behr TM, Juweid ME, Sharkey RM, et al. Thyroid radiation doses during radioimmunotherapy of CEA-expressing tumours with ¹³¹I-labelled monoclonal antibodies. *Nucl Med Commun* 1996;17:767–80.
28. Hajjar G, Sharkey RM, Burton J, et al. Phase I radioimmunotherapy trial with iodine-131 – labeled humanized MN-14 anti-carcinoembryonic antigen monoclonal antibody in patients with metastatic gastrointestinal and colorectal cancer. *Clin Colorectal Cancer* 2002;2:31–42.
29. Behr TM, Liersch T, Greiner-Bechert L, et al. Radioimmunotherapy of small-volume disease of metastatic colorectal cancer. *Cancer* 2002;94:1373–81.
30. Liebmann J, Cook JA, Fisher J, Teague D, Mitchell JB. Changes in radiation survival curve parameters in human tumor and rodent cells exposed to paclitaxel (Taxol). *Int J Radiat Oncol Biol Phys* 1994;29:559–64.
31. Kabbinnavar F, Hurwitz HI, Fehrenbacher L, et al. Phase II, randomized trial comparing bevacizumab plus fluorouracil (FU)/leucovorin (LV) with FU/LV alone in patients with metastatic colorectal cancer. *J Clin Oncol* 2003;21:60–5.
32. Tschmelitsch J, Barendsward E, Williams C Jr, et al. Enhanced antitumor activity of combination radioimmunotherapy (¹³¹I-labeled monoclonal antibody A33) with chemotherapy (fluorouracil). *Cancer Res* 1997;57:2181–6.
33. Wong JY, Shibata S, Williams LE, et al. A phase I trial of 90Y-anti-carcinoembryonic antigen chimeric T84.66 radioimmunotherapy with 5-fluorouracil in patients with metastatic colorectal cancer. *Clin Cancer Res* 2003;9:5842–52.
34. Van Cutsem E, Twelves C, Cassidy J, et al. Oral capecitabine compared with intravenous fluorouracil plus leucovorin in patients with metastatic colorectal cancer: results of a large phase III study. *J Clin Oncol* 2001;19:4097–106.

Clinical Cancer Research

Phase I Trial of ¹³¹I-huA33 in Patients with Advanced Colorectal Carcinoma

Geoffrey Chong, Fook Thean Lee, Wendie Hopkins, et al.

Clin Cancer Res 2005;11:4818-4826.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/11/13/4818>

Cited articles This article cites 31 articles, 19 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/11/13/4818.full#ref-list-1>

Citing articles This article has been cited by 11 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/11/13/4818.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/11/13/4818>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.