

Response of Some Head and Neck Cancers to Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors May Be Linked to Mutation of *ERBB2* rather than *EGFR*

Ezra E. W. Cohen,¹ Mark W. Lingen,^{1,2,3} Leslie E. Martin,² Patricia L. Harris,⁴ Brian W. Brannigan,⁴ Sara M. Haserlat,⁴ Ross A. Okimoto,⁴ Dennis C. Sgroi,^{4,5} Sonika Dahiya,^{4,5} Beth Muir,^{4,5} John R. Clark,⁴ James W. Rocco,^{4,6} Everett E. Vokes,^{1,3} Daniel A. Haber,⁴ and Daphne W. Bell⁴

Abstract Purpose: Small-molecule tyrosine kinase inhibitors (TKI) of the epidermal growth factor receptor (EGFR) have shown modest yet reproducible response rates in patients with squamous cell carcinoma of the head and neck (SCCHN). Somatic mutations in *EGFR* have recently been shown to be predictive of a clinical response in patients with non-small cell lung cancer (NSCLC) treated with these inhibitors. The objective of this study was to determine if such mutations, or recently reported mutations in *ERBB2*, also underlie EGFR-TKI responsiveness in SCCHN patients. **Experimental Design:** We sequenced the kinase domain of *EGFR* and exon 20 of *ERBB2* in tumor specimens from eight responsive patients. In addition, mutational analysis was done on tumor specimens from nine gefitinib nonresponders and 65 unselected cases of SCCHN. **Results:** None of eight TKI-responsive specimens had mutations within the kinase domain of *EGFR*. *EGFR* amplification was also not associated with drug responsiveness. However, a single responsive case had a somatic missense mutation within exon 20 of *ERBB2*. **Conclusion:** Our data indicate that unlike NSCLC, *EGFR* kinase mutations are rare in unselected cases of SCCHN within the United States and are not linked to gefitinib or erlotinib responses in SCCHN. Alternative mechanisms, including *ERBB2* mutations, may underlie responsiveness in this tumor type.

Squamous cell carcinoma of the head and neck (SCCHN) is diagnosed annually in ~40,000 individuals in the United States (1). Despite current therapeutic modalities, as many as half these will recur, or develop metastases, and eventually succumb to their disease, highlighting the need for alternative forms of therapeutic intervention. Because epidermal growth factor receptor (EGFR: ERBB1 and HER1) is almost universally expressed in SCCHN and high levels of expression have been correlated with a poor clinical prognosis (2–6), clinical trials were undertaken to assess the efficacy of kinase inhibitors that target this growth factor, including gefitinib and erlotinib (7),

reporting significant responses in up to 11% of patients refractory to other therapies (8).

Although the response rate of SCCHN to gefitinib and erlotinib is comparable with that seen in non-small cell lung cancer (NSCLC), the demographic and histologic variables associated with responsive cases differ remarkably. In lung cancer, drug responses are characteristically seen in adenocarcinomas arising in nonsmokers, with an increased frequency among women and Asians (9, 10). In contrast, responsive cases of SCCHN were indistinguishable from nonresponsive cases in terms of their histology or associated history of tobacco exposure (8).

In NSCLC, clinical responses to gefitinib or erlotinib have recently been ascribed to the presence of activating mutations within the kinase domain of *EGFR* (11–16). Experimentally, cells expressing these mutant receptors seem dependent upon altered survival signals transduced by these receptors, which themselves also display increased sensitivity to inhibition by gefitinib (17). Thus, at least in a subset of NSCLC, somatic mutations in *EGFR* seem to drive tumorigenesis and provide a biologically relevant target for therapeutic intervention. Clinically, these mutations are associated with dramatic responses to EGFR tyrosine kinase inhibitors (TKI) and may define a biological subset of NSCLC patients. To determine if this molecular mechanism extends to SCCHN, we sequenced the kinase domain of *EGFR* (exons 18–24) in tumors from gefitinib or erlotinib responsive cases as well as untreated cases from the United States.

Authors' Affiliations: Departments of ¹Medicine, ²Pathology, and ³Radiation and Cellular Oncology, University of Chicago, Chicago, Illinois; ⁴Cancer Center and ⁵Department of Pathology, Massachusetts General Hospital, Charlestown, Massachusetts; and ⁶Massachusetts Eye and Ear Infirmary, Boston, Massachusetts. Received 4/29/05; revised 6/20/05; accepted 8/23/05.

Grant support: National Cancer Institute/NIH federal funds under contract nos. N01-CM-17102 and P30-CA14599-30 (clinical trials from which tissue samples were derived), University of Chicago Cancer Research Center, AstraZeneca Pharmaceuticals, grant P01 95281 (D.W. Bell and D.A. Haber), and Doris Duke Charitable Foundation (D.A. Haber).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Daniel A. Haber, Massachusetts General Hospital Cancer Center, CNY7, 149 13th Street, Charlestown, MA 02129. Phone: 617-724-7805; Fax: 617-724-6919; E-mail: haber@helix.mgh.harvard.edu.

© 2005 American Association for Cancer Research.
doi:10.1158/1078-0432.CCR-05-0926

Materials and Methods

Tissue samples. Tissue samples were collected from patients treated at the University of Chicago or Massachusetts General Hospital. Three groups of patient samples were analyzed: those with tumors responding to EGFR-TKI, those with tumors not responding to EGFR-TKI, and randomly selected patients not previously treated with EGFR-TKI. Samples from patients who were treated with EGFR-TKI were collected before therapy with either gefitinib (500 or 250 mg) or erlotinib (150 mg in combination with 15 mg/kg bevacizumab; Table 1). The Response Evaluation Criteria in Solid Tumors were used to evaluate response in all cases. Consent to obtain, store, and analyze tissue was obtained on all patients according to institutional review board guidelines.

Mutational analysis of EGFR and ERBB2. H&E-stained sections of formalin-fixed, paraffin-embedded tissue or of snap-frozen tissue were reviewed by a pathologist to estimate the overall tumor cell content. In cases where tumor cells comprised <70% of the tissue, laser capture microdissection was employed to isolate a homogenous tumor cell population. DNA was isolated by standard phenol-chloroform extraction and ethanol precipitated using glycogen as a carrier followed by resuspension in 30 μ L TE buffer.

Exons 18 to 24 of *EGFR* and exon 20 of *ERBB2* were amplified using nested PCR consisting of an initial primary PCR followed by a secondary PCR. Primer pairs (sense and antisense) used in the initial amplification of *EGFR* were as follows: exon 18, CAAAGTCCCGTGTCC-TGGCACCCAAGC and CCAAACACTCAGTCAAACAAAGAG; exon 19, GCAATATCAGCCTTAGGTGCGGCTC and CATAGAAAGTGAACATT-TAGGATGTG; exon 20, CCATGAGTACGTATTTTGAAGCTC and CATA-TCCCATGGCAAACCTCTTGC; exon 21, CTAACGTTCCGCCAGC-CATAAGTCC and GCTGCGAGCTACCCAGAATGTCTGG; exon 22, GAGCAGCCCTGAACTCCGTCAGACTG and CTCAGTACAATAGATA-GACAGCAATG; exon 23, CAGGACTACAGAAATGTAGGTTTC and GTGCCCTGCCTTAAGTAATGTGATGAC; exon 24, GACTGGAAGTGTG-CATCACCAATG and GGTTTAATAATGCGATCTGGGACAC. The primer pair (CTGTGGGCCATGGCT GTGGTTTG and GGAATGGGAAGCACCATGTAGACC) was used in an initial amplification of exon 20 of

ERBB2. Primary reactions contained 2 μ L genomic DNA as a template in a 25- μ L reaction consisting of 1 \times buffer, 50 μ mol/L deoxynucleotide triphosphate, 200 nmol/L sense primer, 200 nmol/L antisense primer, and 0.8 unit Expand Taq (Roche Diagnostics, Mannheim, Germany).

Nested primers pairs used in secondary amplification of *EGFR* were exon 18, GCACCCAAGCCCATGCCGTGGCTGC and GAAACAAAGAG-TAAAGTAGATGATGG; exon 19 CCTTAGGTGCGGCTCCACAGC and CATTAGGATGTGGAGATGAGC; exon 20, GAAACTCAAGATCGCATT-CATGC and GCAAACCTCTTGTATCCACAGGAG; exon 21, CAGCCA-TAAGTCCTCGACGTGG and CATCCTCCCC TGCATGTGTAAAC; exon 22, GACGGGTCTGGGGTGATCTGGCTC and CTCAGTACAATAGATA-GACAGCAATG; exon 23, GTAGGTTTCTAAACATCAAGAAAC and GTGATGACATTTCTCCAGGGATGC; exon 24, CATCACCAATGCC-TTCTTTAAGC and GCTGGAGGGTTTAATAATGCGATC. Exon 20 of *ERBB2* was amplified using nested primers CTGTGGGCCATGGC-TGTGGTTTG and GGAAGCACCCATGTAGACCTTCTGG. Nested reactions contained 2 μ L primary PCR product as a template in a 25- μ L reaction consisting of 1 \times buffer, 50 μ mol/L deoxynucleotide triphosphate, 200 nmol/L sense primer, 200 nmol/L antisense primer, and 0.8 unit Expand Taq (Roche Diagnostics). Cycling conditions for both primary and secondary PCR reactions were 95°C for 15 minutes followed by 40 cycles consisting of 95°C for 30 seconds, 58°C for 30 seconds, 72°C for 45seconds followed by 72°C for 5 minutes. PCR amplicons were purified using exonuclease I (U.S. Biochemical, Cleveland, OH) and shrimp alkaline phosphatase (U.S. Biochemical) and diluted in water before bidirectional capillary sequencing, which was done using BigDye Terminator v1.1 chemistry (Applied Biosystems, Foster City, CA) in combination with an ABI3100 instrument according to the manufacturer's instructions. Electropherograms were aligned and reviewed using Sequence Navigator software.

Determination of EGFR copy number. *EGFR* copy number was determined by Taqman real-time quantitative PCR with Taqman Universal PCR mastermix and an ABI Prism 7900HT sequence detection system (Applied Biosystems). The primers (5'-3') and fluorogenic probe used for *EGFR* were CAATTGCCAGTTAACGTCTTCCCTT (sense primer), TTTCTCACCTTCTGGGATCCA (antisense primer), and TCTCTCTGTCATAGGGAC (probe). For the control gene, *PCDH7*,

Table 1. Characteristics of patients on EGFR inhibitor clinical trials who demonstrated an objective response

Case no	Age at start of therapy	Sex	Race	Primary site	Histology*	Smoking history [†]	Prior therapy	Treatment [‡]	Response	Duration of response (mo)	Overall survival (mo)
1	50	F	W	OC (tongue)	VC	4	CRT, CT	G 500 mg	PR	11.0	18
2	51	M	W	OP (pharynx)	PD	3	CT	G 500 mg	PR	3.1	3.2
3	53	M	W	OC (buccal)	PD	2	CRT	G 500 mg	CR	4.7	6
4	64	M	W	OP (BoT)	PD	1	CRT	G 500 mg	PR	8.5	20
5	72	M	W	SGL	MD	5	CRT	G 500 mg	Minor	9.0	37.4
6	62	M	W	OP (pharynx)	WD	4	CRT	G 250 mg	PR	5.5	5.5
7	75	F	W	OC (buccal)	MD	4	CRT	E 150 mg + B 15 mg/kg	CR	>8.0 [§]	>8.0
8	59	M	W	OP (tonsil)	MD-PD	3	CRT, CT	G 500 mg	PR	4	7

NOTE: Response was assessed using the Response Evaluation Criteria in Solid Tumors in all patients, and all responses were verified by independent review. Duration of response is defined from first evidence of response until first evidence of disease progression measured in months. Overall survival is defined from start of therapy until death measured in months.

Abbreviations: M, male; F, female; W, White; OC, oral cavity; OP, oropharynx; AR, alveolar ridge; BoT, base of tongue; SGL, supraglottic larynx; VC, verrucous carcinoma; WD, well differentiated; MD, moderately differentiated; PD, poorly differentiated; CRT, chemoradiotherapy; CT, chemotherapy alone; G, gefitinib; E, erlotinib; B, bevacizumab; PR, partial response; CR, complete response.

*Only patients with squamous cell carcinoma were eligible to participate in the clinical trials.

[†]1, never used tobacco products; 2, used pipe, cigar, or chewing tobacco; 3, <20 pack-years cigarette use; 4, 20-40 pack-years cigarette use; 5, >40 pack-years cigarette use. Pack years is defined as the number of packs smoked per day multiplied by the number of years smoked.

[‡]All EGFR-TKIs were administered orally once daily without interruption. Bevacizumab was administered intravenously every 3 weeks.

[§]This patient remains on therapy at time of article submission.

Table 2. Number of cases analyzed from each clinical cohort

Clinical cohort	No. cases
EGFR-TKI Responders	8
EGFR-TKI Nonresponders	9
Unselected SCCHN	
Smokers	50
Never smokers	15

these were GCTGCAATCTCCTCCCTGAA (sense primer), TGCCTT-TTCTCACCTGCATTC (antisense primer), and CCACTGCTCCGACATG (probe).

Results

The kinase domain of *EGFR* (exons 18-24) was sequenced in tumors from eight gefitinib or erlotinib responsive cases of SCCHN (Table 1) treated at the University of Chicago (seven cases) or at Massachusetts General Hospital (one case). Of these, two cases had a complete response, and others had partial responses. We also sequenced the *EGFR* kinase domain in nine tumors from gefitinib nonresponders. No somatic mutations were detected in any of these TKI-treated cases of SCCHN (Table 2). To determine whether such mutations are present in unselected cases of SCCHN, we genotyped 50 cases of characteristic SCCHN cancers from the Massachusetts Eye and Ear Infirmary tumor bank and an additional 15 cases of SCCHN arising in never smokers (Table 2). None of these cases had detectable mutations within the kinase domain of *EGFR*. Given the recent report of *EGFR* gene amplification in some NSCLC cases responsive to gefitinib (18), as well as the occurrence of *EGFR* amplification in a subset of untreated cases (19, 20), we also determined the copy number of *EGFR* in treated SCCHN cases using quantitative PCR. No amplification of *EGFR* was detected among five responsive cases, whereas a single nonresponsive case had low level (2- to 4-fold) amplification (data not shown).

In addition to specific mutations in the kinase domain of *EGFR*, similar mutations have recently been reported in the related growth factor *ERBB2* (*HER2*) within a subset of NSCLC (21, 22). *ERBB2* heterodimerizes with *EGFR*, such that inhibition of *EGFR* kinase activity by TKI is expected to reduce *ERBB2*-dependent signaling. Furthermore, although both gefitinib and erlotinib have high specificity for *EGFR*, they also exhibit some *in vitro* activity towards *ERBB2* (gefitinib IC_{50} = 3,700 nmol/L; erlotinib IC_{50} = 350 nmol/L). We therefore tested for mutations in the reported *ERBB2* hotspot (exon 20) in our cohort of TKI-responsive cases of SCCHN. Of four evaluable cases of responsive SCCHN, one (case 2, treated with gefitinib 500 mg oral dose) harbored a heterozygous mutation within *ERBB2*, which was not present in surrounding normal stromal tissue (Fig. 1). The variant resulted in a nucleotide alteration from T to C at position 2318 leading to an amino acid substitution of alanine for valine at codon 773. Importantly, this residue forms part of the recurrent insertions (ins774AYVM and ins776YVMA) previously described in untreated NSCLC (21, 22), raising the possibility that such mutations may be linked to EGFR-TKI responsiveness. No

somatic mutations of *ERBB2* were identified in tumors from nine unresponsive SCCHN cases ($P = 0.307$) or among 56 untreated cases resected from smokers (50 cases) or never smokers (six cases).

Discussion

We have found that neither *EGFR* kinase domain mutations nor gene amplification explain the response of a subset of SCCHN to erlotinib and gefitinib, and that *EGFR* mutations are rare among cases of SCCHN within the U.S. population. While this work was in progress, Lee et al. reported the presence of *EGFR* kinase domain mutations in ~7% of unselected cases of SCCHN from Korea (23). The fact that we did not observe such mutations among 65 cases of unselected SCCHN from the United States, 17 cases of patients treated with EGFR-TKI, nor among seven SCCHN cell lines reported previously (11), suggests that the frequency of *EGFR* mutations in SCCHN cancer may differ among ethnic groups. Such a difference in the prevalence of *EGFR* mutations between East Asian and European populations is now well established for NSCLC (12, 14–16, 24–26), and its extension to SCCHN is of




Fig. 1. Mutation of *ERBB2* within a gefitinib-responsive tumor. A single base substitution of C for T at nucleotide 2318 (arrow) leads to an amino acid substitution of alanine for valine at codon 773 in the tumor from case 2. A wild-type sequence is shown for comparison.

considerable interest in searching for genetic and environmental risk factors. It will be important to determine whether *EGFR* mutations in Eastern Asian SCCHN patients are associated with response to EGFR-TKIs.

Our findings further suggest that *ERBB2* mutations may be linked to EGFR-TKI sensitivity in at least some of these responsive cases. Although *ERBB2* mutations within this region of the gene have been described in NSCLC their functional relevance to gefitinib or erlotinib responsiveness, or dual inhibitors of EGFR/*ERBB2*, such as lapatinib or HKI-272, has not yet been evaluated. This observation has clinical significance, as *EGFR* mutational testing enters the clinical arena as a predictor of therapeutic response to TKI in lung cancer. Alternative mechanisms must be sought to explain the majority of responses in SCCHN.

The absence of *EGFR* mutations in TKI-responsive SCCHN studied here is perhaps not surprising, given the unique subset of lung cancers defined by such mutations (i.e., adenocarcinomas arising in never smokers), with apparent modulation by both sex and ethnic factors (11–16, 24–26). In our study, no such distinct phenotype was evident in responsive cases of

SCCHN. In this context, it is interesting to note that a dose-response effect for gefitinib may exist in SCCHN cancers (response rates of 1% in a study using 250-mg daily dosing, versus 11% in a second study using a 500-mg dose; refs. 8, 27). Thus, in contrast to the exquisite sensitivity of tumors with mutant *EGFR* to gefitinib and the absence of different responses at these two doses (9, 10), inhibition of EGFR-dependent signaling even in responsive cases of SCCHN may require higher drug levels.

In conclusion, we note that most SCCHN cases responsive to gefitinib or erlotinib do not seem to have mutations in *EGFR* or *ERBB2*, as does a small subset of responsive NSCLC. Further studies are required to determine whether a unifying molecular mechanism may underlie TKI responsiveness in such cases and potentially in other tumor types where occasional gefitinib and erlotinib responses have been reported.

Acknowledgments

We thank Janet Dancey, M.D. for her assistance in conducting the clinical trials and Cyndy Bajda and Rosalyn Williams for their assistance with data management.

References

- Jemal A, Tiwari RC, Murray T, et al. Cancer statistics, 2004. *CA Cancer J Clin* 2004;54:8–29.
- Grandis JR, Tweardy DJ. TGF- α and EGFR in head and neck cancer. *J Cell Biochem* 1993;17F:188–91.
- Ang KK, Berkey BA, Tu X, et al. Impact of epidermal growth factor receptor expression on survival and pattern of relapse in patients with advanced head and neck carcinoma. *Cancer Res* 2002;62:7350–6.
- Chen BK, Ohtsuki Y, Furihata M, et al. Co-overexpression of p53 protein and epidermal growth factor receptor in human papillary thyroid carcinomas correlated with lymph node metastasis, tumor size and clinicopathologic stage. *Int J Oncol* 1999;15:893–8.
- Rubin Grandis J, Melhem MF, Gooding WE, et al. Levels of TGF- α and EGFR protein in head and neck squamous cell carcinoma and patient survival. *J Natl Cancer Inst* 1998;90:824–32.
- Etienne MC, Pivrot X, Formento JL, et al. A multifactorial approach including tumoural epidermal growth factor receptor, p53, thymidylate synthase and dihydropyrimidine dehydrogenase to predict treatment outcome in head and neck cancer patients receiving 5-fluorouracil. *Br J Cancer* 1999;79:1864–9.
- Cohen EE, Lingen MW, Vokes EE. The expanding role of systemic therapy in head and neck cancer. *J Clin Oncol* 2004;22:1743–52.
- Cohen EE, Rosen F, Stadler WM, et al. Phase II trial of ZD1839 in recurrent or metastatic squamous cell carcinoma of the head and neck. *J Clin Oncol* 2003;21:1980–7.
- Kris MG, Natale RB, Herbst RS, et al. Efficacy of gefitinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in symptomatic patients with non-small cell lung cancer: a randomized trial. *JAMA* 2003;290:2149–58.
- Fukuoka M, Yano S, Giaccone G, et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2003;21:2237–46.
- Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004;350:2129–39.
- Paez JG, Janne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004;304:1497–500.
- Pao W, Miller V, Zakowski M, et al. EGF receptor gene mutations are common in lung cancers from “never smokers” and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci U S A* 2004;101:13306–11.
- Huang SF, Liu HP, Li LH, et al. High frequency of epidermal growth factor receptor mutations with complex patterns in non-small cell lung cancers related to gefitinib responsiveness in Taiwan. *Clin Cancer Res* 2004;10:8195–203.
- Tokumo M, Toyooka S, Kiura K, et al. The relationship between epidermal growth factor receptor mutations and clinicopathologic features in non-small cell lung cancers. *Clin Cancer Res* 2005;11:1167–73.
- Mitsudomi T, Kosaka T, Endoh H, et al. Mutations of the epidermal growth factor receptor gene predict prolonged survival after gefitinib treatment in patients with non-small-cell lung cancer with postoperative recurrence. *J Clin Oncol* 2005;11:1–8.
- Sordella R, Bell DW, Haber DA, Settleman J. Gefitinib-sensitizing EGFR mutations in lung cancer activate anti-apoptotic pathways. *Science* 2004;305:1163–7.
- Amann J, Kalyankrishna S, Massion PP, et al. Aberrant epidermal growth factor receptor signaling and enhanced sensitivity to EGFR inhibitors in lung cancer. *Cancer Res* 2005;65:226–35.
- Weichselbaum RR, Dunphy EJ, Beckett MA, et al. Epidermal growth factor receptor gene amplification and expression in head and neck cancer cell lines. *Head Neck* 1989;11:437–42.
- Rikimaru K, Tadokoro K, Yamamoto T, Enomoto S, Tsuchida N. Gene amplification and overexpression of epidermal growth factor receptor in squamous cell carcinoma of the head and neck. *Head Neck* 1992;14:8–13.
- Stephens P, Hunter C, Bignell G, et al. Lung cancer: intragenic *ERBB2* kinase mutations in tumours. *Nature* 2004;431:525–6.
- Shigematsu H, Takahashi T, Nomura M, et al. Somatic mutations of the HER2 kinase domain in lung adenocarcinomas. *Cancer Res* 2005;65:1642–6.
- Lee JW, Soung YH, Kim SY, et al. Somatic mutations of EGFR gene in squamous cell carcinoma of the head and neck. *Clin Cancer Res* 2005;11:2879–82.
- Kosaka T, Yatabe Y, Endoh H, Kuwano H, Takahashi T, Mitsudomi T. Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. *Cancer Res* 2004;64:8919–23.
- Marchetti A, Martella C, Felicioni L, et al. EGFR mutations in non-small-cell lung cancer: analysis of a large series of cases and development of a rapid and sensitive method for diagnostic screening with potential implications on pharmacologic treatment. *J Clin Oncol* 2005;23:857–65.
- Shigematsu H, Lin L, Takahashi T, et al. Clinical and biological features associated with epidermal growth factor receptor gene mutations in lung cancers. *J Natl Cancer Inst* 2005;97:339–46.
- Kane MA, CE M, List B, et al. Phase II study of 250 mg gefitinib in advanced squamous cell carcinoma of the head and neck (SCCHN). In ASCO, Vol. 22. New Orleans: ASCO, 2004.

Clinical Cancer Research

Response of Some Head and Neck Cancers to Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors May Be Linked to Mutation of *ERBB2* rather than *EGFR*

Ezra E.W. Cohen, Mark W. Lingen, Leslie E. Martin, et al.

Clin Cancer Res 2005;11:8105-8108.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/11/22/8105>

Cited articles This article cites 26 articles, 14 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/11/22/8105.full#ref-list-1>

Citing articles This article has been cited by 29 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/11/22/8105.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/11/22/8105>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.