

Aberrantly Methylated *CDKN2A*, *MGMT*, and *MLH1* in Colon Polyps and in Fecal DNA from Patients with Colorectal Polyps

Zsolt Petko,¹⁰ Mahan Ghiassi,¹ Anthony Shuber,⁷ Janice Gorham,⁷ Walter Smalley,^{1,3,4,6} M. Kay Washington,² Stephen Schultenover,^{2,5} Shiva Gautam,⁹ Sanford D. Markowitz,⁸ and William M. Grady^{10,11,12}

¹Division of Gastroenterology, Department of Medicine, ²Departments of Pathology, ³Preventive Medicine, and ⁴Surgery, Vanderbilt University Medical School; ⁵Pathology Service and ⁶Medical Service, Department of Veterans Affairs Tennessee Valley Health Care System, Nashville, Tennessee; ⁷EXACT Sciences, Marlborough, Maryland; ⁸Ireland Cancer Center, University Hospitals of Cleveland, Case Western Reserve University and Howard Hughes Medical Institute, Cleveland, Ohio; ⁹Beth Israel Deaconess Medical Center, Boston, Maryland; ¹⁰Division of Clinical Research, Fred Hutchinson Cancer Research Center; ¹¹Division of Gastroenterology, Department of Medicine, University of Washington Medical School; and ¹²Department of Veterans Affairs Puget Sound Health Care System, Seattle, Washington

ABSTRACT

Colon cancer is the third leading cause of cancer-related death in the United States, affecting ~147,000 people each year. Most colon cancers arise from benign neoplasms and evolve into adenocarcinomas through a stepwise histologic progression sequence that starts from adenomas or hyperplastic polyps/serrated adenomas. Genetic alterations and, more recently, epigenetic alterations have been associated

with specific steps in this polyp-adenocarcinoma sequence and likely drive the histologic progression of colon cancer. Consequently, we have assessed in colon adenomas and hyperplastic polyps the methylation status of *MGMT*, *CDKN2A*, and *MLH1* to determine the timing and frequency of these events in the polyp-carcinoma progression sequence and subsequently to analyze the potential for these methylated genes to be molecular markers for adenomas and hyperplastic polyps. We have found that methylated *MGMT*, *CDKN2A*, and *MLH1* occur in 49%, 34%, and 7% of adenomas and in 5%, 10%, and 7% of hyperplastic polyps, respectively, and that they are more common in histologically advanced adenomas. Furthermore, analysis of fecal DNA from persons who have undergone colonoscopic exams revealed methylated *CDKN2A*, *MGMT*, and *MLH1* in fecal DNA from 31%, 48%, and 0% of individuals with adenomas and from 16%, 27%, and 10% of individuals with no detectable polyps, respectively. These results show that aberrant methylated genes can be detected frequently in sporadic colon polyps and that they can be detected in fecal DNA. Notably, improvements in the specificity and sensitivity of the fecal DNA-based assays will be needed to make them clinically useful diagnostic tests for polyps.

INTRODUCTION

Colorectal cancer affects ~147,000 people in the United States each year and is most effectively treated when diagnosed at an early stage (1). Colon cancers develop as the result of the transformation of normal colon epithelium to cancer via a progression of histologic changes and concurrent molecular changes that has been termed the adenoma to carcinoma progression sequence and the hyperplastic polyp-serrated adenoma to carcinoma sequence (2, 3). The adenoma to carcinoma sequence is characterized by recognizable histologic changes that start with dysplastic aberrant crypt foci and benign tubular adenomas. These lesions then have the potential to progress to advanced adenomas (characterized by size >1 cm, villous histology, and high-grade dysplasia), which have a significant potential to transform into invasive adenocarcinomas (4). The evidence that supports a hyperplastic polyp-serrated adenoma to carcinoma sequence suggests that these polyps, like adenomas, can evolve through a histologic progression sequence that culminates in adenocarcinomas of the colon, and that hyperplastic polyps that occur in the right colon and in the setting of hyperplastic polyposis have the greatest potential to transform into adenocarcinomas (5, 6).

Genetic alterations and epigenetic alterations are believed to play a pathogenic role in driving colon neoplasms through the polyp-carcinoma progression sequence (2). The most extensively studied epigenetic alteration in neoplasms, CpG island DNA methylation, has been shown to affect tumor suppressor genes during the adenoma to carcinoma and hyperplastic polyp-serrated

Received 7/29/04; revised 10/13/04; accepted 11/2/04.

Grant support: National Cancer Institute Early Detection Research Network Associate Membership, Great Lakes North East collaborative group (U01 CA094986), Damon-Runyon/Lilly Clinical Investigator Award from the Damon Runyon Cancer Research Foundation, Mallinckrodt Scholar Award from the Edward Mallinckrodt Jr. Foundation, VA Advanced Research Career Development Award from the Office of Research and Development, Medical Research Service, Department of Veterans Affairs, and Vanderbilt Physician-Scientist Development Award (to W.M. Grady), and with resources provided by the Vanderbilt Ingram Cancer Center and National Cancer Institute SPORE in GI Cancer grant 50 CA95103. J. Gorham and A. Shuber are employees of EXACT Sciences, Marlborough, MA and A. Shuber is stockholder in the company. Funding for a portion of the laboratory equipment and supplies used for experiments conducted in the laboratory of W.M. Grady was provided by EXACT Sciences. S. Markowitz is an investigator of the Howard Hughes Medical Institute.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: William M. Grady, Medical Service, Department of Veterans Affairs Puget Sound Health Care System and Division of Clinical Research, Fred Hutchinson Cancer Research Center, 1100 Fairview Avenue North, D4-100, Seattle, WA 98109. Phone: 206-667-1107; Fax: 206-667-2917; E-mail: wgrady@fhcrc.org.

©2005 American Association for Cancer Research.

adenoma to carcinoma sequences. CpG island DNA methylation is an epigenetic mechanism that represses gene transcription in a variety of normal cellular processes including imprinting and X-chromosome inactivation, but it becomes excessive and aberrant in many neoplasms. The aberrant DNA methylation affects CpG-rich regions, called "CpG islands," in the 5' region of genes and results in transcriptional silencing through complex effects on transcription factor binding and associated changes in chromatin structure. The aberrant methylation of certain tumor suppressor genes, such as *CDKN2A*, has been reported in colonic adenomas and aberrant crypt foci (7–9). These hypermethylated genes are not only probable pathogenic events in the polyp-cancer progression sequence but also are neoplasm-specific molecular events that have the potential to be used as molecular markers for pre-malignant tumors in the colon. In fact, a novel approach to colon cancer early detection that has the potential to be less expensive and better accepted by patients than currently available screening methods is the use of fecal DNA-based tests that can detect tumor-specific DNA alterations in patients with cancer or colonic adenomas (10, 11). Pilot studies have shown the feasibility of an approach using tumor-derived DNA mutations for the early detection or prevention of colon cancer; however, the use of assays based on methylated genes for the detection of colon polyps, which is the histologic stage of colon neoplasm that is most easily treated, has not been well studied to date (11, 12).

Consequently, we have developed modified, high-sensitivity methylation-specific PCR (MSP) assays for *MLH1*, *CDKN2A*, and *MGMT*, three genes previously shown to be methylated in pre-malignant neoplasms in the colon, and have applied these assays to DNA extracted from colon polyps to assess the prevalence of aberrant DNA methylation in neoplasms at different steps of the polyp-carcinoma progression sequence and to analyze the feasibility of using these methylated genes as molecular markers for colon polyps (13). We have detected methylation of these genes in adenomas and hyperplastic polyps and have found increased frequencies of methylated *CDKN2A*, *MGMT*, and *MLH1* in the subset of polyps with the greatest potential for malignant transformation. Furthermore, we have shown it is feasible to detect the methylated genes in fecal DNA using MSP-based assays.

MATERIALS AND METHODS

Patients and Collection of Tissue and Fecal DNA Samples. Sixty-six patients undergoing colonoscopy for routine clinical indications or for colon cancer screening in the Department of Veterans Affairs Tennessee Valley Health Care System (TVHCS) or at Vanderbilt University Medical Center (VUMC) at Nashville, Tennessee, were enrolled in this study over an 18-month time period using protocols approved by the Institutional Review Board of VUMC and the TVHCS. None of the patients had a clinically apparent polyposis syndrome or Hereditary Nonpolyposis Colon Cancer syndrome. Colon lavage effluent samples were collected from all patients during colonoscopy before polypectomy. After collection, the samples were kept at 4°C until being processed. Within 12 hours after collection, the samples were washed with 1× PBS and centrifuged at 1,800 *ref* for 15 minutes to pellet the solid stool. The supernatant was discarded and the pellet was stored at –80°C.

Tissues from the polyps that were resected at the time of colonoscopy were obtained from the formalin-fixed, paraffin-embedded tissue blocks stored in the pathology archives. These tissues were used for DNA analysis after the cases had undergone pathologic review. In addition, 25 normal colon specimens that were resected for non-cancer-related indications were obtained from the VUMC pathology archives. DNA was extracted from the microdissected epithelial layer of formalin-fixed, paraffin-embedded tissue sections from these normal colon specimens.

Control Cell Lines. DNA from the colon cancer cell lines VACO5 and RKO were used as controls in the studies described below. The establishment and maintenance of the cell lines has been described previously. Genomic DNA from the cell lines was extracted using previously published protocols (14, 15).

DNA Isolation from Tissue and Fecal Samples. DNA was extracted from 5 μ formalin-fixed, paraffin-embedded sections using InstaGene Matrix (Bio-Rad, Hercules, CA) as previously described (16). H&E-stained sections from each tumor sample were examined by an experienced pathologist to confirm the histological diagnosis, and the right-sided hyperplastic polyps were reviewed by an experienced gastrointestinal pathologist (M.K.W.).

For fecal DNA extraction, the fecal pellet was thawed at room temperature and resuspended in 7× TNE (EXACT Sciences, Maynard, MA). This fecal-TNE solution was centrifuged to pellet residual particulate matter, and then the supernatant was incubated at 37°C for 30 to 60 minutes after the addition of RNase (4 mg/mL; Sigma Chemical Co., St. Louis, MO). The DNA was precipitated in 1/10 volume 3 mol/L sodium acetate (Fisher Scientific, Pittsburgh, PA) and an equal volume of isopropyl-alcohol (EM Science, Gibbstown, NJ), then centrifuged and washed in 70% ethanol. After a final centrifugation, the pellet was air-dried and resuspended in 1× TE. The DNA solution was incubated at room temperature overnight and stored at –20°C.

Bisulfite Modification. Genomic DNA was modified with sodium bisulfite as previously described (13, 17). The set of known methylated and unmethylated control DNA samples used in the MSP assays was included in each round of bisulfite treatment.

Methylation Specific PCR. The bisulfite-modified DNA was subject to MSP in a blinded manner using primer pairs designed to amplify specifically the methylated or unmethylated alleles for the respective genes (Table 1). Each PCR reaction mix consisted of a total volume of 20 μL containing 10× PCR buffer (Qiagen, Valencia, CA), 200 pmol/L deoxynucleotide triphosphate mix (Sigma), 500 pmol/L of each primer (Sigma Genosys, The Woodlands, TX), 1 unit of HotStar Taq enzyme (Qiagen), and bisulfite-modified DNA. The thermocycler conditions were in general as follows: 95°C for 15 minutes, 41 cycles (45 cycles for CLE DNA) of 92°C for 30 seconds, specific annealing temperature for 30 seconds, 72°C for 30 seconds, followed by a final extension at 72°C for 10 minutes. All MSP assays were repeated at least twice to validate the results. Six control samples were included in each MSP assay run for both the methylated and unmethylated reactions and included the following: DNA from peripheral blood leukocytes, DNA extracted from colorectal cancer cell lines known to be methylated for the three target

Table 1 Primer sequence, annealing temperature, and product size for MSP assays

CpG status	Gene	Forward primer (5' → 3')	Reverse primer (5' → 3')	Annealing temperature (°C)	Product size (bp)*
M	<i>CDKN2A</i>	GGTTACGGTCGCGGTTTCG	CTAAATCGACCTCCGACCG	65.1	143
U		GTTATGGTTGTGGTTTGGGGTTG	CCACCTAAATCAACCTCCAACCA	65.1	146
M	<i>MGMT</i>	TTTCGACGTTCTGTAGGTTTTCGC	GCACTCTTCGAAAACGAAACG	62.0	121
U		TTTGTGTTTTGATGTTTTGTAGGTTTTTGT	AACTCCACACTTTCAAAAACAAAACA	62.0	133
M	<i>MLH1</i>	CGGATAGCGATTTTAAACGC	CCTAAAACGACTACTACCCG	58.5	120
U		AATGAATTAATAGGAAGAGTGGATAGT	TCTCTTCATCCCTCCCTAAAACA	57.5	136

NOTE. M, methylated; U, unmethylated.

*All of the primers were modified with a 20 bp GC-rich tail (5'-GCGGTCCCAAAAGGGTCACT-3') at their 5' end.

genes, a bisulfite treated water control, and a no template control for cross-contamination assessment. The MSP products were then subjected to horizontal gel electrophoresis through a 3% agarose gel, stained with ethidium bromide and visualized with UV transillumination using the Quantity One Image Analyzer system (Bio-Rad).

Statistical Analysis. To compare characteristics of the different groups of patients and biologic samples, *t* tests, χ^2 tests, and Fisher exact tests were used as appropriate. All statistical tests were two sided. All statistical tests were done using SAS software version 8.02 (SAS Institute, Inc., Cary, NC).

RESULTS

Aberrantly Methylated *CDKN2A*, *MGMT*, and *MLH1* Occur in Adenomas and Become More Frequent in Histologically Advanced Adenomas. DNA from 42 separate adenomas obtained from 29 patients was analyzed by methylation-specific PCR (MSP) to analyze the prevalence of methylated *CDKN2A*, *MGMT*, and *MLH1* in the adenomas. Methylated *CDKN2A*, *MGMT*, and *MLH1* were detected in 34% ($n = 14/41$), 49% ($n = 19/39$), and 7% ($n = 3/41$) of the adenomas, respectively. Fifty-nine percent ($n = 23/39$) of the adenomas that were successfully analyzed for all three genes carried at least one methylated gene. The prevalence of methylated *CDKN2A* and *MGMT* was substantially higher in the tubulovillous and villous adenomas compared with the tubular adenomas. The prevalence of methylated *CDKN2A* and methylated *MGMT* increased from 13% ($n = 3/23$) to 61% ($n = 11/18$) and from 38% ($n = 8/21$) to 61% ($n = 11/18$), respectively, in the tubular adenomas versus tubulovillous/villous adenomas. In contrast, the prevalence of methylated *MLH1* was 9% and 5% in these tubular adenomas and tubulovillous/villous adenomas, respectively (Fig. 1). The average size of the adenomas with at least one methylated gene was significantly larger than the average size of the adenomas with no methylated genes (15.6 versus 7.0 mm, respectively; $P = 0.0067$). Analysis of the frequency of the methylated genes in left and right sided adenomas revealed that 50% ($n = 11/22$), 27% ($n = 6/22$), and 4.5% ($n = 1/22$) of right sided and 47% ($n = 8/17$), 42% ($n = 8/19$), and 10.5% ($n = 2/19$) of left sided adenomas showed methylated *MGMT*, *CDKN2A*, and *MLH1*, respectively. There was no statistically significant association between any of the methylated genes and the location of the polyp(s). The overall prevalence of right sided and left sided adenomas with at least one methylated gene

was 54.5% ($n = 12/22$) and 64.7% ($n = 11/17$; $P = 0.74$), and the average number of methylated genes among right and left sided adenomas was 0.82 and 1.0, respectively.

In light of the detection of methylated *CDKN2A*, *MGMT*, and *MLH1* in tubular adenomas, we determined the methylation status of these genes in normal colonic mucosa by analyzing DNA extracted from the normal mucosal layer of 25 people who had undergone partial colectomy for benign colonic disease. None of these samples had detectable methylated *CDKN2A*, *MGMT*, or *MLH1*. This observation shows that the MSP assays we have designed do not detect low-level methylation in normal colon mucosa, which has been observed for some genes, such as *ESR*, *p14^{ARF}*, and *DAPK* (18, 19).

Aberrantly Methylated *CDKN2A*, *MGMT*, and *MLH1* Also Occur in Hyperplastic Polyps. In addition to the analysis of the adenomas, 44 hyperplastic polyps from 17 patients who had hyperplastic polyps were collected and analyzed. None of the individuals in this part of the study met the clinical criteria for hyperplastic polyposis. Interestingly, in the hyperplastic polyps, aberrantly methylated *CDKN2A*, *MGMT*, and *MLH1* were detected in 5% ($n = 2/40$), 10% ($n = 4/40$), and 7.5% ($n = 3/40$) of the polyps, respectively (Fig. 1). Twenty percent of all the hyperplastic polyps that could




Fig. 1 Prevalence of methylated *CDKN2A*, *MGMT*, and *MLH1* in normal colonic mucosa, hyperplastic polyps (HP), tubular adenomas (TA), and tubulovillous and villous adenomas (TVA and VA) from patients undergoing colonoscopy for routine clinical indications, including colon cancer screening. The histologically normal colonic mucosa was obtained from surgical resection specimens from individuals with no neoplastic disease and showed no detectable methylation of the genes studied.

be analyzed for all three genes ($n = 8/39$) carried at least one methylated gene. Among the 17 patients with hyperplastic polyps, 35% ($N = 6/17$) had methylated gene(s) detected in their polyps. In the hyperplastic polyp group, the average size of the polyps with and without at least one methylated gene was 3.4 and 3.8 mm, respectively. These polyps were located throughout the colon but were almost exclusively from the left colon. Of note, the likelihood of identifying a polyp with a methylated gene was independent of the number of hyperplastic polyps present.

Aberrantly Methylated Genes in Colon Polyps Can Be Detected in Fecal DNA. Following the performance of the MSP assays on the DNA extracted from the polyps, we assessed fecal DNA from a group of study subjects who had matched fecal and polyp DNA samples to determine if we could detect methylated genes in the fecal DNA from individuals with colon polyps. Sixteen study subjects had an adenoma with methylated *CDKN2A* and eight (50%) of these cases had methylated *CDKN2A* in the fecal DNA as well. Nine subjects did not have methylated *CDKN2A* in either their adenomas or fecal samples, and one had methylated *CDKN2A* only in his fecal DNA but not in his adenoma. For *MGMT*, 12 of 17 study subjects had concordant methylated *MGMT* in their adenoma DNA and fecal DNA. Nine subjects had no methylated *MGMT* in either their adenomas or fecal samples, and two subjects had methylated *MGMT* only in the fecal DNA. None of the three individuals whose adenomas carried methylated *MLH1* had detectable methylated *MLH1* in their fecal samples, and all of the remaining 25 subjects with adenomas that had unmethylated *MLH1* showed unmethylated *MLH1* in their fecal samples. Comparison of the performance characteristics of *CDKN2A* and *MGMT* MSP assays showed that the assays could detect 50% and 71% of individuals with adenomas that carried methylated *CDKN2A* or *MGMT*, respectively. Overall, methylation of at least one of the candidate genes was detected in the fecal DNA from 55% of patients with adenoma(s) ($N = 16/29$), and 14 of these individuals had at least one gene that showed matching methylation in the adenoma(s) and fecal DNA (Table 2; Fig. 2).

With regards to the study subjects with hyperplastic polyps, we restricted our analysis to individuals who only had hyperplastic polyps to simplify the interpretation of our results.

Ten study subjects with matched fecal and polyp DNA were found to only have hyperplastic polyps during colonoscopy. Two of these individuals had methylated *MGMT* in their polyps, and both of these individuals tested positive for methylated *MGMT* in their fecal DNA. Only two individuals had methylated *CDKN2A* in a hyperplastic polyp, and one of these two cases had detectable methylated *CDKN2A* in his fecal DNA. One additional subject had methylated *CDKN2A* in the fecal sample but had no detectable methylated *CDKN2A* in any of his polyps (Table 2).

After detecting methylated genes in the fecal DNA of individuals with colon polyps, fecal DNA samples from another 25 individuals without detectable polyps during colonoscopy were also tested for the presence of methylated *CDKN2A*, *MGMT*, or *MLH1*. The methylation analysis revealed that 37% ($N = 7/19$) of these individuals had methylated genes detected in their fecal DNA despite having no polyps detected during colonoscopy (Table 2). The MSP assays failed to generate products from the DNA of six subjects with normal colonoscopic exams, most likely because of an insufficient amount of DNA in their fecal samples.

DISCUSSION

We have identified aberrantly methylated genes in colorectal polyps and found that over 59% of adenomas carry at least one methylated allele of *MGMT*, *CDKN2A*, or *MLH1*. Notably, the prevalence of methylated *MGMT* and *CDKN2A*, but not *MLH1*, is higher in the more histologically progressed adenomas. We have also found that aberrant DNA methylation occurs in hyperplastic polyps although significantly less frequently than it does in adenomas. Finally, we have now shown for the first time to our knowledge the results of studies assessing the use of methylated genes as fecal DNA-based molecular markers for colon polyps.

Our results support the observation that the aberrant methylation of genes occurs early in the tumorigenesis process in the gastrointestinal tract, arising in the premalignant stage of these colon neoplasms. Consistent with our observation, aberrant CpG island methylation has been detected by some investigators in the precursor lesions for colon cancer, including aberrant crypt




Fig. 2 Methylation analysis of *CDKN2A*, *MGMT*, and *MLH1* in matched polyp-fecal DNA pairs by methylation-specific PCR using primers for methylated (*M*) and unmethylated (*U*) alleles of bisulfite-modified DNA. The gene analyzed and case numbers are indicated above each gel. DNA from lymphocytes (*N1* and *N2*), and colon cancer cell lines [RKO and VACO5 (*V5*)] was used for unmethylated and methylated controls, respectively, and distilled UV light-treated water without DNA was used as a negative control. The unmethylated alleles detected in the polyp and fecal DNA are presumed secondary to contaminating DNA from normal cells.

Table 2 Correlation between colonoscopic findings and tissue and fecal DNA MSP assay results

Endoscopic findings	Case ID	Genes and DNA sources		
		<i>CDKN2A</i> , tissue/feces	<i>MGMT</i> , tissue/feces	<i>MLH1</i> , tissue/feces
Normal colonoscopy (no polyps)	1n	-/U	-/U	-/U
	2n	-/U	-/U	-/U
	3n	-/U	-/U	-/U
	4n	-/U	-/U	-/U
	5n	-/U	-/U	-/U
	6n	-/U	-/U	-/U
	7n	-/U	-/U	-/U
	8n	-/U	-/na	-/U
	9n	-/U	-/U	-/U
	10n	-/U	-/U	-/U
	11n	-/U	-/U	-/U
	12n	-/U	-/U	-/U
	13n	-/U	-/U	-/M
	14n	-/M	-/U	-/U
	15n	-/U	-/M	-/U
	16n	-/U	-/M	-/U
	17n	-/U	-/M	-/U
	18n	-/M	-/M	-/U
	19n	-/M	-/M	-/M
	20n	-/na	-/na	-/na
	21n	-/na	-/na	-/na
	22n	-/na	-/na	-/na
	23n	-/na	-/na	-/na
	24n	-/na	-/na	-/na
	25n	-/na	-/na	-/na
Hyperplastic polyps	1h	na/na	na/U	na/U
	2h	U/na	U/U	U/U
	3h	U/U	U/U	U/U
	4h	U/U	U/U	U/U
	5h	U/na	U/U	U/U
	6h	U/U	U/U	U/U
	7h	U/M	U/U	U/U
	8h	U/U	M/M	U/U
	9h	U/U	M/M	U/U
	10h	M/U	M/M	U/U
Adenomas	1a	U/na	U/na	U/na
	2a	U/U	U/U	U/U
	3a	U/U	U/U	U/U
	4a	U/U	U/U	U/U
	5a	U/U	U/U	U/U
	6a	U/U	U/U	U/U
	7a	U/na	U/M	U/U
	8a	U/U	U/M	U/U
	9a	M/U	U/U	U/U
	10a	M/U	U/U	U/U
	11a	U/U	M/U	U/U
	12a	na/U	M/U	M/U
	13a	M/U	M/U	U/U
	14a	M/U	M/U	U/U
	15a	M/U	M/U	U/U
	16a	M/M	U/U	U/U
	17a	M/M	U/U	U/U
	18a	U/U	M/M	U/U
	19a	U/U	M/M	M/U
	20a	U/M	M/M	U/U
21a	M/U	M/M	U/U	
22a	M/U	M/M	U/U	
23a	M/U	M/M	U/U	
24a	M/M	M/M	U/U	
25a	M/M	M/M	U/U	
26a	M/M	M/M	U/U	
27a	M/M	M/M	U/U	
28a	M/M	M/M	U/U	
29a	M/M	M/M	M/U	

foci and adenomas (7–9, 20–23). In fact, Chan et al. (8) showed that methylated *MINT1*, *MINT31*, and *MGMT* could be found in sporadic dysplastic aberrant crypt foci, which are the most likely precursors of adenomas, suggesting that aberrant DNA methylation is present at the earliest histologically detectable stage of colon cancer formation. In light of these previous findings, we focused our analysis on three genes that likely play a role in driving the initiation and promotion of colon adenomas, *CDKN2A*, *MGMT*, and *MLH1*. We have found that methylated *CDKN2A* and *MGMT* can be found in 34% and 49% of adenomas and that they increase in frequency in histologically progressed adenomas (7, 24). These findings suggest that methylation of these two genes may contribute to the progression of the polyps. On the contrary, methylated *MLH1* was found in 9% and 5% of tubular adenomas and tubulovillous/villous adenomas, respectively, which, in light of the 10% to 20% frequency of methylated *MLH1* observed in colon cancers, suggests that methylation of *MLH1* may contribute to the rapid progression to colon cancer (9, 20). Indeed, the marked increase in the frequency of methylated *MLH1* from tubular adenomas to cancers is in contrast to the stepwise increase observed for *CDKN2A* and *MGMT* and would be congruent with methylation of *MLH1* being associated with a rapid polyp-cancer progression sequence.

Notably, our results regarding the methylation status of colon adenomas are consistent with those of Rashid and Esteller but contradict those of Lee and Bariol who found lower frequencies of *CDKN2A* methylation in colon adenomas (7, 9, 20, 24). There are multiple potential reasons for these differences including assay-related differences and differing patient populations. It is notable that the two studies done on populations outside of the United States showed substantially less methylation of *CDKN2A* and *MGMT*, suggesting the possibility of environmental factors having a role in the pattern of aberrant DNA methylation observed in colon neoplasms. Also, it is important to note the degree of methylation present for a gene can vary across the 5' part of the gene, which can result in variable results for different MSP assays that are assessing the same gene but not necessarily the same location in the 5' part of the gene (17, 20, 25). It is conceivable that differences in the location of the annealing sites for the MSP primers used by the investigators noted above may account for the different results.

In addition to analyzing adenomas, we have also assessed sporadic hyperplastic polyps that were found in the study subjects at the time of colonoscopy. Aberrantly methylated *CDKN2A*, *MINT1*, *MINT2*, and *MINT31* have been observed in hyperplastic polyps but the hyperplastic polyps assessed in these studies have mainly been those occurring in the setting of hyperplastic polyposis or concurrent cancer (5, 26). We have assessed a

NOTE. MSP assay results from tissue samples and fecal DNA are shown respectively for study subjects with endoscopically detected polyps. Subjects were placed into the group with hyperplastic polyps if they only had hyperplastic polyps and were placed into the group with adenomas if any of the polyps they had detected on endoscopy included an adenoma. If an individual had more than one polyp and any of these polyps had a methylated allele, the polyp DNA was scored as positive. Subjects with normal colonoscopic findings have single methylation results obtained from analysis of DNA from the fecal material.

Abbreviations: U, unmethylated; M, methylated; CRC, colorectal cancer; na, no PCR product generated.

collection of sporadic hyperplastic polyps and, consistent with the results of Bariol et al, we found methylated *CDKN2A*, *MGMT*, and *MLH1* in a subset of these hyperplastic polyps (20, 26). In light of recent studies that have suggested that some serrated polyps have the potential to transform into cancer and that aberrantly methylated genes are commonly found in serrated polyps, it is interesting to speculate that hyperplastic polyps that have aberrantly methylated genes may have the potential to transform (5, 6). Further studies will be needed to determine if such a causal relationship exists between these two observations.

Finally, based on our findings in the adenomas and hyperplastic polyps, we have studied the use of aberrantly methylated genes as fecal-based molecular markers for colon adenomas, owing to these being the polyps that have the best-demonstrated risk for transformation. We have shown that an MSP-based assay panel employing two genes, *CDKN2A* and *MGMT*, when applied to DNA from fecal material, identified 55% of the patients with adenomatous polyps. Notably, not all adenomas carry the methylated gene being detected by the MSP assays we used, suggesting that adding more MSP assays that assess genes other than *CDKN2A* and *MGMT* may improve the overall performance of the MSP assay panel to detect colon adenomas (27). In light of the known heterogeneity of molecular alterations in adenomas, this approach should improve the sensitivity of MSP assays to detect adenoma DNA from fecal material. However, it is important to note that the addition of MSP assays to an assay panel carries the potential to decrease the specificity of the assay panel and requires the selection of highly specific individual assays to generate an assay panel with a high positive predictive value for adenomas (28). Also, the positive association that we and others have found between the frequency of aberrant DNA methylation and advanced histological features suggests that MSP assays may be more accurate for the detection of advanced adenomas and colon cancer than for polyps in general.

We also found that this assay panel only detected 55% of patients with colon adenomas and detected methylated *CDKN2A* or *MGMT* in 37% of people with no polyps found by colonoscopy. We speculate that the MSP assays used in this study have suboptimal sensitivity because the PCR-based assays we have used either are not sensitive enough to detect the small amount of polyp DNA in the fecal DNA samples of the false-negative cases or because the technique we have used to extract the fecal DNA has resulted in loss of the tumor DNA in the samples that generated false-negative results. In addition, an important question raised by our results is the significance of the detection of methylated genes in 37% of the fecal DNA samples from patients with normal colonoscopic exams, which is in contrast with the complete lack of methylated genes detected in the normal mucosal samples obtained from a separate control group that was age-matched. Possible explanations for this finding include technical errors, differences between the populations from whom we obtained normal mucosa samples and from whom we obtained fecal DNA samples after negative colonoscopic exams, low-level DNA methylation in the normal tissue of some people, and false-negative colonoscopic exams due to colonoscopy having at least a 28% miss rate for small adenomas (19, 29–33). Furthermore, it is possible that the methylated genes we detected in the fecal DNA were derived

from the normal colon mucosa or aberrant crypt foci in the study subjects rather than from the polyps, which could explain the low sensitivity and specificity we observed (8, 19). The design of our study did not permit us to investigate these possible causes for the false-positive results.

In summary, we have shown that aberrantly methylated *CDKN2A*, *MGMT*, and *MLH1* occur in colon adenomas and hyperplastic polyps and that the frequency of methylated genes is higher in histologically advanced adenomas, which are believed to have the highest likelihood of progressing to colon cancer. We have also shown the feasibility of using aberrantly methylated genes detected in fecal material as molecular markers for these polyps. Our findings suggest that the detection of aberrantly methylated genes in fecal material from colon lavage effluent, and likely from stool, has potential value in the noninvasive and early diagnosis of colorectal neoplasms, but that the low sensitivity and specificity of the assays we have used necessitates further assay development before this approach can be clinically useful as a diagnostic assay for colon polyps. The identification of additional genes that are frequently, and not concurrently, methylated in polyps may improve the performance characteristics of the current assay panel. In light of the success of molecular markers that detect DNA mutations and the known genetic and epigenetic heterogeneity of colon cancer, it is possible that molecular marker assays that employ both genetic and epigenetic alterations could be effective for the early detection of colon cancer (34–37).

ACKNOWLEDGMENTS

We thank Caroline Gilmore, Stephen Vaughn, the staff in the Nashville VA GI endoscopy lab, and Colin Crosby for their efforts with helping in patient recruitment, assistance during colonoscopy and clinical record management, and performance of the molecular laboratory assays.

REFERENCES

1. Jemal A, Murray T, Samuels A, et al. Cancer statistics, 2003. *CA Cancer J Clin* 2003;53:5–26.
2. Grady WM, Markowitz SD. Genetic and epigenetic alterations in colon cancer. *Annu Rev Genomics Hum Genet* 2002;3:101–28.
3. Jass JR, Young J, Leggett BA. Evolution of colorectal cancer: change of pace and change of direction. *J Gastroenterol Hepatol* 2002;17:17–26.
4. Kinzler K, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;87:159–70.
5. Wynter CV, Walsh MD, Higuchi T, et al. Methylation patterns define two types of hyperplastic polyp associated with colorectal cancer. *Gut* 2004;53:573–80.
6. Goldstein NS, Bhanot P, Odish E, Hunter S. Hyperplastic-like colon polyps that preceded microsatellite-unstable adenocarcinomas. *Am J Clin Pathol* 2003;119:778–96.
7. Rashid A, Shen L, Morris JS, et al. CpG island methylation in colorectal adenomas. *Am J Pathol* 2001;159:1129–35.
8. Chan AO, Broaddus RR, Houlihan PS, et al. CpG island methylation in aberrant crypt foci of the colorectum. *Am J Pathol* 2002;160:1823–30.
9. Lee S, Hwang KS, Lee HJ, et al. Aberrant CpG island hypermethylation of multiple genes in colorectal neoplasia. *Lab Invest* 2004;84:884–93.
10. Song K, Fendrick AM, Ladabaum U. Fecal DNA testing compared with conventional colorectal cancer screening methods: a decision analysis. *Gastroenterology* 2004;126:1270–9.
11. Ahlquist DA, Skoletsky JE, Boynton KA, et al. Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a

- multitarget assay panel. *Gastroenterology* 2000;119:1219–27.
12. Muller HM, Oberwalder M, Fiegl H, et al. Methylation changes in faecal DNA: a marker for colorectal cancer screening? *Lancet* 2004;363:1283–5.
 13. Herman J, Graff J, Myohanen S, et al. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci U S A* 1996;93:9821–6.
 14. Markowitz S, Wang J, Myeroff L, et al. Inactivation of the type II TGF- β receptor in colon cancer cells with microsatellite instability. *Science* 1995;268:1336–8.
 15. Willson J, Bittner G, Oberley T, et al. Cell culture of human colon adenomas and carcinomas. *Cancer Res* 1987;47:2704–13.
 16. Grady W, Rajput A, Myeroff L, et al. Mutation of the type II transforming growth factor- β receptor is coincident with the transformation of human colon adenomas to malignant carcinomas. *Cancer Res* 1998;58:3101–4.
 17. Grady WM, Rajput A, Lutterbaugh J, Markowitz S. Detection of aberrantly methylated hMLH1 promoter DNA in the serum of patients with microsatellite unstable colon cancer. *Cancer Res* 2001;61:900–2.
 18. Shen L, Kondo Y, Hamilton SR, et al. P14 methylation in human colon cancer is associated with microsatellite instability and wild-type p53. *Gastroenterology* 2003;124:626–33.
 19. Ahuja N, Li Q, Mohan M, et al. Aging and DNA methylation in colorectal mucosa and cancer. *Cancer Res* 1998;58:5489–94.
 20. Bariol C, Suter C, Cheong K, et al. The relationship between hypomethylation and CpG island methylation in colorectal neoplasia. *Am J Pathol* 2003;162:1361–71.
 21. Li H, Myeroff L, Smiraglia D, et al. SLC5A8, a sodium transporter, is a tumor suppressor gene silenced by methylation in human colon aberrant crypt foci and cancers. *Proc Natl Acad Sci U S A* 2003;100:8412–7.
 22. Moinova HR, Chen WD, Shen L, et al. HLF gene silencing in human colon cancer. *Proc Natl Acad Sci U S A* 2002;99:4562–7.
 23. Suzuki H, Watkins DN, Jair KW, et al. Epigenetic inactivation of SFRP genes allows constitutive WNT signaling in colorectal cancer. *Nat Genet* 2004;36:417–22.
 24. Esteller M. Epigenetic lesions causing genetic lesions in human cancer. promoter hypermethylation of DNA repair genes. *Eur J Cancer* 2000;36:2294–300.
 25. Deng G, Chen A, Hong J, et al. Methylation of CpG in a small region of the hMLH1 promoter invariably correlates with the absence of gene expression. *Cancer Res* 1999;59:2029–33.
 26. Chan AO, Issa JP, Morris JS, et al. Concordant CpG island methylation in hyperplastic polyposis. *Am J Pathol* 2002;160:529–36.
 27. Laird PW. The power and the promise of DNA methylation markers. *Nat Rev Cancer* 2003;3:253–66.
 28. Lofton-Day C, Lesche R. DNA methylation markers in patients with gastrointestinal cancers. Current understanding, potential applications for disease management and development of diagnostic tools. *Dig Dis* 2003;21:299–308.
 29. Nakagawa H, Nuovo GJ, Zervos EE, et al. Age-related hypermethylation of the 5' region of MLH1 in normal colonic mucosa is associated with microsatellite-unstable colorectal cancer development. *Cancer Res* 2001;61:6991–5.
 30. Palmisano WA, Divine KK, Saccomanno G, et al. Predicting lung cancer by detecting aberrant promoter methylation in sputum. *Cancer Res* 2000;60:5954–8.
 31. Belinsky SA, Palmisano WA, Gilliland FD, et al. Aberrant promoter methylation in bronchial epithelium and sputum from current and former smokers. *Cancer Res* 2002;62:2370–7.
 32. Rex DK, Cutler CS, Lemmel GT, et al. Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies. *Gastroenterology* 1997;112:24–8.
 33. Hixson LJ, Fennerty MB, Sampliner RE, et al. Prospective study of the frequency and size distribution of polyps missed by colonoscopy. *J Natl Cancer Inst* 1990;82:1769–72.
 34. Hermsen M, Postma C, Baak J, et al. Colorectal adenoma to carcinoma progression follows multiple pathways of chromosomal instability. *Gastroenterology* 2002;123:1109–19.
 35. Traverso G, Shuber A, Levin B, et al. Detection of APC mutations in fecal DNA from patients with colorectal tumors. *N Engl J Med* 2002;346:311–20.
 36. Ahlquist DA, Skoletsky JE, Boynton KA, et al. Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a multitarget assay panel. *Gastroenterology* 2000;119:1219–27.
 37. Dong SM, Traverso G, Johnson C, et al. Detecting colorectal cancer in stool with the use of multiple genetic targets. *J Natl Cancer Inst* 2001;93:858–65.

Clinical Cancer Research

Aberrantly Methylated *CDKN2A*, *MGMT*, and *MLH1* in Colon Polyps and in Fecal DNA from Patients with Colorectal Polyps

Zsolt Petko, Mahan Ghiassi, Anthony Shuber, et al.

Clin Cancer Res 2005;11:1203-1209.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/11/3/1203>

Cited articles This article cites 35 articles, 12 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/11/3/1203.full#ref-list-1>

Citing articles This article has been cited by 19 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/11/3/1203.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/11/3/1203>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.