

Cytochrome P450 2D6 and Homeobox 13/Interleukin-17B Receptor: Combining Inherited and Tumor Gene Markers for Prediction of Tamoxifen Resistance

Matthew P. Goetz,^{1,4} Vera J. Suman,² Fergus J. Couch,³ Matthew M. Ames,⁴ James M. Rae,⁵ Mark G. Erlander,⁶ Xiao-Jun Ma,⁶ Dennis C. Sgroi,⁷ Carol A. Reynolds,³ Wilma L. Lingle,³ Richard M. Weinshilboum,⁴ David A. Flockhart,⁸ Zeruesenay Desta,⁸ Edith A. Perez,⁹ and James N. Ingle¹

Abstract Purpose: Genetic variation in *cytochrome P450 2D6* (*CYP2D6*) and the gene expression ratio of the homeobox 13 (*HOXB13*) to interleukin-17B receptor (*IL17BR*) are associated with tamoxifen resistance. We sought to determine the combined effect of inherited (*CYP2D6*) and somatic (*HOXB13/IL17BR*) gene variation in tamoxifen-treated breast cancer.

Experimental Design: Retrospective analysis of women with node-negative breast cancer randomized to receive 5 years of tamoxifen (North Central Cancer Treatment Group 89-30-52). *CYP2D6* metabolism (extensive or decreased) was based on *CYP2D6**4 genotype and presence/absence of a *CYP2D6* inhibitor. Reverse transcription-PCR profiles for *HOXB13* and *IL17BR* and the cut point separating patients into high- and low-risk categories according to disease-free survival (DFS) were used. A risk factor (*CYP2D6:HOXB13/IL17BR*) representing the four categories of combining *CYP2D6* metabolism (extensive or decreased) and *HOXB13/IL17BR* (low or high) was created. The association between *CYP2D6:HOXB13/IL17BR* and DFS and overall survival (OS) was assessed using the log-rank test and proportional hazards modeling.

Results: *CYP2D6* metabolism and *HOXB13/IL17BR* gene ratio was available in 110 of 160 (69%) patients. The combined *CYP2D6:HOXB13/IL17BR* risk factor was significantly associated with DFS (log-rank $P = 0.004$) and OS ($P = 0.009$). Relative to women with extensive *CYP2D6* metabolism and low *HOXB13/IL17BR*, those with either decreased metabolism or a high *HOXB13/IL17BR* ratio had significantly worse OS (adjusted hazard ratio, 2.41; 95% confidence interval, 1.08-5.37; $P = 0.031$), whereas women with both decreased metabolism and high *HOXB13/IL17BR* ratio had the shortest survival (adjusted hazard ratio, 3.15; 95% CI, 1.17-8.52; $P = 0.024$).

Conclusions: An index composed of inherited (*CYP2D6*) and tumor (*HOXB13/IL17BR*) gene variation identifies patients with varying degrees of resistance to tamoxifen.

Genetic markers are increasingly used to identify patients at increased risk for relapse (prognosis) and to predict the likelihood of a therapeutic response (predictive markers). Given that both tamoxifen and aromatase inhibitors are

effective therapies for the treatment of estrogen receptor (ER)-positive breast cancer (1, 2), there is great interest in identifying biomarkers that may allow the individualization of therapy.

Authors' Affiliations: Departments of ¹Oncology, ²Biostatistics, ³Pathology, and ⁴Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic College of Medicine, Rochester, Minnesota; ⁵Division of Hematology/Oncology, Department of Medicine, University of Michigan School of Medicine, Ann Arbor, Michigan; ⁶AviaraDx, San Diego, California; ⁷Department of Pathology, Harvard Medical School, Molecular Pathology Research Unit, Massachusetts General Hospital, Boston, Massachusetts; ⁸Division of Clinical Pharmacology, Department of Medicine, Indiana University, Indianapolis, Indiana; and ⁹Division of Hematology and Oncology, Department of Medicine, Mayo Clinic, Jacksonville, Florida. Received 3/7/08; revised 5/14/08; accepted 5/15/08.

Grant support: Paul Calabresi Program in Clinical-Translational Research at Mayo Clinic grant CA 90628 (M.P. Goetz), Mayo Clinic Breast Cancer Specialized Program of Research Excellence grant CA116201 (J.N. Ingle, F.J. Couch, M.P. Goetz, V.J. Suman, and C.A. Reynolds), Mayo Comprehensive Cancer Center grant CA 15083 (M.M. Ames), CA87898 (F.J. Couch), Breast Cancer Research Foundation (J.M. Rae and E.A. Perez), National Institute of General Medical Sciences (Bethesda, MD) grant U-01 GM61388 (R.M. Weinshilboum, M.P. Goetz, and J.N. Ingle), and CA-25224 (North Central Cancer Treatment Group).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

This work was presented in part at the San Antonio 2006 Breast Cancer Symposium.

All study funding was from public grants for scientific research. The funding organizations had no role in the design and conduct of the study; the collection, analysis, and interpretation of the data; or the preparation, review, or approval of the manuscript.

M.P. Goetz had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Requests for reprints: Matthew P. Goetz, Department of Oncology, Mayo Clinic, 200 First Street Southwest, Rochester, MN 55905. Phone: 507-284-4857; Fax: 507-284-1803; E-mail: goetz.matthew@mayo.edu.

©2008 American Association for Cancer Research.

doi:10.1158/1078-0432.CCR-08-0619

Cytochrome P450 2D6 (CYP2D6) is a polymorphic enzyme responsible for the metabolic activation of tamoxifen to metabolites (endoxifen and 4-OH tamoxifen) with a significantly greater affinity for the ER and greater ability to inhibit tumor cell proliferation compared with the parent drug (tamoxifen) or its primary metabolite, *N*-desmethyltamoxifen (3, 4). Both genetic (inherited traits) and environmental (drug induced) factors that alter CYP2D6 enzyme activity affect the concentrations of the active tamoxifen metabolites (5, 6). We and others have shown the importance of CYP2D6 as a predictor of tamoxifen benefit in the prevention (7), metastatic (8), and adjuvant settings (9–11). Based on these data, a Food and Drug Administration Clinical Pharmacology Subcommittee recently recommended that the tamoxifen label should be updated to reflect the increased risk of breast cancer recurrence for women who are CYP2D6 poor metabolizers, resulting from genetic variation or drug-induced inhibition of CYP2D6.

In a genome-wide microarray analysis of tumors from women with ER-positive breast cancer treated with adjuvant tamoxifen, Ma et al. (12) discovered the *homeobox 13* (*HOXB13*)/*interleukin-17B receptor* (*IL17BR*) gene ratio as an independent predictor of treatment outcome and showed that ectopic expression of *HOXB13* in a nontransformed human mammary epithelial cell confers increased cell migration and invasion. Follow-up studies have shown that both *HOXB13* and *IL17BR* are regulated by estradiol in an ER-dependent manner and this regulation is abrogated by tamoxifen (13). In hormone-sensitive cell lines, *HOXB13* expression rendered cells less sensitive to tamoxifen-induced apoptosis (14). Further clinical analysis of ~2,700 patients from four independent clinical databases (15–18) has repeatedly validated the *HOXB13/IL17BR* ratio as a prognostic marker (18) and a biomarker predictive of tamoxifen benefit (15–17), with the effect most pronounced in node-negative, ER-positive breast cancer (15, 16, 18).

In a cohort of postmenopausal women with resected ER-positive breast treated with adjuvant tamoxifen only, we reported that women with decreased CYP2D6 metabolism (defined as the presence of one or two *CYP2D6*4* alleles or the documented coprescription of a CYP2D6 inhibitor) had significantly shorter disease-free survival (DFS) compared with women with extensive CYP2D6 metabolism (defined as women who did not carry a *CYP2D6*4* allele and were not coprescribed a CYP2D6 inhibitor; ref. 9). In this same cohort, we showed that a high *HOXB13/IL17BR* ratio was associated with significantly worse DFS and overall survival (OS) but only in lymph node–negative patients (15).

We hypothesized that because CYP2D6 and *HOXB13/IL17BR* represent biologically independent markers of tamoxifen resistance, a combined index may provide a better indicator of tamoxifen treatment benefit. Therefore, using the same study population from which we derived our original findings about CYP2D6 (9, 10) and *HOXB13/IL17BR* (15), we evaluated the combined effect of CYP2D6 and *HOXB13/IL17BR* on the outcomes of DFS and OS in women with lymph node–negative breast cancer treated with adjuvant tamoxifen monotherapy.

Materials and Methods

Patients. The North Central Cancer Treatment Group (NCCTG) conducted a randomized phase III clinical trial in postmenopausal women with resected ER-positive breast cancer to assess the value of administering the androgen fluoxymesterone for 1 y during the standard 5 y of tamoxifen adjuvant therapy (NCCTG 89-30-52). A description of the clinical trial and its outcome has been published (19). The protocol was amended to assess genetic and environmental factors, which may be associated with increased risk of breast cancer relapse. The trial and its amendments were approved by the

Table 1. Baseline characteristics

Characteristic	HOXB13/IL17BR and CYP2D6 phenotype available (n = 110)	HOXB13/IL17BR and CYP2D6 phenotype not available (n = 50)
Median age (range)	65 (42-84)	63 (47-82)
Race		
Caucasian	105 (95%)	42 (84%)
African-American	1 (1%)	1 (2%)
Unknown	4 (4%)	7 (14%)
Extent of surgery		
Mastectomy	91(83%)	38 (76%)
Breast sparing	19 (17%)	12 (24%)
ER status		
10-49 fmol	27 (25%)	7 (14%)
≥50 fmol	65 (59%)	34 (68%)
Positive	18 (16%)	9 (18%)
Tumor size ≥3 cm	22 (20%)	7 (14%)
Tumor grade		
1-2	86 (78%)	22 (44%)
3	17 (15%)	5 (10%)
Unknown	7 (6%)	23 (46%)
Immunohistochemical HER2 expression		
0	13 (12%)	2 (4%)
1+	34 (31%)	10 (20%)
2+	35 (32%)	9 (18%)
3+	21 (19%)	6 (12%)
Unknown	7 (6%)	23 (46%)

Table 2. Risk groups as defined by CYP2D6 metabolism and HOXB13/IL17BR gene ratio

Risk group	HOXB13/IL17BR gene ratio	CYP2D6 metabolism phenotype (CYP2D6*4 genotype: CYP2D6 inhibitor use)	n = 110	Estimated 5-y DFS rate (95% CI)
1	<-1.339	Extensive (Wt/Wt: no)	46 (46)	93.5% (86.6-99.9%)
2	<-1.339	Decreased (Unknown: yes) (Wt/Wt: yes) (Wt/*4: yes) (Wt/*4: no) (*4/*4: no)	18 (2) (5) (1) (9) (1)	83.8% (67.8-99.9%)
3	≥-1.339	Extensive (Wt/Wt: no)	32 (32)	75.0% (61.4-91.6%)
4	≥-1.339	Decreased (Wt/Wt: yes) (Wt/*4: no) (*4/*4: no)	14 (1) (11) (2)	57.1% (36.3-89.9%)

NOTE: The Kaplan-Meier estimates of 5-y DFS rates and corresponding CIs for each group are given. CYP2D6 inhibitors coadministered during tamoxifen in nine patients were as follows: cimetidine ($n = 4$), fluoxetine ($n = 2$), paroxetine ($n = 2$), sertraline ($n = 1$), and haloperidol ($n = 1$); one patient was coprescribed two CYP2D6-inhibiting drugs.

institutional review board of Mayo Clinic Rochester and the individual NCCCTG sites that enrolled patients onto the clinical trial.

The means by which CYP2D6 genotyping (10) and HOXB13/IL17BR (15) were measured and quantified has been previously published. Additionally, we published a comprehensive analysis of both CYP2D6 genotype and CYP2D6 enzyme inhibition on tamoxifen treatment outcome (9). Briefly, a woman who carried either one or two CYP2D6*4 alleles or had any CYP2D6 genotype but coprescribed a CYP2D6 inhibitor was considered to have decreased CYP2D6 metabolism. A woman without a CYP2D6*4 allele and who was not coprescribed a CYP2D6 inhibitor was considered to have extensive metabolism. A woman was said to have a low HOXB13/IL17BR gene ratio if her tumor HOXB13/IL17BR gene ratio fell below <-1.339 (15).

Study design and end points. The primary objective of this study was to assess the association of effect of CYP2D6 metabolism and the HOXB13/IL17BR gene ratio on DFS and OS in women with lymph node-negative breast cancer receiving adjuvant tamoxifen only. DFS was defined as the time from randomization to documentation of the first of the following events: any recurrence (local, regional, or distant) of breast cancer, a contralateral breast cancer, a second primary cancer, or death due to any cause. Patients who were alive without any of these events were censored at the date of their last disease evaluation. OS was estimated as the time from randomization to death due to any cause.

The overall distributions of DFS and OS were estimated using the Kaplan-Meier method. Log-rank tests and univariate proportional hazard models were used to assess whether the end point differed with respect to any one of the following factors: age >65 (yes versus no), ER status at time of entry into the trial (10-49 fmol versus ≥50 fmol versus positive by immunohistochemistry), tumor size ≥3 cm (yes versus no), Nottingham grade (3 versus 1 or 2), and HER2 expression (3+ versus 0, 1+, or 2+). For each clinical outcome, multivariate proportional hazard modeling was done to obtain a subset of the potential prognostic factors, which provided an adequate fit to the data. Residual plots were examined. A risk factor (CYP2D6:HOXB13/IL17BR) representing the four categories of combining CYP2D6 metabolism (extensive versus decreased) and HOXB13/IL17BR (low versus high) was created. Representing CYP2D6:HOXB13/IL17BR as three indicator variables, multivariate proportional hazard modeling was done to assess whether any one of these factors made a significant contribution to the previous established model for that clinical end point.

Results

Our study population is derived from women with node-negative breast cancer ($n = 160$) who were randomized from December 31, 1990 to April 6, 1995 to the tamoxifen only arm of NCCCTG 89-30-52. In 110 of 160 patients, both HOXB13/IL17BR gene expression data and a comprehensive assessment of CYP2D6 metabolism (consisting of CYP2D6*4 genotype and medication history) were known. The characteristics of these 110 patients are listed in Table 1 and the combined CYP2D6 and HOXB13/IL17BR phenotype is listed in Table 2.

As of October 25, 2006, 12 women were alive with disease progression, a second primary, or contralateral breast disease, and 37 women had died. The median length of follow-up for those women still alive is 12.5 years (range, 5.7-15.5 years).




Fig. 1. Kaplan-Meier estimates of DFS in lymph node-negative patients according to the presence of no high-risk features (extensive CYP2D6 metabolism and HOXB13/IL17BR <-1.339), only one high-risk feature (either decreased CYP2D6 metabolism or HOXB13/IL17BR ≥-1.339), and two high-risk features (both decreased CYP2D6 metabolism and HOXB13/IL17BR ≥-1.339).




Fig. 2. Kaplan-Meier estimates of OS in lymph node-negative patients according to the presence of no high-risk features (extensive CYP2D6 metabolism and *HOXB13/IL17BR* <-1.339), only one high-risk feature (either decreased CYP2D6 metabolism or *HOXB13/IL17BR* \geq -1.339), and two high-risk features (both decreased CYP2D6 metabolism and *HOXB13/IL17BR* \geq -1.339).

Univariate analyses of clinical outcomes. Four risk groups were formed by combining CYP2D6 metabolism and *HOXB13/IL17BR* (Table 2). DFS and OS were found to differ significantly with respect to these four patient categories ($P = 0.004$ and 0.009 , respectively). Table 2 presents the Kaplan-Meier estimates of 5-year DFS rate and its corresponding confidence interval (CI) for each risk group. To evaluate the cumulative effect of carrying one or more risk factors, we analyzed DFS and OS based on the absence of a risk factor (extensive CYP2D6 and *HOXB13/IL17BR* low), only one high-risk feature (either CYP2D6 metabolism decreased or *HOXB13/IL17BR* high), and finally two high-risk features (both CYP2D6 decreased and *HOXB13/IL17BR* high). In this analysis, both DFS ($P = 0.001$) and OS ($P = 0.005$) differed significantly according to risk groups (Figs. 1 and 2).

Multivariate analyses of clinical outcomes. For each end point, proportional hazard modeling was done using age, extent of surgery, ER status, tumor size, tumor grade, and HER2 expression. Once tumor size was accounted for, none of the other traditional factors made a significant contribution to explaining the variability in these clinical outcomes. Further

multivariate proportional hazard modeling showed that after adjusting for tumor size, DFS and OS were significantly different among patients with zero, one, or two risk factors as determined by CYP2D6 and *HOXB13/IL17BR* gene ratio. Relative to women with no risk factors (extensive CYP2D6 and *HOXB13/IL17BR* low), women with at least one risk factor had significantly worse DFS [adjusted hazard ratio (HR), 2.03; 95% CI, 1.04-3.94; $P = 0.037$] and OS (adjusted HR, 2.41; 95% CI, 1.08-5.37; $P = 0.031$). An even greater effect was observed in women with two risk factors (decreased CYP2D6 metabolism and *HOXB13/IL17BR* high). These women had the shortest DFS (adjusted HR, 3.10; 95% CI, 1.34-7.17; $P = 0.008$) and OS (adjusted HR, 3.15; 95% CI, 1.17-8.52; $P = 0.024$; Table 3).

Discussion

Our analysis of this cohort of ER-positive, lymph node-negative patients treated with adjuvant tamoxifen strongly suggests that a combination of inherited (*CYP2D6*) and tumor (*HOXB13/IL17BR*) genetic variation influences the risk of breast cancer recurrence and death, independent of standard prognostic factors. Furthermore, we showed a stepwise increased risk of recurrence and death, with the greatest risk in those patients who carry both high-risk features (e.g., both decreased CYP2D6 metabolism and a high *HOXB13/IL17BR* gene ratio). In contrast to prior studies in lymph node-negative breast cancer that have simply focused on tumor genetic variation (20), the paradigm of accounting for host genetic variation affecting drug metabolism, as well as tumor somatic variation, offers the potential for a more accurate estimate of drug benefit and the identification of groups of patients for whom alternative or additional therapies could be considered.

The potential synergy between *HOXB13/IL17BR* and *CYP2D6* as biomarkers for tamoxifen resistance may be related to the differential effects of tamoxifen and its metabolites in the setting of tumor cells with dysfunctional estrogen signaling. Endoxifen exhibits a significantly greater affinity for the ER than tamoxifen (3) and multiple studies suggest that only two tamoxifen metabolites, endoxifen and 4-OH tamoxifen, have significant antiproliferative activity (3, 22). Tamoxifen and other metabolites exhibit only weak antagonist/agonist properties (23). The importance of these pharmacologic differences may be magnified in women with a high *HOXB13/IL17BR* gene ratio, as *in vitro*, *HOXB13* and *IL17BR* are regulated by estradiol, which suppresses *HOXB13* and augments *IL17BR*

Table 3. Results of multivariate modeling of DFS and OS

Factor	Clinical outcome	
	DFS	OS
Tumor size \geq 3 cm	2.64 (1.42-4.92)	2.52 (1.23-5.14)
Either <i>HOXB13/IL17BR</i> high or decreased CYP2D6 metabolism	2.03 (1.04-3.94)	2.41 (1.08-5.37)
Both <i>HOXB13/IL17BR</i> high and decreased CYP2D6 metabolism	3.10 (1.34-7.17)	3.15 (1.17-8.52)

NOTE: Multivariate modeling based on tumor size, and the three risk group models comparing presence of one high-risk feature (defined as the presence of either decreased CYP2D6 metabolism or a *HOXB13/IL17BR* gene ratio \geq -1.339) or two high-risk features (defined as the presence of both decreased CYP2D6 metabolism and a *HOXB13/IL17BR* gene ratio \geq -1.339) relative to CYP2D6 extensive metabolizers with a *HOXB13/IL17BR* <-1.339 HRs and corresponding 95% CIs.

expression (13). In tumors with a high *HOXB13/IL17BR* ratio, the loss of both suppression of *HOXB13* expression and induction of *IL17BR* expression may not only be a marker for impaired ER signaling but also for increased growth factor signaling. Based on studies showing that ER-positive tumors with a high *HOXB13/IL17BR* ratio are more likely to over-express HER2 (13), *in vivo* ER binding of tamoxifen and other nonpotent metabolites (as opposed to endoxifen) may activate growth factor signaling and thus tumor growth (18). This hypothesis is supported by preliminary data wherein non-transformed human mammary epithelial cells expressing *HOXB13* showed a synergistic increase in cell motility and invasion in the presence of epidermal growth factor (12).

Most of the published breast cancer gene signatures seem to provide similar information in the setting of ER-positive breast cancer (24) and proliferation-related genes seem to be an essential component accounting for the concordance between these expression signatures (12, 20, 25–28). Although “proliferation” signatures seem to identify those patients who derive the greatest benefit from adjuvant chemotherapy (29), the role of *HOXB13* and *IL17BR* as a tool for clinical decision making has been unclear. However, our data suggest that postmenopausal patients with decreased CYP2D6 metabolism (especially CYP2D6 poor metabolizers) may preferentially benefit from

alternative hormonal therapy (e.g., aromatase inhibitor) in the setting of tumors with a high *HOXB13/IL17BR* ratio. Further retrospective studies of large adjuvant trials comparing tamoxifen with aromatase inhibitors are necessary to validate this hypothesis.

In summary, our data suggest that an index composed of CYP2D6 and *HOXB13/IL17BR* may provide a robust indicator of tamoxifen resistance. A prospective approach incorporating both factors may lead to the improved individualization of endocrine therapy.

Disclosure of Potential Conflicts of Interest

X-J. Ma, M.G. Erlander, and D.C. Sgroi are named inventors on a patent to use the *HOXB13/IL17BR* expression ratio to ascertain breast cancer prognosis. X-J. Ma and M.G. Erlander are employees of AviraDx. M.P. Goetz, J.N. Ingle, M.M. Ames, and F.J. Couch are named inventors on a pending patent application about the combined index of CYP2D6 and *HOXB13/IL17BR* as a predictor for tamoxifen resistance. M.P. Goetz and J.N. Ingle have been consultants for Roche (2007). M.P. Goetz, J.N. Ingle, and F.J. Couch received research funding from Arcturus in 2005.

Acknowledgments

We thank the women who participated in this clinical trial as well as the NCCCTG investigators and clinical research associates at each site who made this translational research study possible.

References

- Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomized trials. *Lancet* 2005; 365:1687–717.
- Howell A, Cuzick J, Baum M, et al. Results of the ATAC (Arimidex, Tamoxifen, Alone or in Combination) trial after completion of 5 years' adjuvant treatment for breast cancer. *Lancet* 2005;365:60–2.
- Johnson MD, Zuo H, Lee KH, et al. Pharmacological characterization of 4-hydroxy-*N*-desmethyl tamoxifen, a novel active metabolite of tamoxifen. *Breast Cancer Res Treat* 2004;85:151–9.
- Lim YC, Desta Z, Flockhart DA, Skaar TC. Endoxifen (4-hydroxy-*N*-desmethyl-tamoxifen) has anti-estrogenic effects in breast cancer cells with potency similar to 4-hydroxy-tamoxifen. *Cancer Chemother Pharmacol* 2005;55:471–8.
- Gjerde J, Hauglid M, Breilid H, et al. Effects of CYP2D6 and SUL1A1 genotypes including SUL1A1 gene copy number on tamoxifen metabolism. *Ann Oncol* 2008;19:56–61.
- Jin Y, Desta Z, Stearns V, et al. CYP2D6 genotype, antidepressant use, and tamoxifen metabolism during adjuvant breast cancer treatment. *J Natl Cancer Inst* 2005;97:30–9.
- Bonanni B, Macis D, Maisonneuve P, et al. Polymorphism in the CYP2D6 tamoxifen-metabolizing gene influences clinical effect but not hot flashes: data from the Italian Tamoxifen Trial. *J Clin Oncol* 2006;24:3708–9, author reply 9.
- Lim HS, Ju Lee H, Seok Lee K, Sook Lee E, Jang IJ, Ro J. Clinical implications of CYP2D6 genotypes predictive of tamoxifen pharmacokinetics in metastatic breast cancer. *J Clin Oncol* 2007;25:3837–45.
- Goetz MP, Knox SK, Suman VJ, et al. The impact of cytochrome P450 2D6 metabolism in women receiving adjuvant tamoxifen. *Breast Cancer Res Treat* 2007; 101:113–21.
- Goetz MP, Rae JM, Suman VJ, et al. Pharmacogenetics of tamoxifen biotransformation is associated with clinical outcomes of efficacy and hot flashes. *J Clin Oncol* 2005;23:9312–8.
- Schroth W, Antoniadou L, Fritz P, et al. Breast cancer treatment outcome with adjuvant tamoxifen relative to patient CYP2D6 and CYP2C19 genotypes. *J Clin Oncol* 2007;25:5187–93.
- Ma XJ, Wang Z, Ryan PD, et al. A two-gene expression ratio predicts clinical outcome in breast cancer patients treated with tamoxifen. *Cancer Cell* 2004;5: 607–16.
- Wang Z, Dahiya S, Provencher H, et al. The prognostic biomarkers *HOXB13*, *IL17BR*, and *CHDH* are regulated by estrogen in breast cancer. *Clin Cancer Res* 2007;13:6327–34.
- Miao J, Wang Z, Provencher H, et al. *HOXB13* promotes ovarian cancer progression. *Proc Natl Acad Sci U S A* 2007;104:17093–8.
- Goetz MP, Suman VJ, Ingle JN, et al. A two-gene expression ratio of homeobox 13 and interleukin-17B receptor for prediction of recurrence and survival in women receiving adjuvant tamoxifen. *Clin Cancer Res* 2006;12:2080–7.
- Jansen MP, Sieuwerts AM, Look MP, et al. *HOXB13*-to-*IL17BR* expression ratio is related with tumor aggressiveness and response to tamoxifen of recurrent breast cancer: a retrospective study. *J Clin Oncol* 2007;25:662–8.
- Jerevall PL, Brommesson S, Strand C, et al. Exploring the two-gene ratio in breast cancer-independent roles for *HOXB13* and *IL17BR* in prediction of clinical outcome. *Breast Cancer Res Treat* 2008;107:225–34.
- Ma XJ, Hilsenbeck SG, Wang W, et al. The *HOXB13:IL17BR* expression index is a prognostic factor in early-stage breast cancer. *J Clin Oncol* 2006;24: 4611–9.
- Ingle JN, Suman VJ, Mailliard JA, et al. Randomized trial of tamoxifen alone or combined with flouxymesterone as adjuvant therapy in postmenopausal women with resected estrogen receptor positive breast cancer. North Central Cancer Treatment Group Trial 89-30-52. *Breast Cancer Res Treat* 2006;98:217–22.
- Paik S, Shak S, Tang G, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 2004;351: 2817–26.
- Goetz MP, Reid J, Safgren S, et al. UGT1A1*28 genotype determines the maximally tolerated dose and pharmacokinetics of irinotecan-based chemotherapy: a phase I dose-escalation trial. *ASCO Gastrointestinal Cancers Symposium* 2007.
- Buck MB, Collier JK, Murdter TE, Eichelbaum M, Knabbe C. TGFβ2 and TβRII are valid molecular biomarkers for the antiproliferative effects of tamoxifen and tamoxifen metabolites in breast cancer cells. *Breast Cancer Res Treat* 2008;107:15–24.
- Johnson MD, Westley BR, May FE. Oestrogen activity of tamoxifen and its metabolites on gene regulation and cell proliferation in MCF-7 breast cancer cells. *Br J Cancer* 1989;59:727–38.
- Fan C, Oh DS, Wessels L, et al. Concordance among gene-expression-based predictors for breast cancer. *N Engl J Med* 2006;355:560–9.
- Loi S, Haibe-Kains B, Desmedt C, et al. Definition of clinically distinct molecular subtypes in estrogen receptor-positive breast carcinomas through genomic grade. *J Clin Oncol* 2007;25:1239–46.
- Oh DS, Troester MA, Usary J, et al. Estrogen-regulated genes predict survival in hormone receptor-positive breast cancers. *J Clin Oncol* 2006;24: 1656–64.
- van de Vijver MJ, He YD, van't Veer LJ, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* 2002;347:1999–2009.
- Wang Y, Klijn JG, Zhang Y, et al. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* 2005;365: 671–9.
- Paik S, Tang G, Shak S, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol* 2006;24:3726–34.

Clinical Cancer Research

Cytochrome P450 2D6 and Homeobox 13/Interleukin-17B Receptor: Combining Inherited and Tumor Gene Markers for Prediction of Tamoxifen Resistance

Matthew P. Goetz, Vera J. Suman, Fergus J. Couch, et al.

Clin Cancer Res 2008;14:5864-5868.

Updated version	Access the most recent version of this article at: http://clincancerres.aacrjournals.org/content/14/18/5864
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2021/03/12/14.18.5864.DC1

Cited articles	This article cites 28 articles, 12 of which you can access for free at: http://clincancerres.aacrjournals.org/content/14/18/5864.full#ref-list-1
Citing articles	This article has been cited by 4 HighWire-hosted articles. Access the articles at: http://clincancerres.aacrjournals.org/content/14/18/5864.full#related-urls

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://clincancerres.aacrjournals.org/content/14/18/5864 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.