

Cancer Therapy: Clinical

Phase I Study of Recombinant Human Interleukin-7
Administration in Subjects with Refractory Malignancy

Claude Sportès¹, Rebecca R. Babb¹, Michael C. Krumlauf¹, Frances T. Hakim¹, Seth M. Steinberg², Catherine K. Chow⁵, Margaret R. Brown⁶, Thomas A. Fleisher⁶, Pierre Noel⁶, Irina Maric⁶, Maryalice Stetler-Stevenson³, Julie Engel⁷, Renaud Buffet⁷, Michel Morre⁷, Robert J. Amato⁸, Andrew Pecora⁹, Crystal L. Mackall⁴, and Ronald E. Gress¹

Abstract

Purpose: Interleukin-7 (IL-7) has critical and nonredundant roles in T-cell development, hematopoiesis, and postdevelopmental immune functions as a prototypic homeostatic cytokine. Based on a large body of preclinical evidence, it may have multiple therapeutic applications in immunodeficiency states, either physiologic (immunosenescence), pathologic (HIV), or iatrogenic (postchemotherapy and posthematopoietic stem cell transplant), and may have roles in immune reconstitution or enhancement of immunotherapy. We report here on the toxicity and biological activity of recombinant human IL-7 (rhIL-7) in humans.

Design: Subjects with incurable malignancy received rhIL-7 subcutaneously every other day for 2 weeks in a phase I interpatient dose escalation study (3, 10, 30, and 60 µg/kg/dose). The objectives were safety and dose-limiting toxicity determination, identification of a range of biologically active doses, and characterization of biological and, possibly, antitumor effects.

Results: Mild to moderate constitutional symptoms, reversible spleen and lymph node enlargement, and marked increase in peripheral CD3⁺, CD4⁺, and CD8⁺ lymphocytes were seen in a dose-dependent and age-independent manner in all subjects receiving ≥10 µg/kg/dose, resulting in a rejuvenated circulating T-cell profile, resembling that seen earlier in life. In some subjects, rhIL-7 induced in the bone marrow a marked, transient polyclonal proliferation of pre-B cells showing a spectrum of maturation as well as an increase in circulating transitional B cells.

Conclusion: This study shows the potent biological activity of rhIL-7 in humans over a well-tolerated dose range and allows further exploration of its possible therapeutic applications. *Clin Cancer Res*; 16(2):727–35. ©2010 AACR.

Interleukin-7 (IL-7) has a wide range of biological activities including a critical role in T-cell development as well as roles in hematopoiesis and postdevelopmental immune functions (1–4). Unlike in the mouse, IL-7 is not required for human B-cell development. IL-7 plays a critical and nonredundant role in maintaining peripheral T-cell homeostasis and is considered a prototypic homeostatic cy-

tokine. It is required to maintain naive CD4⁺ and CD8⁺ cells *in vivo* (5–7) and IL-7 levels are inversely correlated to the T-cell depletion status (7, 8).

A multitude of murine and simian preclinical models have shown the wide array of immune-enhancing properties of IL-7, which may find clinical applications. Recombinant human IL-7 (rhIL-7) improves immune reconstitution in nonhuman primate models. In SIV-infected monkeys, it increases the number of circulating naive and memory CD4⁺ and CD8⁺ T cells (9, 10). IL-7 enhances immune reconstitution, improving CD4⁺ reconstitution following autologous CD34⁺ selected stem cell transplantation in monkeys (11) as well as in mice (12–18). It also enhances anti-infectious cytotoxicity in several animal models following adoptive transfer of specific T cells (19) or bone marrow transplant (20).

IL-7 enhances antitumor immune responses in mice (13, 21, 22). It is critical in the proliferation and *in vivo* persistence of tumor-specific T cells (23) and enhances antitumor responses and survival (24). IL-7 administration post-transplant results in both improved T-cell recovery and prolonged tumor-free survival (12). Immune

Authors' Affiliations: ¹Experimental Transplantation and Immunology Branch, ²Biostatistics and Data Management Section, ³Laboratory of Pathology, and ⁴Pediatric Oncology Branch, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, Maryland; Departments of ⁵Radiology and ⁶Laboratory Medicine, Clinical Center, NIH; ⁷Cytheris, Inc., Rockville, Maryland; ⁸Methodist Hospital, Texas Medical Center, Houston, Texas; and ⁹The Cancer Center at Hackensack University Medical Center, Hackensack, New Jersey

Note: C.L. Mackall and R.E. Gress contributed equally to this work.

Corresponding Author: Claude Sportès, Experimental Transplantation and Immunology Branch, Center for Cancer Research, National Cancer Institute, NIH, 10 Center Drive, Building 10-CRC Room 43142, Bethesda, MD 20892-1203. Phone: 301-435-52-80; Fax: 301-402-75-15; E-mail: csportes@mail.nih.gov.

doi: 10.1158/1078-0432.CCR-09-1303

©2010 American Association for Cancer Research.

Translational Relevance

Interleukin-7 (IL-7) has critical and nonredundant roles in T-cell development, hematopoiesis, and post-developmental immune functions. Based on a large body of preclinical evidence, it is a prototypic homeostatic cytokine that may have multiple therapeutic applications in a wide array of immunodeficiency states, either physiologic (immunosenescence), pathologic (e.g., HIV), or iatrogenic (e.g., postchemotherapy and posthematopoietic stem cell transplant), and, in particular, may have roles in immune reconstitution or enhancement of immunotherapy in multimodality cancer therapy strategies. Indeed, IL-7 has been ranked number 5 on the National Cancer Institute list of "Top 20 Agents with High Potential for Use in Treating Cancer" at the July 12, 2007 National Cancer Institute Immunotherapy Agent Workshop (<http://web.ncicrf.gov/research/brb/workshops/NCI%20Immunotherapy%20Workshop%207-12-07.pdf>). This article shows that recombinant human IL-7 has potent biological activity over a dose range that is well tolerated in humans and therefore will allow further exploration of its possible therapeutic applications.

enhancement after severe immune depletion such as following stem cell transplantation may have clinical value because early lymphocyte reconstitution correlates with clinical outcome following autologous transplantation (25–28). Furthermore, standard or high-dose radiochemotherapy often only induces temporary complete or partial disease responses but no cure, leaving candidates for novel immunotherapy strategies immunologically depleted, which may hinder the effectiveness of such strategies.

The potential therapeutic uses of rhIL-7 are centered on its immunomodulatory effects on T cells and could include generating or optimizing T-cell responses to novel antitumor or anti-infectious immune interventions in a variety of clinical circumstances with severe immune depletion, be they physiologic (e.g., aging), pathologic (e.g., HIV), or iatrogenic (immunodepleting therapies). We conducted a phase I study of the administration of rhIL-7 to determine dose-limiting toxicity (DLT), maximum tolerated dose, and a range of biologically active doses in humans.

Materials and Methods

Eligibility. Adults diagnosed with incurable nonhematologic malignancy with a life expectancy greater than 3 months were eligible. A peripheral CD3⁺ cells count above 300 cells/mm³ was required. Also required were normal left ventricular ejection fraction, DLCO/VA and FEV1 >50%, AST and ALT less than three times the upper limit of normal, absolute neutrophil count >1,000/mm³, platelet count >100,000/mm³, PT/PTT within 1.5 times the up-

per limit of normal, creatinine clearance of >60 mL/min, and Karnofsky performance status >70%. Exclusions were recent use or need for anticoagulation, systemic corticosteroids, and cytotoxic or immunosuppressive therapy; splenectomy, splenomegaly, and prior solid organ or allogeneic stem cell transplantation; infections with HIV and hepatitis B or C; and history of autoimmune disease and severe uncontrolled asthma. Toxicity was evaluated according to the National Cancer Institute "Common Terminology Criteria for Adverse Events." The study was approved by the National Cancer Institute Institutional Review Board and the Food and Drug Administration. Subjects signed written informed consent before enrollment.

Trial design. "CYT 99 007" (*Escherichia coli* rhIL-7; Cytheris) was given subcutaneously every other day for 2 weeks in successive cohorts of three to six patients in a standard phase I design (dose levels: 3, 10, 30, and 60 µg/kg/dose). All subjects, except the first on study, received the same product lot. The primary study objectives were safety, DLT, and maximum tolerated dose determination. Secondary objectives included defining a range of biologically active doses, defining rhIL-7 pharmacokinetics and evaluating possible antitumor effects. Biological activity was defined as 50% increase over baseline in the number of peripheral blood CD3⁺ cells/mm³.

Subjects underwent body computed tomography scan at baseline, end of therapy (day 14), and day 28. Subjects with stable disease at day 28 were followed if possible until disease progression. Bone marrow aspirate and biopsy were done at baseline and day 14.

Flow cytometry. Peripheral blood mononuclear cell analysis: We analyzed fresh, EDTA-anticoagulated peripheral blood at the specified time points as described previously (29).

Bone marrow flow cytometry. Bone marrow aspirates were collected into sterile sodium heparin, prelysed with ammonium chloride, and stained for 30 min at room temperature with antibodies directed against CD13, CD33, and CD36 (Beckman Coulter); CD3, CD10, CD14, CD16, CD19, CD22, CD34, CD56, CD45, and CD71 (BD Biosciences); CD64 (Caltag Laboratories); and CD20, κ, λ (DakoCytomation). Cells were fixed in 1.0% paraformaldehyde and stored at 4°C for up to 12 h before acquisition. Four-color cytometry was done with a Becton Dickinson FACSCalibur. Data were analyzed with CellQuest software (Becton Dickinson). Granulocytes, monocytes, mature lymphocytes, nucleated RBC, and immature hematopoietic precursors were determined based on levels of CD45 expression, side scatter, and pattern of antigen expression.

Immunoglobulin gene rearrangement. DNA was extracted from Ficoll-Hypaque separated bone marrow mononuclear cells and PCR-amplified for detection of immunoglobulin heavy-chain gene rearrangements using primers to framework region III and the joining region of the immunoglobulin heavy-chain gene (FRIII-IGH PCR). Additional PCR was done in duplicate using consensus primers to framework region II and the immunoglobulin heavy-chain

Table 1. Summary of subject characteristics, toxicity, and immunogenicity

| UPN | Cohort no./ ($\mu\text{g}/\text{kg}/\text{dose}$) | Age | Diagnosis | Toxicity (possibly, probably, or definitely related), maximum toxicity grade and type (grade) | Anti-IL7 antibody titer | |
|-----|--|-----|---------------------------------|---|-------------------------|--------------|
| | | | | | Binding | Neutralizing |
| 01 | 1/(3) | 70 | Renal cell carcinoma* | 1 IS (1); EO (1) | Positive (1/1,600) | 0 |
| 02 | | 48 | Hemangiopericytoma | 2 ALT (2) | Negative (0) | NA |
| 03 | | 67 | Melanoma | | Negative (0) | NA |
| 04 | 2/(10) | 56 | Melanoma | 2 IS (2); CS (1) | Negative (1/100) | 0 |
| 05 | | 71 | Renal cell carcinoma | 2 IS (2); CS (1); ALT (1) | Negative (0) | NA |
| 06 | | 28 | Carcinoma of unknown primary | 2 IS (2); CS (2) | Negative (0) | NA |
| 07 | 3/(30) | 21 | Osteogenic sarcoma [†] | 3 AST, bili (3); CS (1) | NA | NA |
| 08 | | 20 | Rhabdomyosarcoma | 1 IS (1); CS (1) | Positive (1/400) | 0 |
| 09 | | 28 | Synovial cell sarcoma | 1 IS (1); CS (1); SM (1) | Negative (1/100) | 0 |
| 10 | | 48 | Duodenal carcinoma | 2 IS (2); CS (1) | Negative (1/100) | 0 |
| 11 | | 61 | Colon carcinoma | | Positive (1/1,200) | 0 |
| 12 | | 62 | Melanoma | 2 IS (2); CS, SOB (2); ALT, bili (1) | Positive (1/400) | 0 |
| 13 | 4/(60) | 19 | Melanoma | 2 IS (2); CS, bili (2); ALT, AP, Plt (1) | Positive (1/1,200) | 0 |
| 14 | | 35 | Pheochromocytoma [‡] | 3 Cardiac [§] (3); AST, ALT (1) | Negative (1/100) | NA |
| 15 | | 43 | Ovarian carcinoma | 3 ALT (3); AST, IS (2); CS, AP (1) | Positive (1/1,200) | 0 |
| 16 | | 66 | Melanoma | 1 IS, CS, bili (1) | Negative (0) | NA |

Abbreviations: ALT and AST, transaminases; AP, alkaline phosphatase; bili, total bilirubin; CS, constitutional symptoms; EO, eosinophilia; IS, injection site; erythema, induration, pruritus; Plt, decreased platelets; SM, splenomegaly; SOB, shortness of breath.

*Received initial lot of "CYT 99 007."

[†]Received only one dose.

[‡]Received only three doses.

[§]See text.

joining region (FRII-IGH PCR) according to the method of Ramasamy et al. (30).

Serum anti-IL-7 antibody determination. Antibody detection used a two-step ELISA and a bioassay for neutralization of IL-7 bioactivity, both developed for and used during rhIL-7 preclinical development by Cytheris. Pretreatment and day 28 serum samples were assayed. Assays were repeated at days 35 and 42 if equivocal at day 28. Titer $\geq 1/200$ was considered positive for the ELISA, but a neutralization bioassay was done for any titer $\geq 1/100$. A neutralizing antibody titer $\geq 1/400$ at either day 28 or 42 was to be considered positive and a DLT.

Serum rhIL-7 level determination and pharmacokinetics. Serum IL-7 levels were determined serially for pharmacokinetics following the first and last doses. The Diaclone Research (Cell Systems) Sandwich IL-7 ELISA kit was used for validation and sample analysis. The two-site sandwich ELISA used anti-rhIL-7 monoclonal antibodies, one as coating antibody for capture and the other, biotinylated, for detection. The linear relationship between rhIL-7 concentrations and absorbance extended up to 200 pg/mL; concentrations < 12.5 pg/mL were treated as zero for pharmacokinetic calculations (Kinetic Software version 4.2) with the noncompartmental extravascular model. The area under the curves (AUC) were computed using the log lin-

ear method, trapezoidal when $C_n > C_{n-1}$. Half-lives were calculated between $t = 2$ h (T_{\max}) and $T = 24$ h.

Statistics. Using the data presented in Table 2, differences between pretreatment and post-treatment values and the relative difference (post-pre)/pre were formed for the four markers of interest of mature B cells ($\text{CD}19^+/\text{CD}45^{\text{bright}}$ and $\text{CD}19^+/\text{CD}10^-$) and immature B cells ($\text{CD}19^+/\text{CD}45^{\text{dim}}$ and $\text{CD}19^+/\text{CD}10^+$). In the one case in which the value was noted to be < 1.0 , this was changed to be 0.5 for simplicity. It was determined that the association between either the absolute or the relative difference was approximately equally well associated with the pretreatment value; thus, for simplicity, the absolute difference was evaluated to determine if the change was equal to zero. A Wilcoxon signed rank test was used to determine if the change was equal to zero after determining that the differences were not normally distributed. All P values are two-sided and presented without formal adjustment for multiple comparisons.

Results

Twelve men and 4 women, ages 19 to 71 years (median, 49 years), with incurable malignancies, were treated. Table 1 summarizes subject characteristics, toxicity, and immunogenicity data.

Toxicity. "CYT 99 007" was well tolerated. Following most injections, subjects receiving ≥ 10 $\mu\text{g}/\text{kg}$ developed grade 1 to 2 constitutional symptoms 6 to 8 h following administration and mild local reactions with erythema, pruritus, and induration. Two subjects did not complete therapy because of toxicity. One developed rapidly reversible grade 3 liver enzyme and bilirubin elevation following the first dose at the 30 $\mu\text{g}/\text{kg}$ and treatment was terminated. The other developed grade 3 chest pain and hypertension with mild troponin elevation several hours after the third dose of 60 $\mu\text{g}/\text{kg}$; treatment was stopped and all abnormalities abated within 2 days (see Discussion). Subsequently, no significant electrocardiogram changes were seen when systematically performed at 3 and 24 h following every injection on all remaining subjects. No symptoms or signs of autoimmunity were seen.

Biological activity. Starting at 10 $\mu\text{g}/\text{kg}/\text{dose}$, all subjects showed evidence of biological activity (a trend toward activity was seen at 3 $\mu\text{g}/\text{kg}/\text{dose}$). A 50% to 60% drop in absolute lymphocyte counts following the first dose was seen in all subjects on day 1 as in most preclinical models. All patients showed a dose-dependent marked increase in absolute lymphocyte count, which persisted for several weeks after treatment (Fig. 1A). Although several hematologic parameters, serum albumin, total protein, and C-reactive protein showed mild fluctuations during the

study, none, except for absolute lymphocyte count, appeared clinically significant; transient platelet decrease during treatment; monocyte decrease after treatment; and eosinophils peak on day 21, all (except the C-reactive protein values) remaining within the normal institutional range. Lymphoid organ enlargement (spleen and lymph nodes but not thymus) was seen in a dose-dependent manner on computed tomography scan, maximum on day 14, returning to baseline over several weeks (Fig. 1B).

T-cell populations. For all T-cell populations, the absolute cell numbers peaked on day 21 and remained elevated for weeks, whereas lymphopenia following treatment was never observed. In most subjects, CD3^+ $\alpha\beta$ and $\gamma\delta$ T cells increased equally and the increase was similar in CD4^+ and CD8^+ cells as well as CD3^+ (but not CD3^-) CD16^+ or CD56^+ cells. An in-depth analysis of T-cell subset kinetics and their consequences on the circulating T-cell makeup and T-cell repertoire diversity following rhIL-7 therapy has been reported (29) and is summarized in the discussion. Briefly, up to 60% of circulating CD3^+ cells were in cycle (Ki-67^+) on day 7 of treatment. In spite of treatment continuation, the fraction of circulating cells in cycle had decreased by the end of treatment and returned to baseline 1 week after the end of treatment. Both IL-7R α surface expression and mRNA were decreased during the treatment and returned

Table 2. Lymphoid bone marrow expansion with rhIL-7 therapy

| UPN (age, y) | Mature B cells | | | | Immature B cells | | | |
|--------------|---|-------|--------------------------------------|-------|--|--------|--------------------------------------|--------|
| | CD19 ⁺ /CD45 ^{bright} | | CD19 ⁺ /CD10 ⁻ | | CD19 ⁺ /CD45 ^{dim} | | CD19 ⁺ /CD10 ⁺ | |
| | Pre | Post | Pre | Post | Pre | Post | Pre | Post |
| 01 (70) | 0.4 | 1 | 0.4 | 0 | 0.2 | 8 | 0.2 | 0 |
| 02 (48) | 4 | 2 | 2 | 1 | 0.4 | 10 | 1 | 11 |
| 03 (67) | 5 | 3 | 4 | 3 | 1 | <1 | 2 | 1 |
| 04 (56) | 6.9 | 2 | 5.5 | 1 | 7 | 59 | 10 | 55 |
| 05 (71) | 3.5 | 27 | 2 | 1.9 | 11 | 23 | 12 | 47 |
| 06 (28) | 4.2 | 2 | 4.3 | 15 | 1.2 | 18 | 0.8 | 5 |
| 08 (20) | 26 | 35 | 17 | 6 | 17 | 51 | 22 | 80 |
| 09 (28) | 11 | 4 | 9.4 | 4 | 12 | 8 | 13 | 8 |
| 10 (48) | 7.3 | 23 | 6.6 | 4 | 6.5 | 21 | 7 | 39 |
| 11 (61) | 10 | 2 | 10 | 2 | 6 | 6 | 4 | 10 |
| 12 (62) | 10 | 0.8 | 7 | 1 | 2 | 7 | 2 | 6 |
| 13 (19) | 6.9 | 1.5 | 6.3 | 0.7 | 4.6 | 33 | 4.6 | 30 |
| 15 (43) | 14.8 | 1 | 10.7 | 1 | 19.2 | 80 | 11.9 | 80 |
| 16 (66) | 3 | 2 | 3 | 4 | 3 | 11 | 3 | 9 |
| Mean | 8.07 | 7.59 | 6.30 | 3.19 | 6.51 | 23.96 | 6.68 | 27.21 |
| Median | 6.90 | 2.0 | 5.90 | 1.95 | 5.30 | 14.50 | 4.30 | 10.50 |
| Post-Pre | | -0.48 | | -3.11 | | 17.46 | | 20.54 |
| <i>P</i> | | 0.38 | | 0.024 | | 0.0012 | | 0.0029 |

NOTE: Summary of bone marrow flow cytometry data. Percentage of bone marrow mononuclear cells for the marker combinations: mature B cells: $\text{CD19}^+/\text{CD45}^{\text{bright}}$ and $\text{CD19}^+/\text{CD10}^-$; immature B cells: $\text{CD19}^+/\text{CD45}^{\text{dim}}$ and $\text{CD19}^+/\text{CD10}^+$. UPN: unique patient number. *P* value given (Wilcoxon signed rank test) on the absolute difference between "pre" and "post" values for each marker combination.

to baseline after the end of treatment. Interestingly, the subject who received only the first three doses of rhIL-7 showed the same cell cycling and IL-7R α downregulation patterns (29) at day 7 and had a day 28 increase in absolute lymphocyte count similar to that of the other subjects, suggesting that maximum therapeutic effects may be reached early possibly because of IL-7R α downregulation (see Discussion).

IL-2R α (CD25) surface expression was studied in a selected group of patients receiving the higher doses of rhIL-7. There was evidence of increased CD25 surface expression in the CD4 $^+$ non-Tregs and CD8 $^+$ T-cell populations. The mean percent increase over baseline of CD25 $^+$ cells were 55% (95% confidence interval, 40-70) at day 7 and 57% (95% confidence interval, 45-69) at day 14 for CD4 $^+$ cells and 74% (95% confidence interval, 63-85) at day 7 and 85% (95% confidence interval, 72-98) at day 14 for CD8 $^+$ cells with return to baseline at day 21.

Circulating B cells. Mature B cells (CD20 $^+$) consistently showed a modest and not demonstrably dose-dependent decrease in absolute number during therapy and returned to baseline by day 28 (Fig. 1C). Several subjects showed a mild expansion of circulating CD19 $^+$ /CD10 $^+$ B cells [immature/transitional B cells (31, 32)], peaking on day 21 (Fig. 1C) with no correlation between cell number and subjects' age or magnitude of their bone marrow pre-B-cell proliferation (see below). There was no increase in circu-

lating activated germinal center B cells (CD19 $^+$ /CD27 $^+$ /CD10 $^+$).

Bone marrow. All subjects underwent bone marrow aspiration and biopsy evaluation before and at the end of therapy by detailed analysis of H&E, immunostains, and flow cytometry. Although 17 different markers or marker combinations spanning the spectrum of bone marrow lineages and of differentiation stages were analyzed by flow cytometry, most showed variations between pretreatment and post-treatment samples that remained within the normally occurring wide normal range. In general, rhIL-7 treatment resulted, by day 14, in an increased cellularity, usually within normal limits. There were no significant abnormalities in morphology or abundance of the myeloid, erythroid, or megakaryocytic lineages. The proportion of CD4 $^+$ and CD8 $^+$ T cells remained largely unchanged and most subjects showed a statistically significant increase in CD19 $^+$ /CD10 $^+$ and/or CD19 $^+$ /CD45 $^{\text{dim}}$ early pre-B cells (Table 2). In 6 subjects, however, the CD10 and Tdt abnormalities on biopsy immunostains would have triggered further flow cytometry investigation in other clinical settings (Fig. 2), and in these cases, post-treatment early pre-B cells represented 30% to 86% of all bone marrow mononuclear cells by flow cytometry, showing no commensurate increase in mature B cells (Table 2) and no correlation with treatment dose or subjects' age (Fig. 3A). As shown in Fig. 3B, CD19 $^+$ B-cell




Fig. 1. Clinical parameter kinetics with rhIL-7 therapy. Cohorts: 1, black; 2, green; 3, red; 4, blue. A, top left, absolute lymphocyte count cohort mean \pm SE. Others: mean \pm SE for all subjects. B, left, mean spleen size; spleen on computed tomography: baseline, day 14, 2 months. C, left, mature B cells, mean absolute numbers; right, immature B cells, mean percentage of CD19 $^+$ population.




Fig. 2. H&E, immunostaining (CD10 and Tdt; $\times 200$), and Giemsa stain ($\times 400$), before and after rhIL-7 therapy, in subject with marked marrow expansion (UPN08).

immunostaining in the flow cytometry analysis was consistent with ongoing maturation: increasing levels of CD45 and CD19 expression (polygonal gate in Fig. 3A); in gated CD19⁺ B cells: increasing CD20 with decreasing CD10 expression (Fig. 3B) and increasing CD45 with decreasing CD34 expression (Fig. 3C). PCR studies of both bone marrow and peripheral blood mononuclear cells for immunoglobulin gene rearrangement failed to reveal any clonal population (data not shown). Follow-up bone marrow or peripheral blood studies, performed several weeks after the end of treatment in subjects with the highest pre-B-cell increases, consistently showed either declining or resolved pre-B-cell marrow proliferation by flow cytometry, confirmed to have remained polyclonal by PCR.

Immunogenicity. Sera were tested for anti-IL-7 antibodies on day 28 and, if required, on day 42 (Table 1). The only subject treated with the first lot of rhIL-7 (01) developed a 1/1,600 antibody titer. Excluding this subject who received a different formulation, 5 of the 14 subjects evaluable for immunogenicity developed anti-IL-7 antibodies. None of the 5 receiving 3 or 10 $\mu\text{g}/\text{kg}$ and 5 of 9 receiving 30 or 60 $\mu\text{g}/\text{kg}$ developed antibodies on day 28, suggesting a dose response. No subject developed anti-IL-7 neutralizing antibodies. None of the subjects with longer follow-up developed late lymphopenia.

Antitumor activity. Although the study was not designed to evaluate antitumor activity, disease follow-up was carried out as feasible. Subject 02 with central nervous system hemangiopericytoma and abdominal metastases (no central nervous system disease at entry on study) had a substantial improvement of abdominal pain and a 20% reduction in mass size at 3 months. The abdominal disease remained stable at the time of a central nervous system recurrence, 9 months following rhIL-7 therapy. All other patients had progressive disease by day 42.

Pharmacokinetics. Pharmacokinetics studies (performed after the first and last rhIL-7 doses) are provided in Table 3. The rhIL-7 half-life ranged from 6.5 to 9.8 h. Whereas the C_{max} increased from the 10 to 30 $\mu\text{g}/\text{kg}$ dose, the day 1 total AUC increased throughout the dose range, and clearance decreased, whereas apparent half-life increased with the dose. High-affinity clearance mechanisms, mediated by IL-7R, probably explain a better clearance at low dose. The consistently higher clearance and lower AUC at day 14 compared with day 1 at a given dose probably reflect the larger lymphoid mass at day 14. rhIL-7 pharmacodynamics seem to correlate better with AUC and half-life than with C_{max} , consistent with the characteristic pharmacodynamic/pharmacokinetic profile of growth factors and cytokines: prolonged AUCs and half-lives tend to improve efficacy by sustaining blood levels above activity threshold over a prolong period, whereas peak concentrations well above activity threshold (high C_{max}) contribute less to the biological effects.

Discussion

We present here clinical findings of a phase I study of rhIL-7, defining a toxicity profile and a range of biologically active doses. Enrollment was limited to individuals with a stable CD3⁺ peripheral count >300 cells/ mm^3 . Studies in other populations (HIV and allogeneic transplant recipients) are under way. Although the first dose level was designed as a "no effect dose" based on nonhuman primate studies (9), early signs of biological activity were apparent at this dose, suggesting a 10- to 30-fold greater sensitivity in humans.

rhIL-7 was well tolerated with no maximum tolerated dose being reached in the treated subjects. The observation of only mild adverse reactions along with the biological activity in the tested dose range suggests a safe therapeutic window. One subject had liver DLT, concomitant with the onset of a catheter infection. The cardiac DLT occurred in the subject with pheochromocytoma, with typical symptoms for this disease (causing the DLT) occurring concurrently with the expected constitutional symptoms. These two potentially rhIL-7-related toxicities will require further scrutiny in subsequent trials. At the two highest doses, spleen and lymph node enlargements were noticeable but not clinically worrisome, although caution during the period of rapid spleen enlargement remains warranted. rhIL-7 immunogenicity was not limiting and no neutralizing antibodies or lymphopenia developed. A glycosylated

rhIL-7 is presently under clinical development, which may prove less immunogenic and allow the evaluation of repeated courses of treatment. There was no suggestion of the development of autoimmunity.

In a limited number of subjects, there was a marked, self-limited expansion of polyclonal bone marrow pre-B cells (hematogones), consistent with *in vitro* observations that human pro-B cells display a functional IL-7R and expand following IL-7 exposure but subsequently die without reaching mature B-cell stage (33–35). Several subjects showed a mild expansion of circulating CD19⁺/CD10⁺ transitional B cells, a population found in HIV patients to correlate with serum IL-7 levels (31). There was no obvious correlation between bone marrow and peripheral blood expansions of immature B cells and the number of circulating mature B cells was largely unaffected by rhIL-7 treatment. IL-7 effects on B-cell development and physiology accumulated in the murine system are mostly irrelevant to humans and the knowledge that rhIL-7 may induce a self-limited, but often marked, polyclonal pre-B-cell expansion in humans is valuable safety information for the future rhIL-7 clinical trials.

rhIL-7 induced a dose-dependent, but age-independent, increase in circulating CD3⁺ (both $\alpha\beta$ and $\gamma\delta$) lymphocytes, CD3⁺ (but not CD3⁻) CD16⁺ or CD56⁺ cells, and CD3⁺/CD4⁻/CD8⁻ T cells (29). This net increase in circulating lymphocytes was most likely due to proliferation/

expansion taking place in the peripheral blood and/or lymphoid organs with varying contribution by alteration in cell trafficking. The massive increase in the number of cycling circulating lymphocytes is unlikely due solely to trafficking out of lymphoid organs, considering the concomitant spleen and lymph node enlargement as well as the continuing increase in circulating lymphocyte numbers after therapy. Other evidence supported T-cell expansion as resulting from a combination of increased cell cycling and decreased apoptosis (29). The sharp decrease in cell cycling while pharmacologic serum IL-7 levels were sustained during the second week of treatment suggests that downregulation of IL-7R α plays a role in the self-limited effects of rIL-7, a consideration for the design of future trials.

Detailed studies of T-cell subsets, reported elsewhere (29), showed that rhIL-7 preferentially expanded the CD4⁺ naive and central memory T cells as well as CD8⁺ naive T cells resulting in a statistically significant broadening of T-cell repertoire diversity in both CD4⁺ and CD8⁺ T cells. Therefore, within 3 weeks of treatment initiation, rhIL-7 resulted in a marked expansion of T cells, which remained functional (with conserved or increased *in vitro* responsiveness to anti-CD3 stimulation) and exhibited a rejuvenated profile resembling that seen early in life (29). Additionally, CD4⁺/FoxP3⁺ regulatory T cells showed markedly less proliferation, consistent with their low IL-7R α -chain expression (36, 37), expanded less than




Fig. 3. Bone marrow B cells. CD19⁺ cells and subsets: % total bone marrow mononuclear cells (subjects ages <30 years in red). Dxx; 4 to 10 weeks; correlation between bone marrow CD19⁺/CD10⁺ pre-B cells and peripheral blood increases in CD20⁺ (●) and CD19⁺/D10⁺ (○). Flow cytometry scatter plots of marrow early B-cell precursors in subject with marked expansion (UPN08) illustrating progressive maturation. A, bone marrow cells (ungated data), X axis CD45 PerCP and Y axis CD19 APC and CD19⁺ B cells (in polygonal analysis gate), have increasing levels of CD45 and CD19 expression consistent with ongoing maturation. B, CD19⁺ B cells (gate-polygonal gate in A), X axis CD20 FITC and Y axis CD10 PE and CD19⁺ B cells, have increasing levels of CD20 expression and decreasing levels of CD10 expression consistent with ongoing maturation. C, CD19⁺ B cells (gate-polygonal gate in A), X axis CD45 PerCP and Y axis CD34 PE and CD19⁺ B cells, have increasing levels of CD45 expression and decreasing levels of CD34 expression consistent with ongoing maturation.

Table 3. rhIL-7 pharmacokinetics

| Dose | 10 µg/kg | | 30 µg/kg | | 60 µg/kg | |
|-----------------------------|----------|--------|-----------|---------|-----------|-----------|
| rhIL-7 serum levels (pg/mL) | | | | | | |
| Time (h) | d 1 | d 14 | d1 | d 14 | d 1 | d 14 |
| 0 | 6 | 0 | 0 | 10 | 0 | 0 |
| 0.5 | 51 | 35 | 506 | 372 | 534 | 929 |
| 1 | 93 | 48 | 837 | 399 | 908 | 1,019 |
| 1.5 | 121 | 57 | 1,335 | 616 | 1,279 | 1,324 |
| 2 | 148 | 66 | 1,895 | 688 | 1,246 | 1,354 |
| 2.5 | 147 | 69 | 1,766 | 426 | 1,223 | 1,344 |
| 4 | 354 | 83 | 1,531 | 717 | 1,176 | 1,211 |
| 8 | 120 | 53 | 917 | 459 | 821 | 177 |
| 12 | 87 | 58 | 936 | 438 | 832 | 161 |
| 24 | 28 | 6 | 296 | 122 | 478 | 139 |
| Pharmacokinetics parameters | | | | | | |
| Mean total daily dose (µg) | 686 | 686 | 2,322 | 2,322 | 4,763 | 5,280 |
| C _{max} (pg/mL) | 354 | 83 | 1,895 | 717 | 1,846 | 1,697 |
| T _{max} (h) | 4.0 | 4.0 | 2.0 | 4.0 | 4.0 | 4.0 |
| AUC total (pg/mL min) | 168,357 | 69,697 | 1,416,470 | 591,776 | 1,812,870 | 1,483,180 |
| t _{1/2} (h) | 8.54 | 9.80 | 9.03 | 6.46 | 8.92 | 7.23 |
| Clearance (mL/min) | 4,077 | 9,847 | 1,639 | 3,890 | 2,627 | 3,560 |

NOTE: Top: mean serum concentrations for cohorts. Bottom: pharmacokinetics parameters (mean values for a cohort). Results are shown after the first dose (day 1) and after the last dose (day 14) for each cohort.

other CD4⁺ T cells resulting in a decreased proportion of circulating regulatory T cells (29, 38).

In summary, rhIL-7 is well tolerated in humans and has significant biological activity over dose range, indicating a potential safe therapeutic window. It induces an age-independent marked expansion of the T-cell mass resulting in a rejuvenated T-cell profile with an increased T-cell repertoire diversity and a decreased proportion of regulatory T cells. This study paves the way for the systematic investigation of specific clinical applications for enhancement of immune responses or immune reconstitution in a multitude of clinical circumstances of immune depletion, be they iatrogenic (chemotherapy/radiation/transplantation), pathologic (HIV), or physiologic (immunosenescence), and may be considered to optimize strategies of passive or active immunotherapy.

Disclosure of Potential Conflicts of Interest

This research was supported by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research and was made possible through a formal collaboration (Cooperative Research and Development Agreement grant 01649) between the National Cancer Institute and Cytheris, the investigational new drug holder and manufacturer of rhIL-7, currently developing rhIL-7 for therapeutic use. As per the Cooperative Research and Development Agreement between National Cancer Institute and Cytheris, the National Cancer Institute investiga-

tors designed and conducted all the research experiments and analyses presented in the article independently from Cytheris. Three coauthors (J. Engel, R. Buffet, and M. Moore) have financial interest in Cytheris capital: M. Moore is the founder and CEO and J. Engel and R. Buffet were employees of Cytheris. All other coauthors have explicitly denied any conflict of interest.

Acknowledgments

We thank the subjects who enrolled on this investigational trial and provided consent for research studies and the clinical staff of the NIH Clinical Center and the Experimental Transplantation and Immunology Branch for excellent care of these subjects.

Grant Support

National Cancer Institute, NIH, Department of Health and Human Services in collaboration with Cytheris under a Cooperative Research and Development Agreement grant 01649.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 5/22/09; revised 10/8/09; accepted 10/25/09; published OnlineFirst 1/12/10.

References

- Komschlies KL, Grzegorzewski KJ, Wiltrout RH. Diverse immunological and hematological effects of interleukin 7: implications for clinical application. *J Leukoc Biol* 1995;58:623–33.
- Hofmeister R, Khaled AR, Benbernou N, et al. Interleukin-7: physiological roles and mechanisms of action. *Cytokine Growth Factor Rev* 1999;10:41–60.
- Appasamy PM. Biological and clinical implications of interleukin-7 and lymphopoiesis. *Cytokines Cell Mol Ther* 1999;5:25–39.
- Fry TJ, Mackall CL. Interleukin-7: from bench to clinic. *Blood* 2002;99:3892–904.
- Schluns KS, Kieper WC, Jameson SC, Lefrancois L. Interleukin-7 mediates the homeostasis of naive and memory CD8 T cells *in vivo*. *Nat Immunol* 2000;1:426–32.
- Fry TJ, Connick E, Falloon J, et al. A potential role for interleukin-7 in T-cell homeostasis. *Blood* 2001;97:2983–90.
- Napolitano LA, Grant RM, Deeks SG, et al. Increased production of IL-7 accompanies HIV-1-mediated T-cell depletion: implications for T-cell homeostasis. *Nat Med* 2001;7:73–9.
- Bolotin E, Annett G, Parkman R, Weinberg K. Serum levels of IL-7 in bone marrow transplant recipients: relationship to clinical characteristics and lymphocyte count. *Bone Marrow Transplant* 1999;23:783–8.
- Fry TJ, Moniuszko M, Creekmore S, et al. IL-7 therapy dramatically alters peripheral T-cell homeostasis in normal and SIV-infected non-human primates. *Blood* 2003;101:2294–9.
- Beq S, Nugeyre MT, Ho Tsong FR, et al. IL-7 induces immunological improvement in SIV-infected rhesus macaques under antiviral therapy. *J Immunol* 2006;176:914–22.
- Storek J, Gillespy T III, Lu H, et al. Interleukin-7 improves CD4 T-cell reconstitution after autologous CD34 cell transplantation in monkeys. *Blood* 2003;101:4209–18.
- Talmadge JE, Jackson JD, Kelsey L, et al. T-cell reconstitution by molecular, phenotypic, and functional analysis in the thymus, bone marrow, spleen, and blood following split-dose polychemotherapy and therapeutic activity for metastatic breast cancer in mice. *J Immunother* 1993;14:258–68.
- Komschlies KL, Gregorio TA, Gruys ME, et al. Administration of recombinant human IL-7 to mice alters the composition of B-lineage cells and T cell subsets, enhances T cell function, and induces regression of established metastases. *J Immunol* 1994;152:5776–84.
- Bolotin E, Smogorzewska M, Smith S, Widmer M, Weinberg K. Enhancement of thymopoiesis after bone marrow transplant by *in vivo* interleukin-7. *Blood* 1996;88:1887–94.
- Mackall CL, Fry TJ, Bare C, et al. IL-7 increases both thymic-dependent and thymic-independent T-cell regeneration after bone marrow transplantation. *Blood* 2001;97:1491–7.
- Alpdogan O, Schmaltz C, Muriglan SJ, et al. Administration of interleukin-7 after allogeneic bone marrow transplantation improves immune reconstitution without aggravating graft-versus-host disease. *Blood* 2001;98:2256–65.
- Broers AE, Posthumus-van Sluijs SJ, Spits H, et al. Interleukin-7 improves T-cell recovery after experimental T-cell-depleted bone marrow transplantation in T-cell-deficient mice by strong expansion of recent thymic emigrants. *Blood* 2003;102:1534–40.
- Alpdogan O, Muriglan SJ, Eng JM, et al. IL-7 enhances peripheral T cell reconstitution after allogeneic hematopoietic stem cell transplantation. *J Clin Invest* 2003;112:1095–107.
- Wiryan P, Bui T, Faltynek CR, Ho RJ. Augmentation of cell-mediated immunotherapy against herpes simplex virus by interleukins: comparison of *in vivo* effects of IL-2 and IL-7 on adoptively transferred T cells. *Vaccine* 1997;15:561–3.
- Abdul-Hai A, Ben Yehuda A, Weiss L, et al. Interleukin-7-enhanced cytotoxic T lymphocyte activity after viral infection in marrow transplanted mice. *Bone Marrow Transplant* 1997;19:539–43.
- Komschlies KL, Back TT, Gregorio TA, et al. Effects of rhIL-7 on leukocyte subsets in mice: implications for antitumor activity. *Immunol Ser* 1994;61:95–104.
- Murphy WJ, Longo DL. The potential role of NK cells in the separation of graft-versus-tumor effects from graft-versus-host disease after allogeneic bone marrow transplantation. *Immunol Rev* 1997;157:167–76.
- Wang LX, Li R, Yang G, et al. Interleukin-7-dependent expansion and persistence of melanoma-specific T cells in lymphodepleted mice lead to tumor regression and editing. *Cancer Res* 2005;65:10569–77.
- Li B, Vanroey MJ, Jooss K. Recombinant IL-7 enhances the potency of GM-CSF-secreting tumor cell immunotherapy. *Clin Immunol* 2007;123:155–65.
- Porrata LF, Gertz MA, Inwards DJ, et al. Early lymphocyte recovery predicts superior survival after autologous hematopoietic stem cell transplantation in multiple myeloma or non-Hodgkin lymphoma. *Blood* 2001;98:579–85.
- Porrata LF, Litzow MR, Tefferi A, et al. Early lymphocyte recovery is a predictive factor for prolonged survival after autologous hematopoietic stem cell transplantation for acute myelogenous leukemia. *Leukemia* 2002;16:1311–8.
- Porrata LF, Inwards DJ, Micallef IN, et al. Early lymphocyte recovery post-autologous haematopoietic stem cell transplantation is associated with better survival in Hodgkin's disease. *Br J Haematol* 2002;117:629–33.
- Porrata LF, Ingle JN, Litzow MR, Geyer S, Markovic SN. Prolonged survival associated with early lymphocyte recovery after autologous hematopoietic stem cell transplantation for patients with metastatic breast cancer. *Bone Marrow Transplant* 2001;28:865–71.
- Sportes C, Hakim FT, Memon SA, et al. Administration of rhIL-7 in humans increases *in vivo* TCR repertoire diversity by preferential expansion of naive T cell subsets. *J Exp Med* 2008;205:1701–14.
- Ramasamy I, Brisco M, Morley A. Improved PCR method for detecting monoclonal immunoglobulin heavy chain rearrangement in B cell neoplasms. *J Clin Pathol* 1992;45:770–5.
- Malaspina A, Moir S, Ho J, et al. Appearance of immature/transitional B cells in HIV-infected individuals with advanced disease: correlation with increased IL-7. *Proc Natl Acad Sci U S A* 2006;103:2262–7.
- Sims GP, Ettinger R, Shirota Y, et al. Identification and characterization of circulating human transitional B cells. *Blood* 2005;105:4390–8.
- Dittel BN, LeBien TW. The growth response to IL-7 during normal human B cell ontogeny is restricted to B-lineage cells expressing CD34. *J Immunol* 1995;154:58–67.
- LeBien TW. Fates of human B-cell precursors. *Blood* 2000;96:9–23.
- Johnson SE, Shah N, Panoskaltzis-Mortari A, LeBien TW. Murine and human IL-7 activate STAT5 and induce proliferation of normal human pro-B cells. *J Immunol* 2005;175:7325–31.
- Liu W, Putnam AL, Xu-Yu Z, et al. CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4⁺ Treg cells. *J Exp Med* 2006;203:1701–11.
- Seddiki N, Santner-Nanan B, Martinson J, et al. Expression of interleukin (IL)-2 and IL-7 receptors discriminates between human regulatory and activated T cells. *J Exp Med* 2006;203:1693–700.
- Rosenberg SA, Sportes C, Ahmadzadeh M, et al. IL-7 administration to humans leads to expansion of CD8⁺ and CD4⁺ cells but a relative decrease of CD4⁺ T-regulatory cells. *J Immunother* 2006;29:313–9.

Clinical Cancer Research

Phase I Study of Recombinant Human Interleukin-7 Administration in Subjects with Refractory Malignancy

Claude Sportès, Rebecca R. Babb, Michael C. Krumlauf, et al.

Clin Cancer Res 2010;16:727-735. Published OnlineFirst January 12, 2010.

Updated version Access the most recent version of this article at:
[doi:10.1158/1078-0432.CCR-09-1303](https://doi.org/10.1158/1078-0432.CCR-09-1303)

Cited articles This article cites 38 articles, 21 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/16/2/727.full#ref-list-1>

Citing articles This article has been cited by 26 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/16/2/727.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/16/2/727>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.