

Cancer Therapy: Clinical

Estrogen Receptor- α Phosphorylation at Serine 305, Nuclear p21-Activated Kinase 1 Expression, and Response to Tamoxifen in Postmenopausal Breast CancerJosefine Bostner¹, Lambert Skoog², Tommy Fornander³, Bo Nordenskjöld¹, and Olle Stål¹

Abstract

Purpose: *In vitro*, p21-activated kinase 1 (Pak1) phosphorylates the serine 305 residue of the estrogen receptor α (ER α) and influences the response of breast cancer cells to tamoxifen. We investigated the influence of Pak1 and pER α^{ser305} on breast cancer prognosis and results of tamoxifen therapy.

Experimental Design: We examined Pak1 and pER α^{ser305} protein by immunohistochemistry in a series of 912 tumors from node-negative breast cancer patients randomized to tamoxifen or no adjuvant endocrine treatment.

Results: Cytoplasmic Pak1 correlated to large tumors and ER negativity, whereas nuclear Pak1 and pER α^{ser305} correlated to small tumors and ER positivity. Nuclear expression of Pak1 and pER α^{ser305} predicted reduced response to tamoxifen in patients with ER α -positive tumors (tamoxifen versus no tamoxifen: hazard ratio (HR), 1.33; 95% confidence interval (95% CI), 0.42-4.2; $P = 0.63$), whereas patients lacking this combination benefitted significantly from tamoxifen (HR, 0.43; 95% CI, 0.30-0.62; $P < 0.0001$). Similar nonsignificant trends were detected in analyses of the proteins separately. Pak1 in the cytoplasm was an independent prognostic marker, indicating increased recurrence rate (HR, 1.79; 95% CI, 1.17-2.74; $P = 0.0068$) and breast cancer mortality (HR, 1.98; 95% CI, 1.14-3.46; $P = 0.016$) for patients randomized to no adjuvant treatment.

Conclusion: Our results suggest that patients with tumors expressing Pak1 and pER α^{ser305} in combination are a group in which tamoxifen treatment is insufficient. In addition, the pathway may be of interest as a drug target in breast cancer. Furthermore, the findings support previous studies showing that Pak1 has differential roles in the cytoplasm and the nucleus. *Clin Cancer Res*; 16(5); 1624-33. ©2010 AACR.

Breast cancer is the most common type of cancer worldwide in the female population (1). Mammography screening and improved surgery, radiotherapy, and adjuvant systemic treatments correlate with decreased mortality during the last decades.

Tamoxifen is an estrogen competitor, commonly used as adjuvant therapy in estrogen receptor (ER)-positive breast cancer. Nonetheless, many tumors relapse due to *de novo* resistance, or over time, *acquired* resistance (2). Tamoxifen response seems to be affected by ER α modifications, including phosphorylation of the ER α and its coregulators as well as other alterations affecting coregulator

dynamics (3). Phosphorylation of ER α serine, threonine, and tyrosine residues is a posttranslational event changing the secondary structure of the highly flexible receptor and is caused by growth factor-regulated kinases. This influences the receptor in several aspects, such as subcellular localization, dimerization, DNA binding, coregulator interaction, and transcriptional activity (4). There are several known phosphorylation sites in ER α , in which serines 118, 167, and 305, controlled by mitogen-activated protein kinase, Akt, PKA, and Pak1 signaling pathways, seem to be the most interesting in relation to endocrine therapy (5, 6). These sites are involved in estrogen-independent transcriptional activation of some ER target genes, such as *CCND1* (7, 8).

Tharakan and colleagues (9) recently showed that an ER α^{S305E} mutant, mimicking a constitutively phosphorylated state, increased receptor dimerization in the presence of estrogen compared with wild-type ER α . The mutant receptor also increased ER α target gene binding compared with wild-type in the absence of ligand. p21-activated kinase (Pak1) activates the ER α by phosphorylation of the serine residue at position 305, leading to the ER α transcriptional activation of ER target genes (7). Additionally, Pak1 phosphorylates the ER α coactivator NRIF3, leading to a potentiation of the ER α transactivation (10). Various

Authors' Affiliations: ¹Division of Surgery and Clinical Oncology, Department of Clinical and Experimental Medicine, Faculty of Health Sciences, Linköping University, Linköping, Sweden; and Departments of ²Cytology and ³Oncology, Karolinska University Hospital, Stockholm, Sweden

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Corresponding Author: Josefine Bostner, Division of Surgery and Clinical Oncology, Department of Clinical and Experimental Medicine, SE-581 85 Linköping, Sweden. Phone: 46-13-222008; Fax: 46-13-221718; E-mail: josefine.bostner@liu.se.

doi: 10.1158/1078-0432.CCR-09-1733

©2010 American Association for Cancer Research.

Translational Relevance

Adjuvant endocrine treatment with tamoxifen or aromatase inhibitor significantly improves the survival of women with estrogen receptor (ER)-positive breast cancer, but resistance to treatment is a huge clinical problem. Modifications of the ER by phosphorylation of several serine residues, leading to increased transcriptional activation, are part of the cross-talk between ER and growth factor receptors that when perturbed may cause resistance. We investigated the expression of pER^{ser305} and p21-activated kinase 1 (Pak1), which may phosphorylate ER at serine 305, in a large tumor material originating from a randomized tamoxifen trial with postmenopausal patients. Patients with tumors positive for pER^{ser305} and nuclear Pak1 showed decreased benefit from tamoxifen, suggesting that these proteins might become biomarkers for tamoxifen resistance. In addition, cytoplasmic Pak1 was found to be an independent prognostic factor among systemically nontreated patients. Those dual roles of Pak1 may make it an interesting target in cancer therapy.

combinations of phospho modifications of the serines 118, 236, and 305 led to different ER α conformational changes upon antiestrogen treatment, seen by the FRET structural analysis (11). Within the cytoplasm of breast epithelial cells, active Pak1 interferes with numerous cellular events, of which several influence tumor progression (12, 13).

This study was based on the pER^{ser305} and Pak1 immunohistochemical analysis of a large series of tumors from node-negative postmenopausal breast cancer patients, who participated in a randomized tamoxifen trial. We report that phosphorylated ER α ^{ser305}, along with the detection of Pak1 protein in the nucleus, correlate with reduced response to tamoxifen in ER α -positive breast cancer. In addition, Pak1 localization influences prognosis and tamoxifen treatment prediction in this set of patients.

Materials and Methods

Patients and tumors. During the years of 1976 through 1990, a cohort of Swedish postmenopausal breast cancer patients was included in a controlled trial to evaluate tamoxifen as adjuvant treatment (14). Patients with low-risk tumors, defined as node negative and ≤ 30 mm in diameter, were included in the present study. Patients were treated either with modified radical mastectomy or with breast-conserving therapy and radiation therapy to the breast with a total dose of 50 Gy with 2 Gy per fraction, 5 d weekly, for ~ 5 wk. All patients were randomized to either 2 y of tamoxifen (40 mg daily) or no adjuvant en-

docrine therapy. In 1983, a new trial was initiated: recurrence-free patients, after 2 y, were again randomized to either tamoxifen for 3 more y or no further therapy. The standard procedure for tissue collection was fixation at room temperature in 4% phosphate-buffered formalin. ER status was determined by isoelectric focusing and the cutoff level was set to 0.05 fmol/ μ g DNA (14). Follow-up data were collected through regional population registers and the Swedish Cause of Death Registry. The median follow-up period for recurrence-free patients in the present study was 17.8 y. The present study was approved by the Stockholm regional Ethics board.

Tissue microarray. Archived tumor tissues were collected from 912 of the 1,780 patients included in the original study. Representative tumor tissues were selected as donor blocks for the tissue micro array. Sections were cut from each donor block and stained with H&E. From these slides, three morphologically representative regions were chosen in each sample. Three cylindrical core tissue specimens with a diameter of 0.8 mm were taken from these areas in each case and mounted in a recipient block. Sixteen tissue array blocks were constructed using a manual arayer (Beecher, Inc.). Tissue microarray blocks were cut with a microtome into 4- μ m sections and mounted on frost-coated glass slides. The subset in the present study did not significantly differ from the original study including 1,780 patients, with respect to a tumor size of 20 mm or less (79% versus 81%), positive ER status (78% versus 80%), or tamoxifen treatment (52% versus 50%; Supplementary Table).

Hormone receptor status. Retrospectively, additional ER and progesterone receptor (PR) status of the tumors was evaluated with immunohistochemistry using the Ventana automated slide stainer (Ventana Medical Systems). The antibodies used were the monoclonal Ventana Medical Systems' CONFIRM mouse anti-ER primary antibody (clone 6F11) and the monoclonal Ventana Medical Systems' CONFIRM mouse anti-PR primary antibody (clone 16). Cutoff level was set to 10% positively stained tumor cell nuclei. In this study, these data were used for ER and PR status. However, when immunohistochemical data on ER status was missing, results from the cytosol assay were used. In this way, ER status could be defined for 886 of the 912 tumors.

Immunohistochemistry. Formalin-fixed and paraffin-embedded tumor tissue microarray slides were deparaffinized with xylen and rehydrated in a graded ethanol series. The slides later stained with Pak1 antibody were boiled in target retrieval solution (pH 9.9; DAKO) in microwave oven for 4 \times 5 min, at 750 W up to boiling point, and thereafter switching between 160 and 350 W to control boiling. The slides later stained with pER^{ser305} antibody were boiled in citrate buffer (pH 6.0) in a pressure cooker at 125°C for 30 s and withdrawn when the temperature reached 95°C to expose antigenic epitopes. The sections were cooled in room temperature for 20 min, placed in 3% H₂O₂ in methanol for 5 min to inactivate endogenous peroxidase, incubated with serum-free protein block

Table 1. Correlations between Pak1 cytoplasmic expression, Pak1 nuclear expression, pER^{ser305}, and clinicopathologic parameters in postmenopausal breast carcinomas

	Pak1 nuclear expression			Pak1 cytoplasmic expression			pER ^{ser305} nuclear expression		
	n (%)		P	n (%)		P	n (%)		P
	-	+		-	+		-	+	
Total	615 (78)	171 (22)		333 (42)	453 (58)		536 (64)	305 (36)	
Tamoxifen			0.92			0.15			0.87
-	294 (78)	81 (22)		149 (40)	226 (60)		257 (63)	148 (37)	
+	321 (78)	90 (22)		184 (45)	227 (55)		279 (64)	157 (36)	
Tumor size			<0.001			<0.001			<0.001
≤20mm	451 (75)	149 (25)		277 (46)	323 (54)		389 (61)	253 (39)	
>20mm	149 (90)	17 (10)		46 (28)	120 (72)		134 (75)	45 (25)	
ER status			0.01			0.009			0.008
-	144 (86)	24 (14)		57 (34)	111 (66)		131 (72)	51 (28)	
+	458 (77)	138 (23)		270 (45)	326 (55)		390 (61)	246 (39)	
PR status			0.02			0.59			0.07
-	279 (83)	56 (17)		139 (41)	196 (59)		243 (68)	113 (32)	
+	281 (76)	87 (24)		160 (43)	208 (57)		240 (62)	148 (38)	
pER ^{ser305}			0.46			0.001			
-	384 (80)	99 (20)		178 (37)	305 (63)				
+	203 (77)	60 (23)		129 (49)	134 (51)				

(DAKO) for 10 min, and incubated with rabbit polyclonal Pak1 primary antibody (Cell Signaling Technology) diluted at 1:25 at 4°C in a moisturized chamber for 20 h or rabbit polyclonal pER^{ser305} primary antibody (Bethyl Laboratories) diluted at 1:300 at 4°C in a moisturized chamber for 17 h. All slides were washed, incubated with the anti-rabbit DAKO cytomation Envision+ system labeled with horse radish peroxidase antibody (DAKO) for 30 min at 4°C, visualized with 3,3'-diaminobenzidine hydrochloride in phosphate buffer and 0.03% H₂O₂, and counterstained with hematoxylin. All washing steps were done in a phosphate buffer solution with 0.5% bovine serum albumin. Staining intensity was evaluated on three separate biopsies for each tumor. Pak1 nuclear reactivity was graded as positive when >10% of the tumor cells showed staining. Pak1 cytoplasmic protein expression was graded as negative, weak, moderate, and strong, in which moderate and strong staining was considered as abnormal expression of the protein in subsequent statistical analysis. Fourteen percent of the patients ($n = 126$) were excluded in the Pak1 assay due to nonrepresentative or missing tissues. pER^{ser305} nuclear staining was graded with respect to percent positively stained nuclei in four steps: negative, 1% to 24%, 25% to 74%, and ≥75%. As the proportion of positive staining was moderate, cutoff point for positive staining was set to the ≥1% level in the statistical analysis. pER^{ser305} was frequently visible in the cytoplasm, but only nuclear staining was graded. Eight percent of the patients ($n = 71$) in the evaluation of pER^{ser305} were excluded due to nonrepresentative or missing tissues. Only minor differences were detected be-

tween subsets evaluated for protein staining and excluded samples compared with the original study cohort (Supplementary Table). For each antibody, all tumors were stained in one batch and the whole range of staining intensities was represented on each slide, due to the large number of tumors. In addition, normal liver samples on all slides served as control. All tumor samples were evaluated by two independent observers, blinded to pathologic and clinical data, and scored under a Leica DM LS (Leica Microsystems) light microscope. Where discordance was found, cases were reevaluated to reach consensus. Pictures were generated using an Olympus SC20 camera with a Leica ×40 objective.

Antibody validation. Validation of the pER^{ser305} antibody included blocking with immunizing peptide according to manufacturers instruction (Bethyl Laboratories) and dephosphorylation of proteins in acetone-fixed T47D breast cancer cells by λ-phosphatase (λ-ppase; New England Biolabs). In the λ-ppase assay, slides were treated with 1,000 units of λ-ppase or water (control) for 2 h at 37°C followed by immunohistochemical staining according to the protocol used for the pER^{ser305} primary antibody. One concern in the detection of phospho proteins regards tissue collection routines. Validation of several ER phospho antibodies did not show a significant decreased detection of phospho ERs in relation to increased time of collection (15). The specificity of the Pak1 antibody was previously confirmed by Western blot and immunohistochemistry (16).

Statistical analysis. To compare protein expression data with prognostic and clinical characteristics, the Pearson χ^2




Fig. 1. Immunohistochemical detection of Pak1 and pER α^{ser305} in human breast cancer sections and validation of ER α phospho antibody with blocking peptide in breast tumor tissue and with λ -ppase-treated T47D breast cancer cells. Magnification, $\times 40$. Pak1-negative cytoplasmic staining and negative nuclei (A), Pak1-positive cytoplasm and positive nuclei (B), Pak1-positive cytoplasm and negative nuclei (C), pER α^{ser305} -negative nuclear staining (D), pER α^{ser305} -positive staining in $>1\%$ to 25% of nuclei (E), pER α^{ser305} -positive staining in $>75\%$ of nuclei (F), pER α^{ser305} without blocking peptide (G), pER α^{ser305} with blocking peptide (H), pER α^{ser305} without λ -ppase (I), and pER α^{ser305} with λ -ppase (J).

test was applied. Hazard ratios (HR) and 95% confidence intervals (95% CI) were estimated using the Cox proportional hazards model. Recurrence-free survival time was calculated as the time between diagnosis and any of the events: locoregional recurrence, distant metastasis, or breast cancer death. Recurrence-free survival time distributions were compared with the log-rank test and plots were drawn using the Kaplan-Meier technique. Multivariate analysis of recurrence rates and breast cancer mortality rates were done with Cox proportional hazard regression, a method also used for the interaction analysis of different factors and treatment by including the variables: potential predictive factor, tamoxifen treatment, and interaction variable (tamoxifen \times potential predictive factor). Tumor size was also included in the model. Anal-

ysis of prognosis was restricted to patients randomized to no tamoxifen treatment, and treatment prediction of tamoxifen was restricted to patients with ER α -positive tumors. P values of ≤ 0.05 were considered significant, with the exception of the associations presented in Table 1, in which P values of ≤ 0.01 were considered significant to compensate for the effect of multiple comparisons. All statistical data were analyzed using the Statistica 8.0 software program.

Results

Expression analysis of Pak1 and phosphorylated ER α^{ser305} proteins with immunohistochemistry was successful with 786 and 841 tumors, respectively. Pak1

cytoplasmic overexpression was observed in 57.6%; Pak1 nuclear expression was observed in 21.8%; and pER α^{ser305} nuclear expression was detected in 36.3% of the tumors (Fig. 1A-F). As Pak1 previously has been shown to

increase the transcriptional activity of ER α through phosphorylation of its amino acid residue, serine 305, we investigated whether the expression of the two proteins in the nuclear compartment was correlated, but they




Fig. 2. Tamoxifen response among ER-positive patients in relation to nuclear protein expression of Pak1-negative (A), Pak1-positive (B), pER α^{ser305} -negative (C), pER α^{ser305} -positive (D), Pak1- and/or pER α^{ser305} -negative (E), and Pak1- and pER α^{ser305} -positive cells (F).

Table 2. Cox proportional hazard models for Pak1 nuclear expression, pER^{ser305} nuclear expression, Pak1 and pER^{ser305} in combination, and the benefit from tamoxifen in patients with ER-positive tumors

	Recurrence tamoxifen vs no tamoxifen		
	HR (95% CI)	P	P _{interaction} *
Pak1 nuclear expression			
-	0.43 (0.29-0.64)	<0.001	0.32
+	0.62 (0.32-1.20)	0.15	
pER ^{ser305} nuclear expression			
-	0.48 (0.32-0.72)	<0.001	0.24
+	0.66 (0.39-1.13)	0.13	
Pak1 and pER ^{ser305} nuclear expression			
-	0.43 (0.30-0.62)	<0.001	0.029
+	1.33 (0.42-4.19)	0.63	

*For details, see Material and Methods.

were not when all tumors were considered (Table 1). However, restricting the analysis to Pak1-positive tumors, independent of localization, rendered a significant correlation between Pak1 nuclear expression and pER^{ser305} nuclear expression ($P = 0.042$). Pak1 cytoplasmic staining correlated significantly to a tumor size larger than 20 mm and a negative ER status, whereas nuclear staining of Pak1 as well as pER^{ser305} correlated to small size and ER positivity. The expression of pER^{ser305} correlated with ER status. Twenty-eight percent (51 of 182) of the tumors defined as ER negative showed some staining for pER^{ser305}. However, the cutoff level for ER status was set to 10% positive nuclei, whereas cutoff for pER^{ser305} was set to >1% positive nuclei. The correlation was stronger when the cutoff level for pER^{ser305} positivity was raised and also when analyzing the pER^{ser305} staining in four categories with increasing percentage of positive cells ($P = 0.0026$). Tumors defined as ER positive were also positive for pER^{ser305} in 39% of the tumors (246 of 636).

Antibody specificity. The pER^{ser305} antibody specificity was determined using the immunizing peptide used for antibody production, including 12 to 15 amino acids surrounding the phospho site and a phosphorylated serine at the position corresponding to serine 305 of ER α . Incubation of the antibody with the peptide before immunoassay resulted in the loss of staining (Fig. 1G-H). For further evaluation of the phospho specificity of the antibody, T47D breast cancer cells on glass slides were treated with λ -ppase, resulting in the staining of control cells and no visible staining of λ -ppase-treated cells (Fig. 1I-J). Previously, the pER^{ser305} antibody used in this study was validated through lack of detection of the mutated pER^{ser305A} compared with the wild-type pER α on Western blot (17). Taken together, data indicate that the pER α -

^{ser305} antibody specifically recognizes ER α phosphorylated at serine 305.

Tamoxifen treatment prediction. Recurrence-free survival in tamoxifen-treated versus nontamoxifen-treated groups in relation to protein expression levels was analyzed to estimate the benefit from adjuvant tamoxifen treatment in subsets of ER α -positive breast cancer patients. Our data show that patients with no nuclear expression of




Fig. 3. Recurrence-free survival for all patients given no adjuvant systemic treatment in relation to protein expression detected by immunohistochemistry. Pak1 cytoplasmic expression (A), Pak1 nuclear expression (B), and pER^{ser305} nuclear expression (C).

Table 3. Multivariate analysis of patients randomized to no adjuvant tamoxifen treatment using Cox proportional hazard regression

	Recurrence		Breast cancer death	
	HR (95% CI)	P	HR (95% CI)	P
Tumor size				
>20 mm vs ≤20 mm	1.88 (1.24-2.85)	0.0029	2.57 (1.57-4.22)	0.0002
ERα status				
Positive vs negative	0.64 (0.38-1.11)	0.11	0.68 (0.36-1.30)	0.25
PR status				
Positive vs negative	1.69 (1.05-2.71)	0.031	1.18 (0.65-2.12)	0.58
Pak1 cytoplasmic expression				
Overexpressed vs normal	1.79 (1.17-2.74)	0.0068	1.98 (1.14-3.46)	0.016
Pak1 nuclear expression				
Positive vs negative	1.05 (0.64-1.73)	0.84	0.76 (0.39-1.51)	0.44
pER ^{ser305} nuclear expression				
Positive vs negative	0.81 (0.53-1.24)	0.32	0.95 (0.56-1.61)	0.84

Pak1 in the tumor were highly sensitive to tamoxifen treatment ($P = 0.00003$; Fig. 2A), whereas expression of Pak1 in the nucleus was associated with less marked benefit from the treatment ($P = 0.15$; Fig. 2B). Pak1 protein expression in the cytoplasm did not influence the efficacy of tamoxifen in this series of patients. Similarly, lack of pER^{ser305} staining resulted in good response to treatment ($P = 0.0003$; Fig. 2C), whereas for positive staining, the benefit was reduced ($P = 0.13$; Fig. 2D). However, when tested for interaction, neither Pak1 nor pER^{ser305} nuclear staining were significantly correlated with reduced effect of tamoxifen (Table 2). Pak1 is phosphorylating serine 305 on the ER α and this may result in tamoxifen resistance in breast tumors. Analyzing Pak1 and pER^{ser305} nuclear expression together could give a further hint on how tamoxifen response is affected by this pathway. The results showed tamoxifen benefit in patients with normal or no expression of any of the two proteins ($P < 0.00001$; Fig. 2E). Patients with tumors expressing the two proteins simultaneously within the nucleus did not seem to respond to treatment ($P = 0.63$; Fig. 2F), and we found a significant interaction with tamoxifen benefit (Table 2).

In this study, we detected the nuclear staining of pER^{ser305} in a group of patients previously defined as ER α negative by standard methods. Adding these pER^{ser305}-positive tumors to the ER α -positive group when analyzing tamoxifen response reinforced the results for both Pak1 nuclear expression ($P_{\text{interaction}} = 0.15$), pER^{ser305} nuclear expression ($P = 0.098$), as well as for positive staining of the two proteins in combination ($P = 0.013$), showing reduced benefit from tamoxifen treatment when these proteins were expressed.

Prognosis. We used patients randomized to no adjuvant tamoxifen treatment to evaluate prognosis. Cytoplasmic expression of the Pak1 protein correlated with decreased recurrence-free survival (overexpressed versus normal:

HR, 1.80; 95% CI, 1.21-2.67; $P = 0.0027$; Fig. 3A). This was true also when restricting the analysis to ER-positive patients (HR, 2.00; 95% CI, 1.26-3.17; $P = 0.0023$), but not in the ER-negative group ($P = 0.53$). Cytoplasmic Pak1 also correlated with decreased breast cancer-specific survival (HR, 1.95; 95% CI, 1.17-3.26; $P = 0.0086$). In a multivariate analysis, cytoplasmic Pak1 protein was significantly related to increased recurrence rate and increased breast cancer mortality (Table 3). Pak1 nuclear expression was not prognostic (HR, 0.99; 95% CI, 0.63-1.53; $P = 0.95$; Fig. 3B). pER^{ser305}-positive expression in the nucleus indicated better prognosis than no expression (HR, 0.67; 95% CI, 0.46-0.99; $P = 0.039$; Fig. 3C), but this relationship did not remain significant in multivariate analysis (Table 3).

Discussion

Pak1 may contribute to tamoxifen resistance in breast cancer patients by phosphorylation of the serine 305 within the activating function-2 region of the ER α ligand binding domain. Here, we report nuclear and cytoplasmic overexpression of Pak1 in 22% and 58% of tumors, respectively. Nuclear expression of the phosphorylated ER α ^{ser305} was detected in 36%.

Recently, we found that amplification of the gene encoding Pak1 predicts poor response to tamoxifen (18). Other studies showed that Pak1 is a kinase that interacts with ER α and phosphorylates the serine 305 residue, and this modulation triggers activation of the receptor (7, 8). Therefore, we investigated the possible predictive value of this signaling pathway in this series of breast tumors stained for Pak1 and pER^{ser305}. Interestingly, in Pak1-positive tumors, nuclear localization of Pak1 was correlated to pER^{ser305}, but we did not find a significant correlation between Pak1 and pER^{ser305} expression over-

all. An explanation to this result could be another protein, PKA, and possibly other kinases also phosphorylating the ER α at the same position (6). Hypothetically, PKA may be the dominant kinase and comparatively high levels of nuclear active Pak1 could be needed to compete with the action of PKA. A cooperative role of the two proteins may also be considered. Hence, our result does not exclude the importance of Pak1 phosphorylation of the ER α in subgroups of breast tumors. Worth noticing is that the Pak1 antibody used in this study was not selective for kinase-active Pak1. However, among relapsing tamoxifen-treated patients, the nuclear coexpression of Pak1/pER α^{ser305} correlated significantly ($P = 0.023$; data not shown). This may indicate that Pak1 phosphorylated ER α in these patients and counteracted tamoxifen. Pak1 in the nucleus and pER α^{ser305} correlated to similar clinicopathologic factors, seen in Table 1, whereas Pak1 in the cytoplasm correlated inversely to these factors, indicating nuclear interaction of ER α and Pak1.

In line with our previous findings and other experimental studies, we found that Pak1 located to the nucleus indicated reduced response to tamoxifen treatment (16, 18, 19). A similar result was observed for pER α^{ser305} staining in the nucleus, which has been suggested in a recent study of premenopausal breast cancer (20). The expression of these two proteins together suggested a further reduction of tamoxifen response, indicating the importance of this signaling pathway for treatment prediction. However, the group with simultaneous expression, includes <10% of the ER-positive patients and the data must be interpreted with caution.

Activation of the ER α is a highly regulated, not yet fully elucidated process, involving phosphorylation of several amino acid residues and coregulator recruitment in a dynamic and possibly cyclic system (21). An imbalance in this process may cause changes in the gene-expression pattern, regulated by the receptor, and influence receptor-dependent malignant cells to develop drug resistance mechanisms. Other ER α phospho sites have previously been studied in breast cancer. In contrast to results seen for the phosphorylation of the serine 305 site, detection of pER α^{ser118} in the ligand-independent domain of the ER α was shown to associate with a differentiated phenotype, improved survival, and maintenance of tamoxifen sensitivity (22–24). Extracellular signal-regulated kinase 1/2 in the mitogen-activated protein kinase signaling pathway phosphorylates ER α^{ser118} , leading to the dissociation of the receptor with its coactivator AIB1 in the presence of tamoxifen (25). Phosphorylation of this site has been found to recruit both coactivators and corepressors, possibly in a dynamic time-dependent fashion (26, 27). Interestingly, phosphorylation of serine 118 is suggested to enhance the estrogen-driven activity of the ER, driving the oncogenesis forward (28). Ligand-dependent transcription of the ER α target genes, *pS2*, *CCND1*, *c-myc*, and *PR* has been detected in a separate study, reinforcing the hypothesis that detection of pER α^{ser118} in breast tumor is a marker for tamoxifen sensitivity (29).

Expression of pER α^{ser167} increased survival and tamoxifen response after relapse, in a small study based on tamoxifen-treated breast cancer patients (30). In contrast, other results have indicated that activation of Akt, which may be responsible for the phosphorylation of ER α^{ser167} in breast cancer, may be linked to reduced tamoxifen sensitivity (17, 31). From these studies, no consensus about the role of the different phospho sites in the ER α can be established.

Apart from modulating the ER α in the nucleus, Pak1 plays multiple roles in the cytoplasmic compartment. Pak1 is a serine/threonine kinase involved in several signaling pathways in the normal breast epithelial cell, as well as in oncogenic regulation in the neoplastic cell. Cytoskeletal organization, anchorage-independent growth, cell invasion, migration, and antiapoptosis are examples of cellular functions in which Pak1 is important (32–35). Recent studies have examined the role of Pak1 in several different malignancies, such as glioblastoma, hepatocellular carcinoma, renal cell carcinoma, and uveal carcinoma (36–39). They all support the involvement of cytoplasmic Pak1 in invasion, migration, and metastasis. Activation of the Rho family GTPases Cdc42 and Rac1, regulators of Pak1 kinase activity, was suggested to induce antiestrogen resistance in breast cancer cell lines through a Pak1-dependent pathway (40). However, the Pak1 target ER α is thought to localize to the cytoplasm under certain conditions in the breast cancer cell, binding to membrane-associated proteins as for example growth factor receptors (41, 42). Whether Pak1 influences ER α in this setting is yet to be elucidated.

We observed that cytoplasmic Pak1 overexpression indicated worse outcome in the tamoxifen-nontreated group. Functional cell experiments have given Pak1 several roles in tumorigenesis, strengthening its prognostic role (32–35). These results are not fully in line with previous smaller studies in which prognosis did not seem to be related to cytoplasmic Pak1 (16, 18). Shou et al. (41) suggest that tamoxifen may also be an agonist on the ER α , when located to the cell membrane, in which ER α signaling is thought to cross-talk with growth factor signaling pathways. We did not observe membrane-associated staining by the pER α^{ser305} antibody. The explanation to this may be that all tumors analyzed were primary tumors, in which acquired resistance has not yet occurred. This study is limited to postmenopausal patients with an early breast cancer. To confirm the observations made in the present study, analyses of activated Pak1 in primary tumors and metastases might be of importance. In addition, the role of pER α^{ser305} in combination with other ER phospho sites in treatment prediction needs further inquiry.

In conclusion, nuclear Pak1 and pER α^{ser305} may be involved in resistance to tamoxifen treatment in postmenopausal breast cancer and could be of interest as drug targets for patients receiving tamoxifen treatment.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Birgitta Holmlund and Torsten Hägersten for the excellent technical assistance.

Grant Support

The Swedish Cancer Society, the Swedish Research Council and King Gustaf V Jubilee Fund.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 07/03/2009; revised 11/25/2009; accepted 12/16/2009; published OnlineFirst 02/23/2010.

References

- Bray F, McCarron P, Parkin DM. The changing global patterns of female breast cancer incidence and mortality. *Breast Cancer Res* 2004;6:229–39.
- Early Breast Cancer Trialists' Collaborative Group. Tamoxifen for early breast cancer: an overview of the randomised trials. *Lancet* 1998;351:1451–67.
- Girault I, Bieche I, Lidereau R. Role of estrogen receptor α transcriptional coregulators in tamoxifen resistance in breast cancer. *Maturitas* 2006;54:342–51.
- Orti E, Bodwell JE, Munck A. Phosphorylation of steroid hormone receptors. *Endocr Rev* 1992;13:105–28.
- Lannigan DA. Estrogen receptor phosphorylation. *Steroids* 2003;68:1–9.
- Michalides R, Griekspoor A, Balkenende A, et al. Tamoxifen resistance by a conformational arrest of the estrogen receptor α after PKA activation in breast cancer. *Cancer Cell* 2004;5:597–605.
- Wang RA, Mazumdar A, Vadlamudi RK, Kumar R. P21-activated kinase-1 phosphorylates and transactivates estrogen receptor- α and promotes hyperplasia in mammary epithelium. *EMBO J* 2002;21:5437–47.
- Balasenthil S, Barnes CJ, Rayala SK, Kumar R. Estrogen receptor activation at serine 305 is sufficient to upregulate cyclin D1 in breast cancer cells. *FEBS Lett* 2004;567:243–7.
- Tharakan R, Lepont P, Singleton D, Kumar R, Khan S. Phosphorylation of estrogen receptor α , serine residue 305 enhances activity. *Mol Cell Endocrinol* 2008;295:70–8.
- Talukder AH, Li DQ, Manavathi B, Kumar R. Serine 28 phosphorylation of NRIF3 confers its co-activator function for estrogen receptor- α transactivation. *Oncogene* 2008;27:5233–42.
- Zwart W, Griekspoor A, Rondaj M, Verwoerd D, Neeffjes J, Michalides R. Classification of anti-estrogens according to intramolecular FRET effects on phospho-mutants of estrogen receptor α . *Mol Cancer Ther* 2007;6:1526–33.
- Bokoch GM. Biology of the p21-activated kinases. *Annu Rev Biochem* 2003;72:743–81.
- Kumar R, Hung MC. Signaling intricacies take center stage in cancer cells. *Cancer Res* 2005;65:2511–5.
- Rutqvist LE, Johansson H. Long-term follow-up of the randomized Stockholm trial on adjuvant tamoxifen among postmenopausal patients with early stage breast cancer. *Acta Oncol* 2007;46:133–45.
- Skliiris GP, Rowan BG, Al-Dhaheer M, et al. Immunohistochemical validation of multiple phospho-specific epitopes for estrogen receptor α (ER α) in tissue microarrays of ER α positive human breast carcinomas. *Breast Cancer Res Treat* 2008.
- Holm C, Rayala S, Jirstrom K, Stal O, Kumar R, Landberg G. Association between Pak1 expression and subcellular localization and tamoxifen resistance in breast cancer patients. *J Natl Cancer Inst* 2006;98:671–80.
- Al-Dhaheer MH, Rowan BG. Application of phosphorylation site-specific antibodies to measure nuclear receptor signaling: characterization of novel phosphoantibodies for estrogen receptor α . *Nucl Recept Signal* 2006;4:e007.
- Bostner J, Ahnstrom Waltersson M, Fornander T, Skoog L, Nordenskjold B, Stal O. Amplification of CCND1 and PAK1 as predictors of recurrence and tamoxifen resistance in postmenopausal breast cancer. *Oncogene* 2007;26:6997–7005.
- Rayala SK, Talukder AH, Balasenthil S, et al. P21-activated kinase 1 regulation of estrogen receptor- α activation involves serine 305 activation linked with serine 118 phosphorylation. *Cancer Res* 2006;66:1694–701.
- Holm C, Kok M, Michalides R, et al. Phosphorylation of the oestrogen receptor α at serine 305 and prediction of tamoxifen resistance in breast cancer. *J Pathol* 2009;217:372–9.
- Green KA, Carroll JS. Oestrogen-receptor-mediated transcription and the influence of co-factors and chromatin state. *Nat Rev Cancer* 2007;7:713–22.
- Bergqvist J, Elmberger G, Ohd J, et al. Activated ERK1/2 and phosphorylated oestrogen receptor α are associated with improved breast cancer survival in women treated with tamoxifen. *Eur J Cancer* 2006;42:1104–12.
- Murphy L, Cherlet T, Adeyinka A, Niu Y, Snell L, Watson P. Phosphoserine-118 estrogen receptor- α detection in human breast tumors *in vivo*. *Clin Cancer Res* 2004;10:1354–9.
- Murphy LC, Niu Y, Snell L, Watson P. Phospho-serine-118 estrogen receptor- α expression is associated with better disease outcome in women treated with tamoxifen. *Clin Cancer Res* 2004;10:5902–6.
- Glaros S, Atanaskova N, Zhao C, Skafar DF, Reddy KB. Activation function-1 domain of estrogen receptor regulates the agonistic and antagonistic actions of tamoxifen. *Mol Endocrinol* 2006;20:996–1008.
- Bunone G, Briand PA, Miksicsek RJ, Picard D. Activation of the unliganded estrogen receptor by EGF involves the MAP kinase pathway and direct phosphorylation. *EMBO J* 1996;15:2174–83.
- Gburcik V, Picard D. The cell-specific activity of the estrogen receptor α may be fine-tuned by phosphorylation-induced structural gymnastics. *Nucleic Recept Signal* 2006;4:e005.
- Kato S, Endoh H, Masuhiro Y, et al. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* 1995;270:1491–4.
- Weitsman GE, Li L, Skliiris GP, et al. Estrogen receptor- α phosphorylated at Ser118 is present at the promoters of estrogen-regulated genes and is not altered due to HER-2 overexpression. *Cancer Res* 2006;66:10162–70.
- Yamashita H, Nishio M, Kobayashi S, et al. Phosphorylation of estrogen receptor α serine 167 is predictive of response to endocrine therapy and increases postrelapse survival in metastatic breast cancer. *Breast Cancer Res* 2005;7:R753–64.
- Perez-Tenorio G, Stal O. Activation of AKT/PKB in breast cancer predicts a worse outcome among endocrine treated patients. *Br J Cancer* 2002;86:540–5.
- Adam L, Vadlamudi R, Kondapaka SB, Chernoff J, Mendelsohn J, Kumar R. Heregulin regulates cytoskeletal reorganization and cell migration through the p21-activated kinase-1 via phosphatidylinositol-3 kinase. *J Biol Chem* 1998;273:28238–46.
- Schurmann A, Mooney AF, Sanders LC, et al. p21-activated kinase 1 phosphorylates the death agonist bad and protects cells from apoptosis. *Mol Cell Biol* 2000;20:453–61.
- Mazumdar A, Kumar R. Estrogen regulation of Pak1 and FKHR pathways in breast cancer cells. *FEBS Lett* 2003;535:6–10.

35. Vadlamudi RK, Adam L, Wang RA, et al. Regulatable expression of p21-activated kinase-1 promotes anchorage-independent growth and abnormal organization of mitotic spindles in human epithelial breast cancer cells. *J Biol Chem* 2000;275:36238-44.
36. Aoki H, Yokoyama T, Fujiwara K, et al. Phosphorylated Pak1 level in the cytoplasm correlates with shorter survival time in patients with glioblastoma. *Clin Cancer Res* 2007;13:6603-9.
37. Ching YP, Leong VY, Lee MF, Xu HT, Jin DY, Ng IO. P21-activated protein kinase is overexpressed in hepatocellular carcinoma and enhances cancer metastasis involving c-Jun NH2-terminal kinase activation and paxillin phosphorylation. *Cancer Res* 2007;67:3601-8.
38. O'Sullivan GC, Tangney M, Casey G, Ambrose M, Houston A, Barry OP. Modulation of p21-activated kinase 1 alters the behavior of renal cell carcinoma. *Int J Cancer* 2007;121:1930-40.
39. Pavey S, Zuidervaart W, van Nieuwpoort F, et al. Increased p21-activated kinase-1 expression is associated with invasive potential in uveal melanoma. *Melanoma Res* 2006;16:285-96.
40. Cai D, Iyer A, Felekis KN, et al. AND-34/BCAR3, a GDP exchange factor whose overexpression confers antiestrogen resistance, activates Rac, PAK1, and the cyclin D1 promoter. *Cancer Res* 2003;63:6802-8.
41. Shou J, Massarweh S, Osborne CK, et al. Mechanisms of tamoxifen resistance: increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer. *J Natl Cancer Inst* 2004;96:926-35.
42. Chung YL, Sheu ML, Yang SC, Lin CH, Yen SH. Resistance to tamoxifen-induced apoptosis is associated with direct interaction between Her2/neu and cell membrane estrogen receptor in breast cancer. *Int J Cancer* 2002;97:306-12.

Clinical Cancer Research

Estrogen Receptor- α Phosphorylation at Serine 305, Nuclear p21-Activated Kinase 1 Expression, and Response to Tamoxifen in Postmenopausal Breast Cancer

Josefine Bostner, Lambert Skoog, Tommy Fornander, et al.

Clin Cancer Res 2010;16:1624-1633. Published OnlineFirst February 23, 2010.

Updated version	Access the most recent version of this article at: doi:10.1158/1078-0432.CCR-09-1733
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2010/03/01/1078-0432.CCR-09-1733.DC1

Cited articles	This article cites 41 articles, 14 of which you can access for free at: http://clincancerres.aacrjournals.org/content/16/5/1624.full#ref-list-1
Citing articles	This article has been cited by 8 HighWire-hosted articles. Access the articles at: http://clincancerres.aacrjournals.org/content/16/5/1624.full#related-urls

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://clincancerres.aacrjournals.org/content/16/5/1624 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.