

The Design of Phase II Clinical Trials Testing Cancer Therapeutics: Consensus Recommendations from the Clinical Trial Design Task Force of the National Cancer Institute Investigational Drug Steering Committee

Lesley Seymour¹, S. Percy Ivy², Daniel Sargent³, David Spriggs⁴, Laurence Baker⁵, Larry Rubinstein², Mark J. Ratain⁶, Michael Le Blanc⁷, David Stewart⁸, John Crowley⁷, Susan Groshen⁹, Jeffrey S. Humphrey¹⁰, Pamela West¹¹, and Donald Berry⁸

Abstract

The optimal design of phase II studies continues to be the subject of vigorous debate, especially studies of newer molecularly targeted agents. The observations that many new therapeutics “fail” in definitive phase III studies, coupled with the numbers of new agents to be tested as well as the increasing costs and complexity of clinical trials, further emphasize the critical importance of robust and efficient phase II design. The Clinical Trial Design Task Force (CTD-TF) of the National Cancer Institute (NCI) Investigational Drug Steering Committee (IDSC) has published a series of discussion papers on phase II trial design in *Clinical Cancer Research*. The IDSC has developed formal recommendations about aspects of phase II trial design that are the subject of frequent debate, such as endpoints (response versus progression-free survival), randomization (single-arm designs versus randomization), inclusion of biomarkers, biomarker-based patient enrichment strategies, and statistical design (e.g., two-stage designs versus multiple-group adaptive designs). Although these recommendations in general encourage the use of progression-free survival as the primary endpoint, randomization, inclusion of biomarkers, and incorporation of newer designs, we acknowledge that objective response as an endpoint and single-arm designs remain relevant in certain situations. The design of any clinical trial should always be carefully evaluated and justified based on characteristic specific to the situation. *Clin Cancer Res*; 16(6); 1764–9. ©2010 AACR.

Background

Many new drugs targeting molecular pathways are ready for clinical development, necessitating the use of efficient trial designs to quickly and accurately identify promising agents, while also identifying those for which all further development should be stopped. Although the develop-

ment of some drugs is discontinued after phase I, the major drug development decision is generally made on the basis of phase II results. Although traditional oncology trial designs using the endpoint of response and a single arm design seem to have done this task reasonably well for cytotoxic agents, the same does not seem to be true for newer agents in which high rates of tumor shrinkage may not be expected, nor for combinations of agents (such as a new drug combined with standard treatments). Certainly, success rates for phase III trials seem to be decreasing (1). This decrease has led to considerable scientific discussion, debating the advantages and disadvantages of using response versus progression (2) or other imaging endpoints (3), single arm versus randomized designs (4), patient enrichment and biomarker endpoints (5), and optimal statistical designs, such as adaptive design or phase I–II designs.

The Investigational Drug Steering Committee (IDSC) of the National Cancer Institute Cancer Therapy and Evaluation Program (NCI CTEP) appointed a Clinical Trial Design Task Force to advise on the design of early (phase I and II) clinical trials (Table 1). In keeping with its broad mandate, Task Force members include IDSC members, as well as external representation from academia and the

Authors' Affiliations: ¹NCIC Clinical Trials Group, Kingston, Ontario, Canada; ²National Cancer Institute, Bethesda, Maryland; ³Mayo Clinic, Division of Biomedical Statistics and Informatics, Rochester, Minnesota; ⁴Memorial Sloan-Kettering Cancer Center, Department of Medicine, New York, New York; ⁵University of Michigan, Department of Medicine and Pharmacology, Ann Arbor, Michigan; ⁶University of Chicago, Department of Medicine, Section of Hematology/Oncology, Chicago, Illinois; ⁷Fred Hutchinson Cancer Research Center, Cancer Research and Biostatistics, Seattle, Washington; ⁸University of Texas M. D. Anderson Cancer Center, Houston, Texas; ⁹Keck School of Medicine, Norris Comprehensive Cancer Center, University of Southern California, Los Angeles, California; ¹⁰Bristol-Myers Squibb, Plainsboro, New Jersey; and ¹¹The Emmes Corporation, Rockville, Maryland on behalf of the Clinical Trial Design Task Force

Corresponding Author: Lesley Seymour, NCIC Clinical Trials Group, 10 Stuart Street, Kingston, ON, Canada K7L 3N6. Phone: 613-533-6430; Fax: 613-533-2941; E-mail: lseymour@ctg.queensu.ca.

doi: 10.1158/1078-0432.CCR-09-3287

©2010 American Association for Cancer Research.

Table 1. Objectives of the clinical trial design task force**Objectives**

Provide guidance to the IDSC on current best practice for all aspects of early clinical trial design
 Identify areas in which further research and investigation are needed to improve the quality and/or efficiency of early clinical trial design
 Assist NCI and the IDSC or as needed with the implementation of recommendations for further research and investigation
 Collaborate with other IDSC task forces to minimize duplication of efforts
 Develop and implement a plan to disseminate and publish the recommendations and guidance formulated by the task force and approved by the IDSC

pharmaceutical industry (Table 2). Members have expertise in early clinical trial design, conduct, and analyses, and the Task Force is composed of statisticians, clinicians, imaging specialists, pharmacologists, biomarker experts, radiation and systems biologists, as well as patient advocates. One of the first initiatives of the Task Force was to coordinate a Phase II Workshop attended by Task Force members, other IDSC members, and a number of invited experts in the field. This Phase II Workshop formed the basis of a series of discussion documents on all aspects of phase II design and conduct, published in this journal (1–5). Subsequently, at the request of the IDSC, the Task Force formulated specific recommendations for the design of phase II clinical trials (Figs. 1 and 2). These recommendations, although based on the original Phase II Workshop and subsequent publications, underwent extensive discussion and revision to ensure broad applicability and acceptance and were formally approved by the IDSC.

We report here on the IDSC's recommendations about phase II trial design.

Review and Discussion

Types of phase II trials. Although the most common grouping of phase II trials is by design, i.e., single arm and randomized trials (Fig. 3), another conceptual grouping is the ultimate aim of the trial. Is the trial to be used to screen for any evidence of activity of a new drug that has recently completed phase I testing, to look for preliminary hints of activity, and to guide selection of tumor types for further study? Or, is the trial designed to provide a go–no go answer to allow the conduct of a definitive registration trial in a specific disease? The Task Force considered a number of ways to categorize phase II trials, including the concept of screening (e.g., IIa) versus decision-making (i.e., IIb, sufficiently robust to support progression to phase III). However, the consensus opinion was that for most drugs, conducting two phase II trials in sequence would be inefficient. We recognized that there are circumstances in which such a “proof of concept” approach might be reasonable, such as seeking a signal about the selection of tumor types for further study (e.g., when not readily apparent from preclinical or phase I studies) or for biomarker-based studies to validate a proposed mechanism of action. Ideally, however, these concepts would

be embedded, possibly adaptively, in a single phase II trial. Thus, we used single-arm versus randomized studies as our primary categorization of phase II trials (Fig. 3).

Selection of the appropriate primary endpoint. The advantages and limitations of objective tumor response as the primary endpoint in phase II trials, and alternative endpoints such as multinomial endpoints (6), response as a continuous variable, progression-free survival, imaging, patient reported outcomes (7), and biomarker endpoints were reviewed by Dhani (2), Shankar (3), and McShane (5), and will not be further addressed here. We believe that the decision on the most appropriate endpoint is critical to inform the appropriate design. Although the Task Force accepted that response-based endpoints are still relevant for some agents (when tumor shrinkage and clinically relevant response rates are expected) and some trials, the recommendations emphasize the need to consider the inclusion of a progression-free survival primary endpoint as more informative (8, 9). Overall survival is not recommended as an endpoint, as subsequent therapy may confound conclusions, and progression is usually substantially earlier, thus shortening the duration of the trial and follow up.

Randomization, blinding, and crossover. The Task Force agreed that randomization was generally required to evaluate the efficacy of combinations of agents (e.g., for approved drugs and investigational agents). Randomization is usually essential for a phase II trial in which progression-free survival is the most appropriate endpoint. Nonetheless, single arm designs are still appropriate for the evaluation of a monotherapy or when a well-defined historical control database is available (10). As for any trial the design and the selected null and alternative hypotheses must be carefully justified. If a randomized design is selected, blinding of the agents (against placebo, other doses of the same agent, or other active agents) should be considered. When the primary endpoint is progression-free survival or response based, designs that allow crossover after progression maintain the integrity of the study and can provide additional data that could inform the future development of the agent.

Biomarkers. Biomarkers are of considerable interest in the setting of the phase II study, but they present significant challenges in their incorporation, measurement, and interpretation. In most instances, the biomarker is

not clinically validated as a predictive marker (of efficacy) early in the development of a new agent (i.e., at the time of the phase II trial). The IDSC's recommendations about biomarker use in early clinical trials are detailed in the current Focus Series (11–16). Because of these limitations, these recommendations encourage the prospective inclusion of molecular markers in phase II trials to evaluate predictive markers, but discourage prospective patient selection on the basis of a biomarker (unless already clinically validated), except in the setting of an appropriate (and explicit) adaptive design. Phase II trials including patients with a specific biomarker but with multiple histological subtypes were considered of particular interest and may be a more efficient screening tool, especially when combined with an adaptive design.

Statistical designs. Improved efficiencies in clinical trial design with associated shortening of development times for effective agents are highly desirable. Numerous designs have been proposed, including randomized selection designs (pick-the-winner), adaptive designs (17), randomized discontinuation designs (18), and other randomized designs (19). Prospectively specified adaptive designs are of particular interest in the context of phase II studies of molecularly targeted agents in which biomarker identification and validation may be emergent during the conduct of the trial, limiting the ability to select patients or identify optimal doses and/or schedules at the trial outset. Such adaptive designs are also particularly useful for trials including patients with a range of histologic subtypes but with biomarkers of interest. Adaptive designs in such settings should be efficient and may result in improved precision. Despite the multiplicity of new designs that have been proposed, their inclusion in new trials has in general been modest at best (20, 21). Reasons postulated include requirements for statistical support as well as concerns about robustness, accrual, and cost.

The Task Force is strongly supportive of designs that improve efficiency and shorten development time, such as adaptive designs, but recognized the need to continue to formally evaluate these designs to encourage wider acceptance and implementation. An ongoing initiative is the cre-

ation of a database to allow the formal testing, *in silico* of newer designs, in order to validate their use in future trials.

Interestingly, although formulated prior to the publication of the editorial, these recommendations are congruent with a review of phase II trials published in the *Journal of Clinical Oncology* (22), as well as with other reviews and recommendations (23).

Consensus Recommendations

Choosing the appropriate primary endpoint

The first and critical decision point for the design of a phase II trial is based on the choice of the most appropriate primary endpoint, which should be tailored to the disease and drug(s) under investigation.

- Response-based endpoints such as those defined by Response Evaluation Criteria In Solid Tumors (RECIST), are standard, especially in early phase II trials. Other qualified biomarkers, such as molecular imaging or tumor markers, may be appropriate in select circumstances. Response-based endpoints are appropriate primary endpoints if unambiguous and clinically relevant direct antitumor activity (such as tumor shrinkage) is hypothesized.
- If a response-based endpoint is not appropriate, especially in later phase II trials, progression-free survival is recommended as the primary endpoint. Other biomarker endpoints (such as tumor burden, tumor markers, novel imaging, tumor response, molecular biomarkers) and patient-reported outcomes are always encouraged as secondary endpoints, especially in the context of studies that aim to qualify such endpoints. It is acknowledged that once qualified, these biomarker endpoints will become appropriate primary endpoints.

Study design: primary endpoint is tumor response

Monotherapy trials. Single arm designs are acceptable. However, randomization should be encouraged to optimize dose and schedule or to benchmark activity against known active therapies.

Table 2. Past and present members of the clinical trial design task force

Position	Member
Chair	Lesley Seymour
Co-chair	Donald Berry
CTEP	S. Percy Ivy
Clinical and/or pharmacology	A. Adjei, S. Yao, L. Baker, S. Lutzker, J. Humphrey, D. Stewart, A. Dowlati, P. Keegan, P. LoRusso, M. Ratain, D. Spriggs, J. Collins, M. Grever, C. Erlichmann
Statistics	J. Crowley, S. Groshen, M. Le Blanc, L. Rubinstein, D. Sargent
Imaging	L. Shankar, A. Shields
Advocate	D. Collyar
Nonvoting	R. Agarwal, L. Minasian, P. Ujhazy, L. Jensen, P. West
Past members	G. Eckhart, S. Arbuck, M. Christian, G. Fyfe, R. Humphrey, M. Sznol, M. Villalona-Calero, M. Weinblatt




Fig. 1. Process for development of recommendations.

Combination trials. With some exceptions (e.g., availability of a well-validated robust control database), randomization is usually required for trials testing combinations of agents to establish efficacy. An example is standard therapy \pm novel agent or combinations of novel agents.

Study design: primary endpoint is progression-free survival

Monotherapy or combination trials

1. With some exceptions (e.g., availability of a robust control database), randomization is required.
2. For randomized trials, blinded designs are encouraged when feasible. Although placebo controlled trials are challenging, they are encouraged whenever possible. Alternatives include dose ranging, randomization versus active controls or other novel agents, and randomized discontinuation and other crossover designs.
3. It may be informative to prospectively incorporate crossover to the standard therapy + novel agent for those patients initially assigned to the standard therapy

alone, although careful consideration should be given to the timing of crossover (e.g., only after the primary endpoint has been observed). Such crossover designs increase the access of patients to investigational agents, and also provide additional information about the activity of the study arms.

Patient selection and enrichment strategies

Monotherapy or combination trials

1. A goal of phase (I and) II development should be to define biomarkers predictive of efficacy and/or toxicity. When feasible and appropriate, molecular biomarkers should be explored in order to identify subsets of patients of interest for future study.
2. Enrollment should, in general, not be limited by biomarker status unless there are strong confirmatory and supportive clinical data justifying the enrichment strategy. Adaptive statistical designs may be used to allow modification of enrollment if data suggest a biomarker is predictive.
3. In an unselected trial (i.e., patients not defined by a biomarker), the patient population of primary




Fig. 2. Process and/or flow or approaches for determination of phase II trial design recommendations. PRO, patient related outcomes; PFS, progression-free survival.

interest (i.e., a cohort defined by a biomarker) should be predefined and the study powered accordingly to detect an effect in that subset.

- Multidisease phase II designs should be considered, especially if the objective is to test a biomarker-focused hypothesis.

Statistical designs

Prospective designs that adapt to what is learned during the trial can improve the efficiency of drug development and provide greater precision. Available adaptations include stopping early, continuing longer than anticipated, dropping arms (or doses), adding arms, focusing




Fig. 3. Types of phase II studies.

on patient subsets, assignment of better performing treatment arms with greater probability, and seamlessly moving from phase I to II or phase II to III during a single trial.

Conclusions

The Task Force formulated recommendations (Fig. 2) for the design of phase II trials of anticancer agents on the basis of consensus gained during a workshop and extensive discussions with members of the IDSC, the Task Force, and external experts. These recommendations were subsequently approved by the IDSC. Although these recommendations in general encouraged the use of progression-free survival as the primary endpoint, the use of randomization, the inclusion of biomarkers, and the use of newer designs, they acknowledge that objective

response and single-arm design remain relevant in appropriate circumstances. The design of any clinical trial should always be carefully evaluated and justified.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The Clinical Trial Design Task Force would like to acknowledge the support of the NCI and the IDSC members and chairs, Amy Gravell, LeeAnn Jensen, the attendees of the original Phase II Workshop, as well as all Task Force members, past and present, especially Drs. Alex Adjei and Michael Christian who originally led the Task Force (Table 2).

Received 12/15/2009; revised 01/08/2010; accepted 01/13/2010; published OnlineFirst 03/09/2010.

References

- Adjei AA, Christian M, Ivy P. Novel designs and end points for phase II clinical trials. *Clin Cancer Res* 2009;15:1866–72.
- Dhani N, Tu D, Sargent DJ, Seymour L, Moore MJ. Alternate end-points for screening phase II studies. *Clin Cancer Res* 2009;15:1873–82.
- Shankar LK, Van den Abbeele A, Yap J, Benjamin R, Scheutze S, FitzGerald TJ. Considerations for the use of imaging tools for phase II treatment trials in oncology. *Clin Cancer Res* 2009;15:1891–7.
- Rubinstein L, Crowley J, Ivy P, LeBlanc M, Sargent D. Randomized phase II designs. *Clin Cancer Res* 2009;15:1883–90.
- McShane LM, Hunsberger S, Adjei AA. Effective incorporation of biomarkers into phase II trials. *Clin Cancer Res* 2009;15:1898–905.
- Goffin JR, Tu D. Phase II Stopping rules that employ response rates and early progression. *J Clin Oncol* 2008;26:3715–20.
- Wagner LI, Wenzel L, Shaw E, Cella D. Patient-reported outcomes in phase II cancer clinical trials: lessons learned and future directions. *J Clin Oncol* 2007;25:5058–62.
- Fleming TR, Rothmann MD, Lu HL. Issues in using progression-free survival when evaluating oncology products. *J Clin Oncol* 2009;27:2874–80.
- Freidlin B, Korn EL, Hunsberger S, et al. Proposal for the use of progression-free survival in unblinded randomized trials. *J Clin Oncol* 2007;25:2122–6.
- Korn E, Liu P-Y, Lee SJ, et al. Meta-analysis of phase II cooperative group trials in metastatic stage IV melanoma to determine progression-free and overall survival benchmarks for future phase II trials. *J Clin Oncol* 2008;26:527–34.
- LoRusso PM, Boerner SA, Seymour LK. An overview of the optimal planning, design and conduct of phase I studies of new therapeutics. *Clin Cancer Res* 2010;16:1710–8.
- Dancey JE, Dobbin KK, Groshen S, et al. Guidelines for the development and incorporation of biomarker studies in early clinical trials of novel agents. *Clin Cancer Res* 2010;16:1745–55.
- Senderowicz A. Information needed to conduct first-in-human oncology trials in the U.S. A view from a former FDA Medical Reviewer. *Clin Cancer Res* 2010;16:1719–25.
- Rock EP, Molloy VJ, Humphrey JS. Good clinical practice data quality in early drug development. *Clin Cancer Res* 2010;16:1756–63.
- Forster MD, Saijo N, Seymour L, Calvert AH. Performing phase I clinical trials of anticancer agents: perspectives from within the European Union and Japan. *Clin Cancer Res* 2010;16:1737–44.
- Ivy SP, Siu L, Garrett-Mayer E, Rubinstein L. Approaches to phase I clinical trial design focused on safety, efficiency and selected patient populations: A report from the Clinical Trial Design Task Force of the National Cancer Institute Investigational Drug Steering Committee. *Clin Cancer Res* 2010;16:1726–36.
- Coffey CS, Kairalla JA. Adaptive clinical trials: progress and challenges. *Drugs R D* 2008;9:229–42.
- Rosner GL, Stadler W, Ratain M. Randomized discontinuation design: application to cytostatic antineoplastic agents. *J Clin Oncol* 2002;20:4478–84.
- Rubinstein LV, Korn EL, Friedlin B, Hunsberger S, Ivy SP, Smith NA. Design issues of randomized phase II trials and a proposal for phase II screening trials. *J Clin Oncol* 2005;23:7199–206.
- El-Maraghi RH, Eisenhauer EA. Review of phase II trial designs used in studies of molecular targeted agents: outcomes and predictors of success in phase III. *J Clin Oncol* 2008;26:1346–54.
- Lee JJ, Feng L. Randomized phase II designs in cancer clinical trials: current status and future directions. *J Clin Oncol* 2005;23:4450–445.
- Cannistra S. Phase II trials in *Journal of Clinical Oncology*. *J Clin Oncol* 2009;27:3073–6.
- Booth CM, Calvert HA, Giaccone G, et al. Design and conduct of phase II studies of targeted anticancer therapy: recommendations from the task force on methodology for the development of innovative cancer therapies (MDICT). *Eur J Cancer* 2008;44:25–9.

Clinical Cancer Research

The Design of Phase II Clinical Trials Testing Cancer Therapeutics: Consensus Recommendations from the Clinical Trial Design Task Force of the National Cancer Institute Investigational Drug Steering Committee

Lesley Seymour, S. Percy Ivy, Daniel Sargent, et al.

Clin Cancer Res 2010;16:1764-1769. Published OnlineFirst March 9, 2010.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-09-3287](https://doi.org/10.1158/1078-0432.CCR-09-3287)

Cited articles This article cites 23 articles, 21 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/16/6/1764.full#ref-list-1>

Citing articles This article has been cited by 27 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/16/6/1764.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/16/6/1764>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.