

NEMO-Binding Domain Peptide Inhibits Constitutive NF- κ B Activity and Reduces Tumor Burden in a Canine Model of Relapsed, Refractory Diffuse Large B-Cell Lymphoma

Anita Gaurnier-Hausser¹, Reema Patel¹, Albert S. Baldwin^{4,5}, Michael J. May², and Nicola J. Mason^{1,3}

Abstract

Purpose: Activated B-cell diffuse large B-cell lymphoma (ABC-DLBCL) is an aggressive, poorly chemoresponsive lymphoid malignancy characterized by constitutive canonical NF- κ B activity that promotes lymphomagenesis and chemotherapy resistance via overexpression of antiapoptotic NF- κ B target genes. Inhibition of the canonical NF- κ B pathway may therefore have therapeutic relevance in ABC-DLBCL. Here, we set out to determine whether dogs with spontaneous DLBCL have comparative aberrant constitutive NF- κ B activity and to determine the therapeutic relevance of NF- κ B inhibition in dogs with relapsed, resistant DLBCL.

Experimental Design: Canonical NF- κ B activity was evaluated by electrophoretic mobility shift assays and immunoblot analyses, and NF- κ B target gene expression was measured by quantitative real time PCR. Primary malignant canine B lymphocytes were treated with the selective IKK complex inhibitor NF- κ B essential modulator-binding domain (NBD) peptide and evaluated for NF- κ B activity and apoptosis. NBD peptide was administered intranodally to dogs with relapsed B-cell lymphoma and NF- κ B target gene expression and tumor burden were evaluated pre- and post-treatment.

Results: Constitutive canonical NF- κ B activity and increased NF- κ B target gene expression were detected in primary DLBCL tissue. NBD peptide inhibited this activity and induced apoptosis of primary malignant B cells *in vitro*. Intratumoral injections of NBD peptide to dogs with relapsed DLBCL inhibited NF- κ B target gene expression and reduced tumor burden.

Conclusions: This work shows that dogs with spontaneous DLBCL represent a clinically relevant, spontaneous, large animal model for human ABC-DLBCL and shows the therapeutic relevance of NF- κ B inhibition in the treatment of ABC-DLBCL. These results have important translational relevance for ABC-DLBCL treatment in human patients. *Clin Cancer Res*; 17(14); 4661–71. ©2011 AACR.

Introduction

Diffuse large B-cell lymphoma (DLBCL) is the most common adult lymphoid malignancy with approximately 30,000 new cases diagnosed each year in the United States (1, 2). Gene expression profiling has revealed the presence of at least 3 subtypes of DLBCL, activated B-cell (ABC-DLBCL), germinal center B-cell (GCB-DLBCL), and primary mediastinal B-cell lymphoma (PMBL), that have distinct molecular signatures and different clinical outcomes (1, 3). ABC-DLBCL is the most aggressive subtype and is less responsive to conventional multiagent che-

motherapy than GCB-DLBCL and PMBL. Even with the addition of rituximab, the overall survival for patients with ABC-DLBCL is 47% (4).

ABC-DLBCL is characterized by the presence of constitutive canonical NF- κ B activity that drives expression of NF- κ B target genes which promote lymphocyte proliferation (e.g., *Cyclin D1* and *D2*) and inhibit apoptosis (e.g., *Bcl-2*, *Bcl-xL*, *A1*, *c-FLIP*, and *XIAP*; refs. 5, 6). Aberrant canonical NF- κ B activity contributes to lymphomagenesis and chemoresistance by promoting malignant cell survival and proliferation (7, 8). In health, NF- κ B activation is highly regulated. NF- κ B dimers are held inactive in the cytoplasm through their association with inhibitory I κ B family members (9). Engagement of cell surface receptors by many different infectious and inflammatory stimuli leads to phosphorylation and activation of the IKK complex, which phosphorylates I κ B α and targets it for ubiquitination and proteosomal degradation. This allows NF- κ B dimers to translocate to the nucleus where they bind to target gene promoters and enhancers and initiate transcription (8). Somatic activating mutations in genes encoding regulatory proteins upstream of the IKK complex have been identified in patients with ABC-DLBCL and lead to

Authors' Affiliations: Departments of ¹Pathobiology, ²Animal Biology, and ³Clinical Studies, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania; ⁴Lineberger Comprehensive Cancer Center and Department of Biology, University of North Carolina; and ⁵TheraLogics, Inc., Chapel Hill, North Carolina

Corresponding Author: Nicola J. Mason, School of Veterinary Medicine, University of Pennsylvania, 315 Hill Pavilion, 380 South University Avenue, Philadelphia, PA 19104-6010. Phone: 215-898-3996; Fax: 215-746-2295; E-mail: nmason@vet.upenn.edu

doi: 10.1158/1078-0432.CCR-10-3310

©2011 American Association for Cancer Research.

Translational Relevance

Activated B-cell diffuse large B-cell lymphoma (ABC-DLBCL) is an aggressive, poorly chemoresponsive lymphoid malignancy characterized by constitutive canonical NF- κ B activity that promotes lymphomagenesis and chemotherapy resistance via overexpression of antiapoptotic NF- κ B target genes. We have investigated NF- κ B as a therapeutic target by using the dog as a clinically relevant, spontaneous, animal model of DLBCL. We show that as in human ABC-DLBCL, malignant lymphocytes from dogs with DLBCL have constitutive canonical NF- κ B activity and overexpress antiapoptotic genes. We show that the selective IKK inhibitor, NBD peptide inhibits NF- κ B activity *in vitro* and leads to apoptosis of malignant B cells. We show *in vivo* that intratumoral injections of NBD peptide inhibit NF- κ B gene expression and reduce tumor burden in dogs with DLBCL. These studies show the therapeutic relevance of NF- κ B inhibition *in vivo* in a canine DLBCL model and have important translational relevance for the treatment of human ABC-DLBCL.

constitutive phosphorylation and activation of the IKK complex (5, 10, 11). Constitutive, canonical NF- κ B activity is essential for ABC-DLBCL cell survival (3, 7) and inhibition of this pathway with small molecule inhibitors of IKK, I κ B α super repressors, proteasome inhibitors, or inhibitors upstream of the IKK complex promotes cell-cycle arrest, chemotherapeutic sensitivity, and apoptosis in ABC-DLBCL cell lines (3, 7, 12). Taken together, these findings suggest that targeted inhibition of aberrant NF- κ B activity represents a promising therapeutic strategy for patients with primary or relapsed ABC-DLBCL.

The high-quality draft genome sequence of the dog has revealed its close phylogenetic relationship with man, emphasizing the potential benefit of canine models in identifying disease genes and evaluating response to novel therapies (13–16). In particular, the dog is being increasingly recognized as a clinically relevant, spontaneous large animal model for human non-Hodgkin's lymphoma (NHL; refs. 17–19). NHL is the most common, spontaneous, hematopoietic malignancy in dogs, with an annual incidence of 30/100,000 (20). DLBCL is the most common subtype of canine NHL and shares similar biologic, behavioral, molecular, and genetic characteristics with DLBCL in humans (17, 20–22). Dogs with DLBCL are treated with the same cytotoxic agents used in human DLBCL patients that inhibit cell division and induce apoptosis. However, as in human patients, clinical remission is not maintained and 85% to 90% of canine patients relapse with lethal, drug-resistant lymphoma within 6 to 9 months of initial diagnosis and treatment (19, 23).

NF- κ B essential modulator (NEMO)-binding domain (NBD) peptide is a selective inhibitor of the IKK complex that consists of 11 amino acids located at the carboxy terminus of the catalytic IKK β subunit that binds to the

scaffold protein NEMO (24). The core of 6 amino acids in the NBD is also present in IKK α and facilitates its association with NEMO (24–26). NBD peptide inhibits the interaction of both IKK α and IKK β with NEMO and prevents assembly of the IKK complex (25). Fusion of NBD peptide to protein transduction domains, such as that within the *Drosophila* antennapedia protein enables it to enter cells and effectively inhibit canonical NF- κ B activation in response to TNF α , lipopolysaccharide and toll-like receptor ligation (24, 25, 27). Systemic administration of NBD peptide results in potent antiinflammatory effects in animal models of inflammation (24, 28–32). NBD peptide also inhibits constitutive NF- κ B activity in tumor cell lines, including pancreatic carcinoma, squamous cell carcinoma, melanoma, cutaneous T-cell lymphoma, and mantle cell lymphoma, and sensitizes these cells to TNF-related apoptosis-inducing ligand or TNF α -induced apoptosis (33–37). Furthermore, NBD peptide inhibits NF- κ B activation that occurs in response to DNA damage induced by cytotoxic chemotherapeutics (38). Taken together, these findings suggest that targeted inhibition of canonical NF- κ B activity by using NBD peptide may represent an effective therapeutic strategy in patients with ABC-DLBCL.

Before evaluating the therapeutic effects of NBD peptide in human patients with ABC-DLBCL, we conducted studies to evaluate its ability to inhibit constitutive NF- κ B activity in dogs with spontaneous B-cell lymphoma. We show that constitutive canonical NF- κ B activity occurs in the malignant lymph nodes of dogs with DLBCL and that this is associated with overexpression of antiapoptotic NF- κ B target genes. We show that NBD peptide inhibits constitutive NF- κ B activity and induces apoptosis of canine malignant B cells *in vitro*. Furthermore, we show that administration of NBD peptide to dogs with relapsed B-cell lymphoma inhibits the expression of NF- κ B target genes and leads to a reduction in tumor burden. These findings provide proof of concept that NBD peptide can inhibit constitutive NF- κ B activity in a clinically relevant spontaneous canine model of DLBCL and begin to pave the way for phased clinical trials in humans with ABC-DLBCL.

Materials and Methods

Cells and reagents

Canine lymphocytes were cultured in complete RPMI media containing 10% FBS. The human GCB-DLBCL cell line SUDHL-6 was maintained in complete RPMI media, and the human ABC-DLBCL cell line OCI-Ly10 was maintained in complete Iscove's modified essential medium containing 20% FBS. Cell lines were obtained from Dr. Anne Novak (Mayo Clinic Cancer Center, Rochester, MN). Malignant lymph node samples were collected from dogs with either primary or relapsed DLBCL following approval from the University of Pennsylvania's Institutional Animal Care and Use Committee. The histopathologic diagnosis of DLBCL and the cytologic diagnosis of large B-cell lymphoma were made by board-certified veterinary pathologists (M.G. and A.D., see Acknowledgements) and by a

board-certified clinical pathologist (R.P.), respectively. NBD peptide was synthesized and purified by Dr. James I. Elliott (at the Howard Hughes Medical Institute Biopolymer-Keck Foundation Biotechnology Resource Laboratory, Yale University, New Haven, CT) using standard tertbutoxycarbonyl chemistry, cleavage with hydrofluoric acid, and purification by reversed-phase high-performance liquid chromatography (26). TNF α was purchased from Sigma-Aldrich.

Western blot analyses

Cells were lysed in 50 mmol/L Tris-HCL containing 1% NP-40, 150 mmol/L NaCl, 2.5 mmol/L EDTA, 5% glycerol, a 1:50 dilution of protease inhibitor cocktail, and 1:50 dilutions of phosphatase inhibitor cocktails 2 and 3 (Sigma-Aldrich). Protein concentrations were determined by micro BCA assay (Thermo Scientific). Proteins were separated by SDS-PAGE (10%) and transferred to polyvinylidene difluoride membrane. Membranes were probed with polyclonal rabbit anti-human antibodies against phospho-IKK α / β (16A6), pan IKK β (L570), phospho-I κ B α (14D4), and β -actin (4,967; Cell Signaling) or rabbit anti-human pan I κ B α (C-21; SantaCruz Biotechnology). A horseradish peroxidase-conjugated donkey anti-rabbit IgG was used as the secondary detection antibody (Amersham). Blots were developed by using ECL Plus (Amersham) or SuperSignal West Femto (Thermo Scientific).

Electrophoretic mobility shift assay

Whole cell extracts of malignant canine lymphocytes were prepared as previously described (26). Electrophoretic mobility shift assays (EMSA) were done by using 5 μ g protein extract and a palindromic NF- κ B binding sequence probe (Santa Cruz Biotechnology) as previously described (26). EMSA supershift assays were done by using 10 μ g of protein extract and polyclonal rabbit anti-human p65

(sc-109X), p50 (sc-114X), and c-Rel (sc-70X) antibodies or control rabbit IgG (sc-2027; Santa Cruz Biotechnology).

Quantitative reverse transcription PCR

Total RNA extraction was done by using the RNeasy Mini Kit (Qiagen). Reverse transcription was done by using random hexamers and Superscript II reverse transcriptase (Invitrogen Corp.). Transcript sequences for canine *A1*, *c-FLIP*, *Cyclin D1*, and *I κ B α* were obtained from the National Center for Biotechnology Information (<http://www.ncbi.nih.gov/Genbank>) and were analyzed for secondary DNA structure by using M-Fold (<http://frontend.bioinfo.rpi.edu/applications/mfold/cgi-bin/dna-form1.cgi>). Primers were designed by using Primer 3 software (<http://frodo.wi.mit.edu/primer3/>) with the maximum self-complementarity score set at 5 and the maximum 3 self-complementarity score set to 0 to minimize primer-dimer formation (Table 1). Primer sequences for canine *Bcl-2*, *XIAP*, and *β -actin* have been previously described (39). Quantitative real time PCR (qRT-PCR) was done by using SYBR Green (Fermentas). Samples were run in triplicate by using standard conditions on an ABI 7500 sequence detector (Applied Biosystems) and data were analyzed by using *β -actin* as an endogenous control. Dissociation curves were done after each experiment to confirm the specificity of product amplification.

Immunohistochemistry

Immunohistochemical analysis was done on formalin fixed, paraffin-embedded tissue sections of lymph nodes from healthy dogs and dogs with DLBCL. Sections were deparaffinized in xylene, rehydrated, and boiled in sodium citrate buffer to unmask antigens. Endogenous peroxidase was blocked with 3% hydrogen peroxide. Immunohistochemistry was done by using a rabbit anti-human p65 antibody (C22B4; Cell Signaling) or rabbit isotype control

Table 1. Primer sequences for NF- κ B target gene expression analyses

Gene	RT-PCR primers		Accession #
<i>A1</i>	Sense:	5'-TCA ATC AGG TGA TGG AGA AGG-3'	XM_545888
	Antisense:	5'-TGC TCG AGG AGT TTC TTG GT-3'	
<i>Actin</i>	Sense:	5'-TCC CTG GAG AAG AGC TAC GA-3'	AF021873
	Antisense:	5'-CTT CTG CAT CCT GTC AGC AA-3'	
<i>Bcl-2</i>	Sense:	5'-TGG ATG ACT GAG TAG CTG AA-3'	NM_001002949
	Antisense:	5'-GGC CTA CTG ACT TCA CTT AT-3'	
<i>c-FLIP</i>	Sense:	5'-TCC AGG AAT CAG GAC CAT TT-3'	XM_545592
	Antisense:	5'-GAT TCC TAG GGG CTT GCT CT-3'	
<i>Cyclin D1</i>	Sense:	5'-CAT CTA CAC TGA CAA CTC CAT CC-3'	NM_001005757
	Antisense:	5'-CAG GTT CCA CTT CAG TTT GTT C-3'	
<i>IκBα</i>	Sense:	5'-CCT ACG TCC AGC CAT CAT TT-3'	XM_537413
	Antisense:	5'-CAG TTC CTC CTT GGG GTT TG-3'	
<i>XIAP</i>	Sense:	5'-ACT ATG TAT CAC TTG AGG CTC TGG TTT C-3'	AY603038
	Antisense:	5'-AGT CTG GCT TGA TTC ATC TTG TGT ATG-3'	

(sc-2027; Santa Cruz Biotechnology) and the Vectastain Elite ABC Kit (Vector Laboratories). Sections were developed by using 3,3'-diaminobenzidine and counterstained with hematoxylin. Slides were viewed by using a Nikon E600 infinity corrected upright microscope. Bright-field images were acquired by using a Nikon Digital Sight DS-Fi1 color camera by using NIS-Element BR3.0 for image analysis.

***In vitro* NBD peptide treatment of malignant B lymphocytes**

Freshly isolated or cryopreserved malignant lymphocytes from lymph nodes of dogs with DLBCL were seeded at 4×10^6 cells/mL and treated with 100 μ m of either NBD or mutant peptide in dimethyl sulfoxide (DMSO). After 12 hours, whole cell extracts were evaluated by immunoblot for canonical NF- κ B activity as described above and for apoptosis by flow cytometry. The human ABC- and GCB-DLBCL cell lines OCI-Ly10 and SUDHL-6 were seeded at 0.5×10^6 cells/mL and treated with 0, 25, or 50 μ mol/L NBD peptide. After 24 hours, cell death was evaluated by Trypan Blue.

Flow cytometry

Detection of apoptotic cell death was done by using the PE Annexin-V Apoptosis Detection Kit (BD Biosciences). Cells were acquired on a FACSCalibur cytometer (BD Biosciences) and analyzed by using TreeStar FlowJo software. Immunophenotyping of primary lymphoma cells was done by using the following antibodies: Fluorescein isothiocyanate (FITC)-conjugated rat anti-canine CD3 (Serotec), adenomatous polyposis coli (APC)-conjugated mouse anti-human CD79a (BD Pharmingen) and mouse anti-canine CD21-like molecule (Serotec). A FITC-labeled goat anti-mouse IgG secondary antibody was used to detect anti-CD21 antibody. For CD79a staining, cells were fixed with 1% paraformaldehyde and then permeabilized with 0.1% saponin before staining.

Pilot trial

All *in vivo* studies were done following the approval from the University of Pennsylvania's Institutional Animal Care and Use Committee and the University of Pennsylvania School of Veterinary Medicine Clinical Review Board. Client-owned dogs were enrolled with the following inclusion criteria: (i) cytologic confirmation of relapsed large B-cell lymphoma, (ii) stage III-V, substage a, with at least 1 set of bilateral, enlarged, measurable, malignant lymph nodes, (iii) life expectancy of greater than 1 month, (iv) no concurrent systemic disease, (v) adequate hematologic, renal, and hepatic function, and (vi) the presence of constitutive NF- κ B activity within malignant lymph nodes as determined by EMSA. All dogs enrolled in the pilot trial had been previously treated with a range of rescue chemotherapeutic agents including L-asparaginase, cyclophosphamide, vincristine, prednisone, lomustine, doxorubicin, mechlorethamine, procarbazine, dacarbazine, and total body irradiation. Signed-informed consent was obtained

from all owners before entry into the study. Baseline evaluation included a complete medical history, full clinical examination, complete blood count (CBC), chemistry screen (CS) and urinalysis, malignant lymph node cytology, and flow cytometric immunophenotyping by using a basic panel of antibodies including CD3, CD21, and CD79a. Eligible dogs received either 5 or 10 mg NBD peptide/100 g malignant lymph node mass. Node mass was calculated based on 2 dimension caliper measurements which were averaged and considered the diameter of the lymph node sphere. Volume was determined based on $V = 2/3\pi r^3$. On the basis of our previous assessments, 1 cm³ of malignant lymph node mass is equivalent to ~379 mg. Peptide was injected directly into 1 malignant node and the same node was biopsied 24 hours later. qRT-PCR for NF- κ B target gene expression was done on node tissue taken 24 hours post-NBD peptide treatment and was normalized to values obtained pre-NBD peptide. No control dogs were included in this pilot study. Following the post-treatment biopsy, patients received rescue chemotherapy as determined by their primary attending oncologist. All dogs received a follow-up full clinical examination plus CBC, CS, and urinalysis 1-week post-NBD peptide injection. Mass of the NBD-treated lymph node and the contralateral, malignant, untreated lymph node was calculated before, and 1-week post-NBD peptide administration.

Results

Constitutive canonical NF- κ B activity occurs in dogs with DLBCL

To determine whether constitutive NF- κ B activity is present in dogs with spontaneous, histologically confirmed DLBCL, EMSA was done on biopsy samples taken from the peripheral lymph nodes of 13 dogs with DLBCL at the time of diagnosis and prior to chemotherapy. Constitutive NF- κ B activity was detected in all 13 samples. In contrast, NF- κ B activity was not detected in normal lymph nodes from healthy dogs (Fig. 1A). To determine whether constitutive NF- κ B activity was associated with the canonical pathway, malignant lymph node tissue from 11 dogs with newly diagnosed, histologically confirmed DLBCL was analyzed by EMSA supershift by using antibodies directed against p50, p65, and c-Rel. In 7 dogs, p50, p65, and c-Rel were all present in the active NF- κ B complexes (Fig. 1B). In 2 dogs, only p50 and p65 were present, and in 2 dogs, only p50 was identified (data not shown). To further confirm the presence of constitutive canonical NF- κ B activity, formalin-fixed, paraffin-embedded lymph node tissues from 6 dogs with DLBCL were evaluated for the presence of nuclear p65 by immunohistochemistry (Fig. 1C). All tissue sections had abundant cytoplasmic and nuclear-localized p65 whereas 4 healthy canine lymph node sections had minimal p65 staining.

To determine whether the detected canonical NF- κ B activity was associated with constitutive IKK activity, the phosphorylation status of the IKK complex and the inhibitory I κ B α protein were evaluated by immunoblot




Figure 1. Constitutive canonical NF- κ B activity in dogs with spontaneous DLBCL. **A**, whole cell protein extracts of normal lymph nodes (NLN) taken from 3 healthy dogs and lymph nodes taken from 4 dogs with primary, untreated DLBCL were evaluated for NF- κ B DNA-binding activity by EMSA. Results shown are representative of 13 dogs with DLBCL. **B**, whole cell protein extracts of lymph nodes taken from 2 dogs with primary, untreated DLBCL were subject to EMSA supershift analysis with antibodies against p65, p50, c-Rel, or control rabbit IgG. Results shown are representative of 7 dogs with DLBCL. **C**, immunohistochemical staining for p65 was done on lymph node sections from healthy dogs and dogs with DLBCL. Rabbit IgG was used as an isotype control. Arrowheads indicate nuclear p65 staining. Results shown are representative of 6 dogs with DLBCL and 4 healthy dogs (NLN). **D**, whole cell protein extracts of NLN and lymph nodes taken from 3 dogs with primary, untreated DLBCL were evaluated for the presence of p-IKK β and p-I κ B α by Western blot. Results shown are representative of 8 dogs with DLBCL. β -Actin was used as an endogenous loading control.

(Fig. 1D). p-IKK β and p-I κ B α were detected in 8 of 9 lymph node biopsy samples taken from dogs with histologically confirmed, untreated DLBCL with constitutively active NF- κ B as determined by EMSA. In contrast, p-IKK β and p-I κ B α were only detected at very low levels in normal lymph nodes taken from healthy dogs (Fig. 1D). Taken together, these findings show that constitutive canonical NF- κ B activity occurs in the malignant tissue of dogs with DLBCL, and that in the majority of dogs this activity is associated with IKK activation and downstream I κ B α phosphorylation. These data parallel the findings in human ABC-DLBCL and suggest that dogs with DLBCL represent a clinically relevant, spontaneous large animal model in which to evaluate the therapeutic effects of NF- κ B inhibition in the treatment of ABC-DLBCL.

Antiapoptotic NF- κ B target genes are upregulated in canine DLBCL

To determine whether NF- κ B target genes are upregulated in canine DLBCL, malignant lymph node tissue from 14 dogs with histologically confirmed DLBCL, and constitutive NF- κ B activity was evaluated by qRT-PCR for *Bcl-2*, *c-FLIP*, and *XIAP* gene expression (Fig. 2). Tissue samples were obtained at the time of diagnosis and prior to chemotherapy. Using healthy canine lymph node tissue as a normalizing control, statistically significant increases in

Bcl-2 gene expression were identified in 7 dogs. In addition, *c-FLIP* and *XIAP* gene expression was significantly increased in 6 and 7 dogs, respectively. These results suggest that constitutive canonical NF- κ B activity in canine DLBCL promotes the overexpression of antiapoptotic NF- κ B target genes. This finding parallels that in human patients with ABC-DLBCL and indicates that constitutive canonical NF- κ B activity in the dog may contribute to lymphomagenesis and chemoresistance.

NBD peptide inhibits constitutive NF- κ B activity and promotes apoptosis in DLBCL cells *in vitro*

To determine whether NBD peptide can inhibit constitutive phosphorylation of IKK and prevent downstream I κ B α phosphorylation, cells from malignant lymph nodes of dogs with DLBCL were treated *in vitro* with either 100 μ mol/L NBD peptide or 100 μ mol/L mutant peptide for 12 hours and then analyzed for the presence of p-IKK β and p-I κ B α by immunoblot (Fig. 3A and B). Cells treated with NBD peptide showed a marked reduction of p-IKK β compared with untreated cells, and those treated with either mutant peptide or DMSO vehicle control. NBD peptide treatment also resulted in complete inhibition of I κ B α phosphorylation (Fig. 3B). In contrast, treatment with mutant peptide or DMSO (vehicle control) did not inhibit I κ B α phosphorylation.




Figure 2. Expression of antiapoptotic NF- κ B target genes in canine DLBCL. RNA was isolated from lymph node biopsies of dogs with DLBCL at the time of diagnosis and prior to chemotherapy. qRT-PCR was done for *Bcl-2*, *c-FLIP*, and *XIAP*. Values are expressed as relative quantification by using β -actin as an endogenous control. RNA isolated from biopsy samples of 3 healthy canine lymph nodes were used as a normalizing control. All data shown represent means \pm SEMs of reactions done in triplicate. Statistical analysis was done by using a 2-tailed Student's *t* test where *, $P < 0.05$.

To determine the effects of NF- κ B inhibition on DLBCL cell apoptosis, malignant lymphocytes harvested from canine DLBCL tumors were treated *in vitro* with NBD or mutant peptide and analyzed by flow cytometry (Fig. 3C). After 12 hours incubation with NBD peptide, more than 50% of DLBCL cells stained positive for Annexin-V while only 15% of cells treated with mutant peptide appeared apoptotic. Comparable results were identified in malignant cells taken from the lymph nodes of 3 dogs with DLBCL. In contrast, when peripheral blood mononuclear cells (PBMC) isolated from healthy dogs were treated with either 100 μ mol/L NBD or mutant peptide for 12 hours, less than

5% of PBMCs stained positive for Annexin-V (Fig. 3D). Collectively, these data show that NBD peptide inhibits constitutive NF- κ B activity in canine DLBCL cells, leading to selective, rapid apoptosis of malignant cells.

To confirm a selective cytotoxic effect of NBD peptide on human ABC-DLBCL cells, OCI-Ly10 cells and the GCB-DLBCL cell line SUDHL-6 were treated with NBD peptide for 24 hours and the percentage increase in cell death compared with untreated cells was calculated. At 24 hours, NBD peptide induced a dose-dependent increase in cell death of the ABC-DLBCL cell line, OCI-Ly10. In contrast, NBD peptide did not induce cell death in the GCB-DLBCL cell line at either concentration used (Fig. 3E).

***In vivo* administration of NBD peptide inhibits NF- κ B target gene expression and leads to a reduction in tumor burden in dogs with relapsed, chemoresistant large B-cell lymphoma**

To determine whether NBD peptide can inhibit NF- κ B target gene expression in malignant lymphoid tissue *in vivo*, privately owned dogs with relapsed, chemoresistant large B-cell lymphoma were enrolled to a small pilot study (Fig. 4). Eligibility criteria included the presence of histologically or cytologically confirmed large B-cell lymphoma and constitutive NF- κ B activity in malignant lymph nodes as determined by EMSA. Response to NBD peptide was determined by evaluating NF- κ B target gene expression pre- and 24 hours post-NBD peptide administration. Clinical response to NBD peptide was determined by comparing the mass of the injected lymph node before NBD peptide and 1 week after administration of peptide plus rescue chemotherapy. The mass of the contralateral malignant lymph node that was not injected with NBD peptide but was exposed to systemic rescue chemotherapy was also determined at both time points and the change in lymph node mass was calculated for both nodes and compared. Six eligible dogs were enrolled to the study, 3 dogs received 5 mg NBD peptide/100 g of node tissue and 3 received 10 mg NBD peptide/100 g node tissue via a single intranodal injection. At the time of enrollment, qRT-PCR analysis of lymph node tissue revealed that all dogs overexpressed at least 2 of 6 NF- κ B target genes that promote cellular proliferation and inhibit apoptosis when compared with normal lymph node tissue (Fig. 5A left). 24 hours post-NBD peptide administration 1 of 3 dogs receiving 5 mg NBD peptide/100 g nodal tissue showed a reduction in the expression of 4/6 NF- κ B target genes within the malignant node when compared with pre-NBD treatment gene expression values. At the higher dose of NBD peptide, all 3 dogs showed a reduction in the expression of at least 3/6 NF- κ B target genes including *Bcl-2*, *Cyclin D1*, and *I κ B α* when compared with pre-NBD treatment values (Fig. 5A right).

Response to NBD peptide was determined by comparing the change in mass of the NBD peptide-treated lymph node with change in mass of the contralateral, malignant, noninjected lymph node. Results were available for 4 of 6




Figure 3. NBD peptide inhibits constitutive NF- κ B activity in canine primary DLBCL *in vitro*. Primary canine DLBCL cells were treated with either 100 μ mol/L NBD peptide or mutant peptide or with DMSO (vehicle control) for 12 hours. 25 μ g of whole cell extract was immunoblotted by using antibodies against (A) p-IKK β and pan-IKK β and (B) p-I κ B α and pan-I κ B α . β -Actin was used as an endogenous loading control. Results are representative of 3 individual dogs. Canine PBMCs with or without TNF- α were included as positive and negative controls for p-IKK β and p-I κ B α . C, primary canine DLBCL cells were either untreated (grey histogram) or treated (open histograms) with either 100 μ mol/L NBD peptide or mutant peptide for 12 hours and then analyzed by flow cytometry for Annexin-V. Results are representative of 3 individual dogs. D, healthy canine PBMCs were treated with 100 μ mol/L NBD peptide or mutant peptide for 12 hours and then analyzed by flow cytometry for Annexin-V. Histograms are gated on CD21 $^{+}$ lymphocytes. E, OCI-Ly10 and SUDHL-6 cells were treated with 0, 25, or 50 μ mol/L NBD peptide for 24 hours. Cell death was quantified by Trypan Blue. Data represent the percent increase in cell death in treated wells compared with untreated cells. Cell counts were done in triplicate, and data shown represents means \pm SEMs. Statistical analysis was done by using a 2-tailed Student's *t* test. *, *P* < 0.05. Results are representative of 3 separate experiments.

treated dogs (Fig. 5B). Rescue chemotherapy protocols administered to each dog after NBD peptide treatment varied among dogs and were determined by the referring oncologist. In 3 of 4 dogs, the NBD peptide-treated lymph node showed between a 1.5- and 6-fold reduction in mass when compared with the contralateral, malignant lymph node that did not receive NBD peptide. Two dogs (dogs 7 and 8) were treated on a compassionate basis and given intranodal injections of 10 mg NBD/100 g tumor into 8 peripheral enlarged lymph nodes, resulting in cumulative NBD doses of 0.33 and 0.8 mg/kg body weight, respectively. This amendment to the original protocol was approved by the University of Pennsylvania's Institutional Animal Care and Use Committee. At enrollment, each dog overexpressed at least 3 NF- κ B target genes (Fig. 5A left). 24 hours after NBD peptide administration, reductions in NF- κ B target gene expression were observed in 3/6 and 6/6 evaluated genes, respectively (Fig. 5A right). In these 2 dogs, a reduction in lymph node mass was observed in 3/8 and 4/8 NBD-

treated lymph nodes (Fig. 5C). Despite NBD peptide and systemic chemotherapy, the remaining lymph nodes continued to increase in size. In both dogs, the lymph nodes that showed a reduction in mass following NBD peptide treatment were the smallest nodes at the time of treatment.

Discussion

Aberrant, constitutive, canonical NF- κ B activity is recognized as a key contributor to oncogenesis and chemotherapy resistance in many different cancers and as such represents a logical target for inhibition in cancer therapy. Although nearly 800 inhibitors of NF- κ B have been identified (40), their ability to inhibit constitutive NF- κ B activity in spontaneous cancers *in vivo* and the therapeutic relevance of such inhibition in patients with cancer is unknown.

In this study, we first set out to determine whether dogs with DLBCL represent an appropriate, spontaneous, large




Figure 4. Schematic of pilot clinical trial.

animal model in which to evaluate the therapeutic effects of NF- κ B inhibition. DLBCL in the dog is often characterized by total effacement of lymph node architecture with large malignant B cells that have a high-mitotic index and multiple, prominent nucleoli (41). This malignancy in dogs is frequently associated with overexpression of c-myc that occurs as a result of chromosomal translocations that place the oncogene under control of the IgH promoter

(17). Although these histocytologic and cytogenetic changes are also found in ABC-DLBCL in humans, it is unknown whether canine DLBCL is characterized by constitutive canonical NF- κ B activity. Here, we show that dogs with histologically confirmed DLBCL have constitutive canonical NF- κ B signaling in malignant lymphoid tissue at the time of diagnosis. In most cases evaluated, p50/p65 and p50/c-Rel heterodimers were present in the activated complex. Only 2 biopsy samples had evidence of active p50 in the apparent absence of active p65 or c-Rel suggesting the presence of p50 homodimers or complexes with RelB in these patients. These differences may be attributed to the specific mechanism(s) responsible for constitutive NF- κ B activity in each tumor sample, as the identity and relative abundance of active homo- and heterodimers depends on the nature of the activating signal or pathway aberrancy (42, 43). Furthermore, as in human ABC-DLBCL patients, constitutive activity was associated with phosphorylation of IKK β and I κ B α in all but 1 dog evaluated, indicating that in the majority of cases signals leading to constitutive activity occur upstream of the IKK complex (7, 10). These findings confirm on a molecular level that human ABC-DLBCL and canine DLBCL have similar aberrant NF- κ B




Figure 5. *In vivo* administration of NBD peptide inhibits NF- κ B target gene expression in malignant lymph nodes of dogs with relapsed/resistant large B-cell lymphoma. A; left, qRT-PCR of malignant lymph nodes of dogs with relapsed large B-cell lymphoma at enrollment. Values are expressed as relative quantification by using β -actin as an endogenous control. RNA isolated from biopsy samples of 3 healthy canine lymph nodes were used as a normalizing control. A; right, qRT-PCR of malignant lymph nodes 24 hours post-NBD peptide injection. RNA isolated from biopsy samples of the same dogs prior to NBD peptide treatment was used as a normalizing control. All data shown represents means \pm SEMs of reactions done in triplicate. B, comparison of percent decrease in mass between NBD-injected lymph node and contralateral, malignant, noninjected lymph node of 4 study dogs. Measurements were taken immediately before 1-week post-NBD peptide administration. C, percent decrease in mass of NBD-injected lymph nodes from 2 dogs that received multinodal injections of NBD peptide.

signaling and that the dog may serve as a clinically relevant large animal model in which to study the therapeutic effects of NF- κ B pathway inhibition in ABC-DLBCL.

Although constitutive NF- κ B activity was present in the malignant lymph nodes of all dogs with untreated DLBCL, the NF- κ B target genes *Bcl-2*, *c-FLIP*, and *XIAP* were overexpressed in only half of the biopsy samples. These findings serve to underline the multiple layers of complexity involved in the regulation of NF- κ B target gene expression. First, the particular combination of activated NF- κ B family members influences the target gene expression profile, in part because these family members have different binding affinities for different gene promoters. Although NF- κ B activation originates upstream of IKK in these dogs, the specific activation triggers have not been investigated and may differ between dogs, resulting in different NF- κ B family member usage and different target gene expression signatures. However, in this small dataset, we did not find any correlation between the expression of NF- κ B target genes and activation of specific NF- κ B family members. Second, other signaling pathways that activate additional transcription factors such as AP-1 and ETS may be concurrently active in these tumors. These transcription factors together with cofactors that modify NF- κ B activity influence gene activation and repression, leading to variable gene expression profiles (44, 45). Taken together these points highlight the complex and multifaceted nature of NF- κ B regulated gene expression. A much larger sample set of histologically confirmed, DLBCL biopsies with complete gene expression profiling together with a thorough evaluation of activated NF- κ B family members will be needed to identify specific NF- κ B gene expression signatures.

In the second part of this study, we show that NBD peptide can inhibit constitutive IKK β and I κ B α phosphorylation in DLBCL biopsy samples which leads to increased apoptosis of malignant lymphocytes. Furthermore, in a small clinical pilot study, NBD peptide showed a dose-dependent inhibition of NF- κ B target gene expression in dogs with relapsed, refractory large B-cell lymphoma. In 2 dogs receiving the lower NBD peptide dose (dogs 1 and 3), all NF- κ B target genes increased up to 4-fold 24 hours post-NBD peptide suggesting that constitutive NF- κ B activity in these dogs may be exerting a transcriptional repressive effect. No correlation was identified between 24-hour gene expression profile results and change in lymph node mass 7 days later.

In 3 of 4 dogs, intranodal administration of NBD peptide led to a marked reduction in tumor mass when compared with the contralateral, malignant lymph node that did not receive NBD peptide, although it is unclear whether NBD peptide alone, or acting synergistically with rescue chemotherapy exerted this effect. In dogs that received multi-node injections of NBD peptide, the apparent effects on tumor mass were variable. Although most nodes continued to increase in size despite NBD treatment and rescue chemotherapy, several nodes in both dogs showed a marked reduction in tumor mass. In both cases, responding nodes were smaller than nonresponsive nodes at the time

of treatment. These results raise the possibility that the NBD peptide dose and its ability to diffuse within malignant nodes may influence its ability to exert a measurable therapeutic effect.

Given the central role of the IKK complex in canonical NF- κ B activation, IKK inhibitors are being investigated as novel therapeutic agents in both solid and hematologic malignancies that depend upon constitutive NF- κ B activity for their survival. IKK β inhibitors have been developed to specifically block canonical NF- κ B signaling; however, in the presence of selective IKK β inhibition, compensatory IKK α activity can drive NEMO-dependent canonical NF- κ B activity in human DLBCL cells (46). This finding suggests that inhibitors of both IKK α and IKK β activity (such as NBD peptide) will provide more complete inhibition and have greater therapeutic efficacy than IKK β inhibitors alone. Furthermore, agents such as NBD peptide that specifically target the scaffolding molecule NEMO are less likely than kinase inhibitors to exhibit off-target effects on other intracellular signaling pathways (27).

Previous studies evaluating the safety and efficacy of NBD peptide as an antiinflammatory agent *in vivo* have shown that both short- and long-term administration of NBD peptide to piglets and rodents at doses ranging from 0.75 to 20 mg/kg is well tolerated (24, 28–31). In this small pilot study, NBD peptide did not cause any systemic toxicity and there were no significant changes in serum chemistries or hematologic profiles over the 10-day assessment period that could be attributable to NBD peptide administration. However, the dose of NBD peptide administered on a mg/kg body weight basis was very low. However, 2 dogs treated with the higher dose of NBD peptide developed submandibular abscesses within 1 week of NBD injection into prescapular lymph nodes. In both cases, the commensal skin bacteria *Staphylococcus aureus* was cultured and successfully treated with drainage and broad-spectrum antibiotics. In these cases, it is possible that commensal skin bacteria introduced with NBD peptide were able to colonize the necrotic node and that inhibition of TLR signaling by NBD peptide may have contributed to this process.

In this pilot study, NBD peptide was administered intranodally rather than systemically to first determine whether it could inhibit constitutive NF- κ B activity in malignant lymphocytes *in vivo*. On the basis of the favorable *in vitro* and *in vivo* results, we are now actively recruiting dogs with ABC-DLBCL to a phase I dose escalation clinical trial in which NBD peptide will be administered systemically via the intravenous route. Although the amount of peptide necessary to achieve complete inhibition of NF- κ B activity in a dog or human are likely to be high, it is unknown whether complete inhibition is required to see a therapeutic effect when used in combination with chemotherapy.

Taken together, the results reported here provide the first *in vivo* evidence that NBD peptide can inhibit antiapoptotic target gene expression driven by constitutive NF- κ B activity and that this inhibition, either alone or in combination with cytotoxic therapy, can lead to improved clinical

responses in dogs with ABC-like DLBCL. Given our reported findings that show the dog is a highly relevant, spontaneous clinical model for ABC-DLBCL in humans, the therapeutic effects of NF- κ B inhibition seen in the canine cancer patient are likely to hold high translational relevance to human patients with ABC-DLBCL.

Disclosure of Potential Conflicts of Interest

M.J. May holds a patent for the NBD peptide and A.S. Baldwin is the founder of TheraLogics and therefore has commercial interests. All other authors declare no competing financial interests.

Grant Support

American Cancer Society ACS-IRG (N.J. Mason), American Kennel Club Canine Health Foundation (N.J. Mason), the Mari Lowe Center for Comparative Oncology (N.J. Mason), and the Canine Hereditary Cancer Consortium/NIH UC2 CA148149 (N.J. Mason). A. Gaurmier-Hausser is supported by the Ruth L. Kirschstein grant/NIH T32 CA 09140.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received December 20, 2010; revised May 2, 2011; accepted May 18, 2011; published OnlineFirst May 24, 2011.

References

- Abramson JS, Shipp MA. Advances in the biology and therapy of diffuse large B-cell lymphoma: moving toward a molecularly targeted approach. *Blood* 2005;106:1164–74.
- Alizadeh AA, Eisen MB, Davis RE, Ma C, Lossos IS, Rosenwald A, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 2000;403:503–11.
- Lam LT, Davis RE, Pierce J, Hepperle M, Xu Y, Hottel M, et al. Small molecule inhibitors of I κ B kinase are selectively toxic for subgroups of diffuse large B-cell lymphoma defined by gene expression profiling. *Clin Cancer Res* 2005;11:28–40.
- Nyman H, Adde M, Karjalainen-Lindsberg ML, Taskinen M, Berglund M, Amini RM, et al. Prognostic impact of immunohistochemically defined germinal center phenotype in diffuse large B-cell lymphoma patients treated with immunochemotherapy. *Blood* 2007;109:4930–5.
- Davis RE, Ngo VN, Lenz G, Tolar P, Young RM, Romesser PB, et al. Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. *Nature* 2010;463:88–92.
- Feuerhake F, Kutok JL, Monti S, Chen W, LaCasce AS, Cattoretto G, et al. NF κ B activity, function, and target-gene signatures in primary mediastinal large B-cell lymphoma and diffuse large B-cell lymphoma subtypes. *Blood* 2005;106:1392–9.
- Davis RE, Brown KD, Siebenlist U, Staudt LM. Constitutive nuclear factor κ B activity is required for survival of activated B cell-like diffuse large B cell lymphoma cells. *J Exp Med* 2001;194:1861–74.
- Karin M, Lin A. NF- κ B at the crossroads of life and death. *Nat Immunol* 2002;3:221–7.
- Gilmore TD. Introduction to NF- κ B: players, pathways, perspectives. *Oncogene* 2006;25:6680–4.
- Compagno M, Lim WK, Grunn A, Nandula SV, Brahmachary M, Shen Q, et al. Mutations of multiple genes cause deregulation of NF- κ B in diffuse large B-cell lymphoma. *Nature* 2009;459:717–21.
- Lenz G, Davis RE, Ngo VN, Lam L, George TC, Wright GW, et al. Oncogenic CARD11 mutations in human diffuse large B cell lymphoma. *Science* 2008;319:1676–9.
- Ferch U, Kloo B, Gewies A, Pfander V, Duwel M, Peschel C, et al. Inhibition of MALT1 protease activity is selectively toxic for activated B cell-like diffuse large B cell lymphoma cells. *J Exp Med* 2009;206:2313–20.
- Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, et al. Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature* 2005;438:803–19.
- Acland GM, Aguirre GD, Bennett J, Aleman TS, Cideciyan AV, Bennicelli J, et al. Long-term restoration of rod and cone vision by single dose rAAV-mediated gene transfer to the retina in a canine model of childhood blindness. *Mol Ther* 2005;12:1072–82.
- Herzog RW, Yang EY, Couto LB, Hagstrom JN, Elwell D, Fields PA, et al. Long-term correction of canine hemophilia B by gene transfer of blood coagulation factor IX mediated by adeno-associated viral vector. *Nat Med* 1999;5:56–63.
- Ponder KP, Melniczek JR, Xu L, Weil MA, O'Malley TM, O'Donnell PA, et al. Therapeutic neonatal hepatic gene therapy in mucopolysaccharidosis VII dogs. *Proc Natl Acad Sci U S A* 2002;99:13102–7.
- Breen M, Modiano JF. Evolutionarily conserved cytogenetic changes in hematological malignancies of dogs and humans—man and his best friend share more than companionship. *Chromosome Res* 2008;16:145–54.
- Paoloni M, Khanna C. Translation of new cancer treatments from pet dogs to humans. *Nat Rev Cancer* 2008;8:147–56.
- Paoloni MC, Khanna C. Comparative oncology today. *Vet Clin North Am Small Anim Pract* 2007;37:1023–32.
- MacEwen EG. Spontaneous tumors in dogs and cats: models for the study of cancer biology and treatment. *Cancer Metastasis Rev* 1990;9:125–36.
- Fournel-Fleury C, Magnol JP, Bricaire P, Marchal T, Chabanne L, Delverdier A, et al. Cytohistological and immunological classification of canine malignant lymphomas: comparison with human non-Hodgkin's lymphomas. *J Comp Pathol* 1997;117:35–59.
- Thomas R, Smith KC, Ostrander EA, Galibert F, Breen M. Chromosome aberrations in canine multicentric lymphomas detected with comparative genomic hybridisation and a panel of single locus probes. *Br J Cancer* 2003;89:1530–7.
- Vail DM, ME, Young KM. Canine lymphoma and lymphoid leukemia. In: Withrow SJ, editor. *Small Animal Clinical Oncology*. Philadelphia: WB Saunders Co.; 2001, p. 558–90.
- May MJ, D'Acquisto F, Madge LA, Glockner J, Pober JS, Ghosh S. Selective inhibition of NF- κ B activation by a peptide that blocks the interaction of NEMO with the I κ B kinase complex. *Science* 2000;289:1550–4.
- May MJ, Marienfeld RB, Ghosh S. Characterization of the I κ B B-kinase NEMO binding domain. *J Biol Chem* 2002;277:45992–6000.
- Solt LA, Madge LA, May MJ. NEMO-binding domains of both IKK α and IKK β regulate I κ B kinase complex assembly and classical NF- κ B activation. *J Biol Chem* 2009;284:27596–608.
- Orange JS, May MJ. Cell penetrating peptide inhibitors of Nuclear Factor- κ B. *Cell Mol Life Sci* 2008;65:3564–91.
- Ankermann T, Reissner A, Wiemann A, Krams M, Kohler H, Krause MF. Topical inhibition of nuclear factor- κ B enhances reduction in lung edema by surfactant in a piglet model of airway lavage. *Crit Care Med* 2005;33:1384–91.
- di Meglio P, Ianaro A, Ghosh S. Amelioration of acute inflammation by systemic administration of a cell-permeable peptide inhibitor of NF- κ B activation. *Arthritis Rheum* 2005;52:951–8.
- Jimi E, Aoki K, Saito H, D'Acquisto F, May MJ, Nakamura I, et al. Selective inhibition of NF- κ B blocks osteoclastogenesis and prevents inflammatory bone destruction in vivo. *Nat Med* 2004;10:617–24.
- Lawrence T, Bebie M, Liu GY, Nizet V, Karin M. IKK α limits macrophage NF- κ B activation and contributes to the resolution of inflammation. *Nature* 2005;434:1138–43.
- Acharya S, Villalta SA, Bakkar N, Bupha-Intr T, Janssen PM, Carathers M, et al. Interplay of IKK/NF- κ B signaling in macrophages and myofibers promotes muscle degeneration in Duchenne muscular dystrophy. *J Clin Invest* 2007;117:889–901.

33. Aggarwal S, Takada Y, Singh S, Myers JN, Aggarwal BB. Inhibition of growth and survival of human head and neck squamous cell carcinoma cells by curcumin via modulation of nuclear factor-kappaB signaling. *Int J Cancer* 2004;111:679–92.
34. Kiessling MK, Klemke CD, Kaminski MM, Galani IE, Krammer PH, Gulow K. Inhibition of constitutively activated nuclear factor-kappaB induces reactive oxygen species- and iron-dependent cell death in cutaneous T-cell lymphoma. *Cancer Res* 2009;69:2365–74.
35. Shishodia S, Amin HM, Lai R, Aggarwal BB. Curcumin (diferuloylmethane) inhibits constitutive NF-kappaB activation, induces G1/S arrest, suppresses proliferation, and induces apoptosis in mantle cell lymphoma. *Biochem Pharmacol* 2005;70:700–13.
36. Thomas RP, Farrow BJ, Kim S, May MJ, Hellmich MR, Evers BM. Selective targeting of the nuclear factor-kappaB pathway enhances tumor necrosis factor-related apoptosis-inducing ligand-mediated pancreatic cancer cell death. *Surgery* 2002;132:127–34.
37. Ianaro A, Tersigni M, Belardo G, Di Martino S, Napolitano M, Palmieri G, et al. NEMO-binding domain peptide inhibits proliferation of human melanoma cells. *Cancer Lett* 2009;274:331–6.
38. Tapia MA, Gonzalez-Navarrete I, Dalmases A, Bosch M, Rodriguez-Fanjul V, Rolfe M, et al. Inhibition of the canonical IKK/NF kappa B pathway sensitizes human cancer cells to doxorubicin. *Cell Cycle* 2007;6:2284–92.
39. Spee B, Arends B, van den Ingh TS, Penning LC, Rothuizen J. Copper metabolism and oxidative stress in chronic inflammatory and cholestatic liver diseases in dogs. *J Vet Intern Med* 2006;20:1085–92.
40. Gilmore TD, Herscovitch M. Inhibitors of NF-kappaB signaling: 785 and counting. *Oncogene* 2006;25:6887–99.
41. Valli V. *Veterinary comparative hematopathology*. Iowa: Blackwell Publishing Ltd.; 2007.
42. Hoffmann A, Levchenko A, Scott ML, Baltimore D. The IkappaB-NF-kappaB signaling module: temporal control and selective gene activation. *Science* 2002;298:1241–5.
43. O’Dea E, Hoffmann A. The regulatory logic of the NF-kappaB signaling system. *Cold Spring Harb Perspect Biol* 2010;2:a000216.
44. Perkins ND. Post-translational modifications regulating the activity and function of the nuclear factor kappa B pathway. *Oncogene* 2006;25:6717–30.
45. Perkins ND. Integrating cell-signalling pathways with NF-kappaB and IKK function. *Nat Rev Mol Cell Biol* 2007;8:49–62.
46. Lam LT, Davis RE, Ngo VN, Lenz G, Wright G, Xu W, et al. Compensatory IKKalpha activation of classical NF-kappaB signaling during IKKbeta inhibition identified by an RNA interference sensitization screen. *Proc Natl Acad Sci U S A* 2008;105:20798–803.

Clinical Cancer Research

NEMO-Binding Domain Peptide Inhibits Constitutive NF- κ B Activity and Reduces Tumor Burden in a Canine Model of Relapsed, Refractory Diffuse Large B-Cell Lymphoma

Anita Gurnier-Hausser, Reema Patel, Albert S. Baldwin, et al.

Clin Cancer Res 2011;17:4661-4671. Published OnlineFirst May 24, 2011.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-10-3310](https://doi.org/10.1158/1078-0432.CCR-10-3310)

Cited articles This article cites 44 articles, 15 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/17/14/4661.full#ref-list-1>

Citing articles This article has been cited by 5 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/17/14/4661.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/17/14/4661>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.