

MicroRNA-125a-5p Is an Independent Prognostic Factor in Gastric Cancer and Inhibits the Proliferation of Human Gastric Cancer Cells in Combination with Trastuzumab

Naohiro Nishida^{1,2}, Koshi Mimori¹, Muller Fabbri³, Takehiko Yokobori¹, Tomoya Sudo¹, Fumiaki Tanaka¹, Kohei Shibata¹, Hideshi Ishii^{1,2}, Yuichiro Doki², and Masaki Mori^{1,2}

Abstract

Purpose: MicroRNA 125a-5p (*miR-125a-5p*) has been reported to be a tumor suppressor in malignancies of the breast, ovary, lung, and central nervous system. However, the clinical significance of *miR-125a-5p* in human gastrointestinal cancer has not been explored. We investigated a tumor inhibitory effect of *miR-125a-5p* in gastric cancer, focusing in particular on the *miR-125a-ERBB2* (*HER2*, *HER-2/neu*) pathway.

Experimental Design: Quantitative RT-PCR was used to evaluate *miR-125a-5p* expression in 87 gastric cancer cases to determine the clinicopathologic significance of *miR-125a-5p* expression. The regulation of *ERBB2* by *miR-125a-5p* was examined with precursor *miR-125a*-transfected cells. Furthermore, we investigated whether *miR-125a-5p* suppresses proliferation of gastric cancer cells in combination with trastuzumab, a monoclonal antibody against *ERBB2*.

Results: Low expression levels of *miR-125a-5p* were associated with enhanced malignant potential such as tumor size ($P = 0.0068$), tumor invasion ($P = 0.031$), liver metastasis ($P = 0.029$), and poor prognosis ($P = 0.0069$). Multivariate analysis indicated that low *miR-125a-5p* expression was an independent prognostic factor for survival. *In vitro* assays showed that *ERBB2* is a direct target of *miR-125a-5p*, which potently suppressed the proliferation of gastric cancer cells, and, interestingly, the growth inhibitory effect was enhanced in combination with trastuzumab.

Conclusions: *miR-125a-5p* is a meaningful prognostic marker. Furthermore, *miR-125a-5p* mimic alone or in combination with trastuzumab could be a novel therapeutic approach against gastric cancer. *Clin Cancer Res*; 17(9); 2725–33. ©2011 AACR.

Introduction

MicroRNAs (miRNA) constitute a class of small (19–25 nucleotides) noncoding RNAs that function as posttranscriptional gene regulators. miRNAs regulate gene expression by binding to their mRNAs (1). Alterations in miRNA expression are involved in the initiation, progression, and metastasis of human cancer, and it is believed that miRNAs function both as tumor suppressors and oncogenes in cancer development (2, 3).

Authors' Affiliations: ¹Department of Surgery and Molecular Oncology, Medical Institute of Bioregulation, Kyushu University, Beppu, Oita; ²Department of Gastroenterological Surgery, Osaka University Graduate School of Medicine, Suita City, Osaka, Japan; and ³Department of Molecular Virology, Immunology and Medical Genetics, The Ohio State University Comprehensive Cancer Center, Columbus, Ohio

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Corresponding Author: Masaki Mori, Department of Gastroenterological Surgery, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita City, Osaka 565-0871, Japan. Phone: 8166-879-3251; Fax: 8166-879-3259; E-mail: mmori@gesurg.med.osaka-u.ac.jp

doi: 10.1158/1078-0432.CCR-10-2132

©2011 American Association for Cancer Research.

Recent studies have shown that the expression of *miR125a-5p* is downregulated in several human cancers such as breast cancer (4–6), ovarian cancer (7), lung cancer (8), and medulloblastoma (9). Li and colleagues reported that a germline mutation in mature miRNA 125a-5p (*miR-125a-5p*) is closely associated with breast cancer tumorigenesis (5). Other reports showed that epidermal growth factor (EGF) receptor signaling suppresses *miR-125a-5p* expression and leads to cancer metastasis in lung cancer (8) and ovarian cancer (10). Furthermore, in squamous cell carcinoma of the oral cavity, the levels of *miR-125a-5p* were significantly downregulated in the saliva of patients (11). These findings strongly suggest that the function of *miR-125a-5p* as a tumor suppressor is not organ specific.

Scott and colleagues revealed that *miR-125a-5p* and its homologue, *miR-125b*, regulate *ERBB2* and *ERBB3* in human breast cancer cells (12). In gastric cancer, *ERBB2* overexpression has been increasingly recognized as a frequent molecular abnormality and as an important therapeutic target similar to breast cancer (13, 14). Preclinical and clinical data have revealed significant efficacy of anti-*ERBB2* therapies, especially trastuzumab (Herceptin), a monoclonal antibody directed at *ERBB2* in gastric cancer (15, 16).

Translational Relevance

Quantitative real-time reverse transcriptase PCR analysis of microRNA 125a-5p (*miR-125a-5p*) in 87 cases of gastric cancer revealed that low expression levels of *miR-125a-5p* were associated with enhanced malignant potential such as tumor size, tumor invasion, liver metastasis, and poor prognosis. To evaluate the function of *miR-125a-5p*, we focused on the *miR-125a-ERBB2* (*HER2*, *HER-2/neu*) pathway. In gastric cancer, *ERBB2* overexpression has been increasingly recognized as an important therapeutic target similar to that in breast cancer. Our data suggested that *miR-125a-5p* directly targets *ERBB2*. *miR-125a-5p* powerfully suppressed the proliferation of gastric cancer cells and, moreover, the growth inhibitory effect was enhanced in combination with trastuzumab, a monoclonal antibody against *ERBB2*. *miR-125a-5p* is a meaningful prognostic indicator. Furthermore, *miR-125a-5p* mimic alone or in combination with trastuzumab could be a novel therapeutic approach against gastric cancer.

In this study, we showed that *miR-125a-5p* functions as a crucial tumor suppressor in human gastric cancer. Low *miR-125a-5p* expression was correlated with more aggressive disease and poorer prognosis and was an independent prognostic factor. Of the numerous target genes of *miR-125a-5p*, we focused on *ERBB2* and discovered that *miR-125a-5p* regulates *ERBB2* in human gastric cancer cells. *miR-125a-5p* potently suppressed the proliferation of gastric cancer cells. Moreover, the growth inhibitory effect was enhanced in combination with trastuzumab. This is the first report describing the clinical significance of *miR-125a-5p* and its growth inhibitory effect in human gastric cancer.

Materials and Methods

Clinical cases

Patients and sample collection. Eighty-seven gastric cancer samples were obtained during surgery and used after obtaining informed consent. All patients underwent resection of the primary tumor at Kyushu University Hospital at Beppu and affiliated hospitals between 1992 and 2000. Written informed consent was obtained from all patients. All patients had a clear histologic diagnosis of gastric cancer, based on the clinicopathologic criteria described by the Japanese Gastric Cancer Association (17). All patients were closely followed every 3 months. The follow-up periods ranged from 0.2 months to 12.3 years, with a mean of 2.6 years. Resected cancer tissues were immediately cut and embedded in Tissue-Tek OCT (optimum cutting temperature) medium (Sakura), frozen in liquid nitrogen, and kept at -80°C until RNA and DNA extraction. Frozen tissue specimens were homogenized in guanidium thiocyanate, and total RNA was obtained by ultracentrifugation through a cesium chlor-

ide cushion. cDNA was synthesized from 8.0 μg of total RNA as previously described (18). Clinicopathologic factors and clinical stage were classified by the criteria of the Japanese Gastric Cancer Association (17). All sample data, including age, gender, tumor size and depth, lymphatic invasion, lymph node metastasis, vascular invasion, liver metastasis, peritoneal dissemination, distant metastasis, clinical stage, and histologic grade, were obtained from the clinical and pathologic records and are summarized in Table 1.

Evaluation of *miR-125a-5p* expression in clinical samples

For *miR-125a-5p* quantitative real-time reverse transcriptase PCR (RT-PCR), cDNA was synthesized from 10 ng of total RNA using TaqMan MicroRNA hsa-*miR-125a-5p* specific primers (Applied Biosystems) and a TaqMan MicroRNA Reverse Transcription kit (Applied Biosystems). RT-PCR protocols are described in Supplementary Data.

Evaluation of *ERBB2*, *DACH1*, and *PDCD6* mRNA expression in gastric cancer cells

For RNA analysis, each cell line was seeded at 2×10^5 cells per well in a volume of 2 mL in 6-well flat-bottomed microtiter plates. Total RNA from cell lines was isolated using the mirVana miRNA Isolation Kit (Ambion) after 48-hour incubation. Quantitative RT-PCR was carried out to measure *ERBB2*, dachshund homolog 1 (*DACH1*), and programmed cell death 6 (*PDCD6*) mRNA expression with the Universal Probe Library Probe (UPL; Roche Diagnostics). Primer sequences corresponding to UPL and RT-PCR protocols are described in Supplementary Data.

Immunohistochemistry

Immunohistochemical studies of *ERBB2* were conducted on formalin-fixed, paraffin-embedded (FFPE) surgical sections obtained from patients with gastric cancer. Tissue sections were deparaffinized, soaked in 0.01 mol/L sodium citrate buffer, and boiled in a microwave oven for 5 minutes at 500 W to retrieve cell antigens. Mouse monoclonal antibody against *ERBB2* (Epitomics, Inc.) diluted 1:400 was used as the primary antibody. All tissue sections were immunohistochemically stained with the avidin-biotin-peroxidase method (LSAB+ System HRP; Dako, Inc.) and counterstained with hematoxylin.

Evaluation of *ERBB2* immunohistochemical staining

The slides were examined and scored independently by 2 experienced pathologists. Evaluation of the results was done according to the criteria recommended by Hofmann and colleagues and other groups (19, 20), by assigning a score from 0 to 3+. Scores were defined as follows: 0, no reactivity or membranous reactivity in less than 10% of \geq cells; 1+, faintly perceptible membranous reactivity in 10% or more of cells; cells are reactive only in part of their membrane; 2+, weak to moderate complete or basolateral membranous reactivity in \geq 10% or more of tumor cells; 3+, moderate to strong complete or basolateral

Table 1. *miR-125a-5p* expression and clinicopathologic factors

Factors	Low expression group (n = 55)		High expression group (n = 32)		P
	n	%	n	%	
Age (mean ± SD)	63.5 ± 1.60		67.2 ± 2.09		
Sex					
Male	36	65.5	20	62.5	0.78
Female	19	34.5	12	37.5	
Histologic grade					
Well and moderately	21	38.2	17	53.1	0.18
Poorly and others	34	61.8	15	46.9	
Size					
50 mm > (small)	18	32.7	20	62.5	0.0068 ^a
50 mm < (large)	37	67.3	12	37.5	
Depth of tumor invasion ^b					
m, sm, mp	12	21.8	21	14	0.031 ^a
ss, se, si	43	78.2	11	18	
Lymph node metastasis					
Absent	15	27.3	13	40.6	0.20
Present	40	72.7	19	59.4	
Lymphatic invasion					
Absent	14	25.5	12	37.5	0.24
Present	41	74.5	20	62.5	
Venous invasion					
Absent	41	74.5	21	65.6	0.38
Present	14	25.5	11	34.4	
Liver metastasis					
Absent	50	90.9	32	91.0	0.029 ^a
Present	5	9.1	0	9.0	
Peritoneal dissemination					
Absent	45	81.8	27	84.4	0.76
Present	10	18.2	5	15.6	
Clinical stage					
I-II	22	40	21	65.6	0.020 ^a
III-IV	33	60	11	34.4	

^aP < 0.05.^bTumor invasion of mucosa (m), submucosa (sm), muscularis propria (mp), subserosa (ss), penetration of serosa (se), and invasion of adjacent structures (si).

membranous reactivity in $\geq 10\%$ or more of tumor cells. Specimens with scores of 0 and 1+ were regarded as being negative for ERBB2 expression, whereas scores of 2+ and 3+ indicated positive expression of ERBB2.

Experimental studies

Cell lines and cell culture. The human gastric cancer cell lines AZ521, KATO, MKN1, MKN45, MKN74, NUGC3, and NUGC4 were provided by the Cell Resource Center of Biomedical Research, Institute of Development, Aging and Cancer, Tohoku University. These cell lines were maintained in RPMI 1640 containing 10% FBS with 100 units/mL penicillin and 100 units/mL streptomycin sulfates and cultured in a humidified 5% CO₂ incubator at 37°C.

Transfection of miRNA-125a precursor (*Pre-miR-125a*)

Using NUGC4, a gastric cancer cell line that expresses a high level of *ERBB2* mRNA, either *Pre-miR-125a* or *Pre-miR* negative control (Ambion *Pre-miR* miRNA Precursors Applied Biosystems), was transfected at 30 nmol/L (final concentration) by using Lipofectamine RNAiMAX (Invitrogen Life Technologies) according to the manufacturer's instruction.

In vitro assays

The MTT assay for gastric cancer cell growth after transfection with *Pre-miR-125a* with/without trastuzumab treatment.

Logarithmically growing NUGC4 cells were transfected with *Pre-miR-125a* or *Pre-miR* negative control with or

without addition of trastuzumab (0.1 or 1 $\mu\text{g}/\text{mL}$) and were seeded at 8.0×10^3 cells per well in 96-well flat-bottomed microtiter plates in a final volume of 100 μL of culture medium per well. Cells were incubated in a humidified atmosphere (37°C and 5% CO_2) for 24, 48, 72, and 96 hours after initiation of transfection. MTT assays were used to measure cell proliferation at each period, as described in Supplementary Data. The assay was carried out with 6 replicates.

Plasmid construction

The 3' untranslated region (3'-UTR) and open reading frame (ORF) of *ERBB2* was amplified by RT-PCR. The amplified product was subcloned and ligated into the pmirGLO Dual-Luciferase miRNA Target Expression Vector (Promega). The resultant reporter vector position was confirmed by sequencing and termed Luc-*ERBB2* WT. To make *miR-125a-5p* binding site mutants, positions 37 to 43 of *ERBB2* 3'-UTR (the sequence; CTCAGGG) were mutated to the sequence CACTGCG (mutated nucleotides are underlined), using the QuikChange Lightning Site-Directed Mutagenesis Kit (Stratagene) according to the manufacturer's protocol. The resultant reporter vector position was confirmed by sequencing and termed Luc-*ERBB2* mutant.

Luciferase assay

Luciferase assays were conducted using 1×10^4 NUGC4 cells plated in a 96-well plate. Transfections were done with Lipofectamine 2000 (Invitrogen) in OptiMEM reduced serum media (GIBCO). Cells were transfected with 30 ng of Luc-*ERBB2* WT vector or Luc-*ERBB2* mutant vector and 100 nmol/L of either *Pre-miR* negative control or *Pre-miR-125a*. Twenty-four hours following transfection, cells were assayed for both firefly and *Renilla* luciferase, using Dual-Glo Luciferase Assay System (Promega). All transfection experiments were conducted in triplicate.

ERBB2 and *miR-125a-5p* expression in the NCI60 panel

For analysis of the correlation between *ERBB2* and *miR-125a-5p* expression in the NCI60 panel (21), the normalized expression levels of the cDNA array and the miRNA array were obtained from the Web site of the Genomics and Bioinformatics Group (<http://discover.nci.nih.gov>). The data were analyzed by JMP 5 for Windows software (SAS Institute, Inc.).

Protein expression analysis

Western blotting was used to confirm the expression of *ERBB2* and phosphorylated AKT, BAK1, and p53 in *Pre-miR-125a*-transfected cells. Primary antibodies and dilutions were as follows: *ERBB2* rabbit monoclonal antibody (Epitomics, Inc.) at a 1:500 dilution; AKT rabbit monoclonal antibody (Cell Signaling Technology, Inc.) at a 1:1,000 dilution; phosphorylated AKT (p-AKT) rabbit monoclonal antibody (Cell Signaling Technology, Inc.) at a 1:2,000 dilution; BAK1 (Cell Signaling Technology, Inc.) at a 1:1,000 dilution; p53 (Dako, Inc.) at a 1:1,000

dilution. Detailed protocols are described in Supplementary Data.

Statistical analysis

Data from RT-PCR analysis and *in vitro* transfected cell assays were analyzed with JMP 5. Overall survival rates were calculated actuarially according to the Kaplan–Meier method and were measured from the day of surgery. Differences between groups were estimated using the χ^2 test, Student's *t* test, repeated-measures ANOVA test, and the log-rank test. Variables with a value of $P < 0.05$ in univariate analysis were used in a subsequent multivariate analysis based on the Cox proportional hazards model. A probability level of 0.05 was chosen for statistical significance.

Results

The clinicopathologic significance of *miR-125a-5p* mRNA expression in gastric cancer

In this study, patients with values less than the average expression level of *miR-125a-5p* (8.66, normalized to RNU6B) were assigned to a low expression group ($n = 55$) whereas those with expression values above average were assigned to a high expression group ($n = 32$). Patients in the low *miR-125a-5p* expression group had a significantly poorer prognosis than those in the high *miR-125a-5p* expression group ($P = 0.0069$; Fig. 1). Clinicopathologic factors were significantly different in the low *miR-125a-5p* expression group. There were greater tumor size ($P = 0.0068$), tumor invasion ($P = 0.031$), liver metastasis




Figure 1. Kaplan–Meier overall survival curves of gastric cancer patients according to the level of *miR-125a-5p* expression. Patients in the low *miR-125a-5p* expression group had a significantly poorer prognosis than those in the high *miR-125a-5p* expression group. The high *miR-125a-5p* expression group ($n = 32$) was composed of patients with higher than average expression levels (8.66, normalized to RNU6B); the low *miR-125a-5p* expression group ($n = 55$) had lower than average expression levels.

($P = 0.029$), and clinical staging ($P = 0.02$) than the high *miR-125a-5p* expression group (Table 1). However, no significant differences were observed regarding age, gender, histology, lymphatic invasion, venous invasion, lymph node metastasis, peritoneal dissemination, and distant metastasis (Table 1). The results of univariate and multivariate Cox proportional hazards regression analyses for overall survival are shown in Table 2. Multivariate analysis indicated that the high expression level of *miR-125a-5p* was an independent and significant prognostic factor for survival (OR, 2.44; CI, 1.04–6.73; $P = 0.041$; Table 2). Expression of *ERBB2*, which is a putative *miR-125a-5p* target, is shown to be an indicator of patient prognosis by univariate analysis ($P = 0.048$), although it is not an independent prognostic factor. For the comparison, using the same RNA samples, we also investigated *DACH1* and *PDCD6* mRNA expression, which was previously reported as prognostic factors for gastric cancer patients (22). However, in univariate analysis for overall survival, the expression levels of those 2 molecules were not superior to *miR-125a-5p* expression as prognostic factors, at least in the group we investigated (Supplementary Table S1).

***ERBB2* mRNA expression in gastric cancer cell lines and the effect of trastuzumab**

ERBB2 mRNA expression was examined in 7 gastric cancer cell lines by using RT-PCR. NUGC4, a human cell line derived from a signet ring cell carcinoma of the stomach, showed a remarkably high level of *ERBB2* mRNA compared with other cell lines, ($P < 0.0001$; Supplementary Fig. S1A) and was chosen for experiments on validation of *ERBB2* suppression by *miR-125a-5p*. MTT assays were carried out to evaluate the growth inhibitory effect of trastuzumab in gastric cancer cell lines, NUGC4, AZ521, and NUGC3. AZ521 and NUGC3 were chosen as representative low *ERBB2* expression cell lines. The results indicated that

trastuzumab exerted its activity selectively on NUGC4, the *ERBB2* high expression gastric cancer cell line. At the maximum concentration of 100 $\mu\text{g}/\text{mL}$, the cell viability of NUGC4 was reduced by $28.3\% \pm 3.98\%$ whereas the viability of NUGC3 and AZ521 remained above 90% (Supplementary Fig. S1B).

***miR-125a-5p* regulates *ERBB2* in gastric cancer cells**

Using *in silico* miRNA target prediction tools, such as miRanda (23), PicTar (24), and TargetScan (25), we identified the sequence of the *miR-125a-5p* binding sites in the 3'-UTRs of transcripts encoding *ERBB2* (Fig. 2A). To investigate binding and repression, a luciferase reporter assay was carried out with a vector which included the ORF sequence and 3'-UTR of *ERBB2* downstream from the luciferase reporter gene (Luc-*ERBB2* WT). Transient cotransfection of NUGC4 cells with the reporter plasmid and *Pre-miR-125a* significantly reduced luciferase activity in comparison with the negative control ($P < 0.05$). However, the activity of the reporter construct with mutant sequence (Luc-*ERBB2* mutant) was unaffected by simultaneous transfection with *Pre-miR-125a* (Fig. 2B). These data suggest that *ERBB2* mRNA is a direct functional target of *miR-125a-5p*.

***miR-125a-5p* expression and *ERBB2* protein expression in clinical samples**

Of the 87 gastric cancer patients we examined for the expression of *miR-125a-5p*, FFPE surgical sections were available in 52 cases. To explore the association between *miR-125a-5p* expression and *ERBB2* protein expression status, we carried out immunohistochemical analysis with these samples. The results showed that in the low *ERBB2* expression group (*ERBB2* staining 0 or 1+; $n = 45$), the expression of *miR-125a-5p* was significantly increased compared with that in the high *ERBB2* expression group (*ERBB2* staining 2+ or 3+; $n = 7$; mean \pm SEM; 4.97 ± 0.88 vs. 2.34

Table 2. Univariate and multivariate analysis for overall survival (Cox proportional regression model)

Factors	Univariate analysis			Multivariate analysis		
	RR	95% CI	P	RR	95% CI	P
Age ($\leq 65 / > 66$)	0.928	0.648–1.314	0.673	–	–	–
Sex (male/female)	0.834	0.552–1.121	0.345	–	–	–
Histology grade (well, moderately/poorly and others)	1.309	0.913–1.946	0.146	–	–	–
Depth of tumor invasion ^a (m, sm, mp/ss, se, si)	8.387	2.531–51.86	0.0001 ^b	2.415	0.650–15.87	0.208
Lymph node metastasis (negative/positive)	4.675	2.162–19.74	0.0001 ^b	2.892	1.219–12.88	0.011 ^b
Lymphatic invasion (negative/positive)	2.684	1.474–6.676	0.0003 ^b	1.177	0.481–5.237	0.761
Venous invasion (negative/positive)	3.324	1.627–6.080	0.0012 ^b	1.316	0.883–1.971	0.176
<i>ERBB2</i> mRNA expression (low/high)	1.466	1.003–2.263	0.0482 ^b	1.010	0.991–1.025	0.277
<i>miR-125a-5p</i> expression (high/low)	3.018	1.372–7.588	0.0051 ^b	2.438	1.035–6.727	0.041 ^b

Abbreviation: RR, Relative risk.

^aTumor invasion of mucosa (m), submucosa (sm), muscularis propria (mp), subserosa (ss), penetration of serosa (se), and invasion of adjacent structures (si).

^b $P < 0.05$.




Figure 2. *miR-125a-5p* targets *ERBB2*. A, top, sequence of the *miR-125a-5p* binding sites in the 3'-UTRs of transcripts encoding *ERBB2*. Bottom, schematic diagram of the luciferase reporters in target validation. B, *miR-125a-5p* represses its target in the luciferase assay in NUGC4. Relative luciferase level = (Sample Luc/Sample *Renilla*)/(Control Luc/Control *Renilla*). Luc, raw firefly luciferase activity; *Renilla*, internal transfection control *Renilla* activity; Pre-miR n.c., Pre-miR negative control. The error bar represents the SD from 6 replicates. C, association of *miR-125a-5p* expression with *ERBB2* protein expression status determined by immunohistochemical analysis in 52 gastric cancer cases. In the low *ERBB2* expression group (*ERBB2* staining 0 or 1+; $n = 45$), the expression of *miR-125a-5p* was significantly increased compared with that in the high *ERBB2* expression group (*ERBB2* staining 2+ or 3+; $n = 7$; $P \leq 0.05$). Dots represent the *miR-125a-5p* expression of each sample. Horizontal lines indicate mean value of each group. D, *miR-125a-5p* and *ERBB2* expression levels of carcinoma cells of different origin from the NCI60 tumor cell panel. *ERBB2* expression is inversely correlated with expression of *miR-125a-5p* in 23 cell lines including colon, lung, prostate, and renal cancer.

± 2.24 ; $P < 0.05$; Fig. 2C). Images of immunohistochemistry are shown in Supplementary Figure S2.

***miR-125a-5p* and *ERBB2* expression levels in carcinoma cells of different origin from the NCI60 tumor cell panel**


To evaluate whether the *miR-125a-ERBB2* pathway functions in cells of different origin, we exploited the cDNA array and the miRNA array data set of NCI60, as described in Material and Methods. *ERBB2* expression was inversely correlated with the expression of *miR-125a-5p* in 23 cell lines including colon, lung, prostate, and renal cancer ($r = -0.5348$, $P = 0.0086$; Fig. 2D).

Both *ERBB2* and its primary downstream signal through AKT were suppressed by *miR-125a-5p*

Using RT-PCR, we confirmed that *miR-125a-5p* expression in Pre-miR-125a-treated cells was significantly

higher than that in untreated cells (parent) and in Pre-miR negative control-treated cells ($P < 0.05$, Fig. 3A). To determine whether *miR-125a-5p* suppresses *ERBB2* and its downstream signaling in the gastric cancer cell line NUGC4, cell lysates of transfected cells were analyzed by Western blotting. Remarkable suppression of *ERBB2* and phosphorylated AKT (p-AKT) was observed in Pre-miR-125a-treated cells, in comparison with untreated cells (parent) or Pre-miR negative control-treated cells (Fig. 3B). However, AKT itself did not show significant reduction. We also investigated the inhibition of previously reported *miR-125a-5p* targets, including apoptosis-related gene *BAK1* (26) and tumor suppressor gene *p53* (27). Western blot analysis showed that protein expression of *BAK1* and *p53* was moderately suppressed in Pre-miR-125a-treated cells; however, the reduction was not as significant as that of *ERBB2* (Supplementary Fig. S3).

Figure 3. *miR-125a-5p* regulates ERBB2. A, *miR-125a-5p* expression after treatment with negative control or *Pre-miR-125a* in NUGC4 (quantitative RT-PCR). *miR-125a-5p* expression in *Pre-miR-125a*-treated cells is significantly higher than in untreated cells (parent) and in *Pre-miR* negative control-treated cells. The results are the mean \pm SD of triplicates. B, Western blotting analysis of ERBB2 and phosphorylated AKT (p-AKT) in NUGC4 cells transfected with *Pre-miR-125a* or negative control (n.c.). These proteins were normalized to the level of β -actin.



miR-125a-5p inhibited the proliferation of gastric cancer cells in combination with trastuzumab


To explain the antitumor efficacy of *miR-125a-5p* in gastric cancer cells, a proliferation assay was carried out with *Pre-miR-125a*-treated cells or negative control cells by using NUGC4. Furthermore, we investigated whether the additional administration of trastuzumab, the ERBB2-targeting antibody, enhanced the antitumor efficacy of *miR-125a-5p*. These experiments were carried out at 2 different concentrations of trastuzumab [0.1 μ g/mL (Fig. 4A) and 1 μ g/mL (Fig. 4B)]. Interestingly, *miR-125a* not only potently suppressed the proliferation of gastric cancer cells by itself but also inhibited growth more potently when combined with trastuzumab ($P = 0.0032$, Fig. 4A; $P = 0.0033$, Fig. 4B). The combined growth inhibitory effect was more robust at the higher concentration of trastuzumab.

Discussion

Recent evidence has shown that altered patterns of miRNA expression correlate with various human cancers. The behavior of miRNAs is complex because they regulate hundreds of targets, resulting in the downregulation of numerous target genes including oncogenes and tumor suppressors. Therefore, exploring their clinical potential is especially worthwhile.

In the current study, we show that altered *miR-125a-5p* expression significantly affected cancer progression and prognosis in human gastric cancer. Multivariate analysis revealed that *miR-125a-5p* is an independent prognostic factor for survival. Clinicopathologic analysis revealed that low *miR-125a-5p* expression contributes to more advanced tumor size and tumor invasion (Table 1). It

Figure 4. *miR-125a-5p* inhibits the proliferation of the gastric cancer cell line NUGC4 in combination with trastuzumab. *Pre-miR-125a* or *Pre-miR* negative control transfected with or without trastuzumab (A, 0.1 μ g/mL; B, 1 μ g/mL) treatment were seeded at 8.0×10^3 cells per well in 96-well plates and cell growth was monitored every 24 hours using the MTT assay. Absorbance at 0 hour was assigned a value of 1. The results are the mean \pm SD of 3 replicates.



suggests that this miRNA primarily achieves its antiproliferative effect through downregulation of proliferation-related genes, including *ERBB2*, a member of the EGF receptor family of receptor tyrosine kinases, which regulate a key initiator of phosphoinositide-3 kinase (PI3K)-AKT and RAS/RAF/mitogen-activated protein kinase signaling (28). *miR-125a-5p* is shown to be a superior biomarker to previously reported gastric cancer biomarkers such as *DACH1* and *PDCD6* (ref. 22; Supplementary Table S1). However, because of the differences in patient backgrounds such as clinical stage and the presence or absence of chemotherapy, further investigation is required for adequate use of these biomarkers.

We confirmed *miR-125a-ERBB2* interaction in the human gastric cancer cell line NUGC4. *MiR-125a-5p* significantly repressed *ERBB2* expression and the phosphorylation of its downstream molecule, AKT (Fig. 3B). In addition, *ERBB2* expression was shown to be inversely correlated with expression of *miR-125a-5p* both *in vitro* and in clinical samples. Overexpression of *Pre-miR-125a* also led to the inhibition of previously reported *miR-125a-5p* targets, such as apoptosis-related gene *BAK1* (26) and tumor suppressor gene *p53* (ref. 27; Supplementary Fig. S3). However, the inhibition of these tumor suppressor genes was modest compared with that of *ERBB2*, suggesting *ERBB2* is a crucial target of *miR-125a-5p*, at least in the gastric cancer cell line NUGC4.

It is noteworthy that the growth inhibitory effect of *miR-125a-5p* was enhanced when combined with trastuzumab (Fig. 4A and B). This could be partly due to the fact that *miR-125a-5p* and trastuzumab share the same target, *ERBB2*. *miR-125a-5p* and trastuzumab silence the *ERBB2* pathway through 2 different mechanisms. *miR-125a-5p* suppresses the molecule at the posttranscriptional level before protein synthesis, whereas trastuzumab is a monoclonal antibody targeted against completed *ERBB2* protein. In other words, *miR-125a-5p* blocks the synthesis of the oncoprotein at an earlier phase than does trastuzumab. These considerations suggest that *miR-125a-5p* mimic and

trastuzumab have the potential to be highly effective against *ERBB2* when used together.

ERBB2-positive gastric cancer patients constitute about 19.0% (8.2%–53.4%) of all gastric cancer patients (19, 29, 30). A recent phase III study (the ToGA trial) combining treatment of trastuzumab and conventional chemotherapy against *ERBB2*-positive gastric cancer showed a statistically significant advantage in overall survival for patients who received combined therapy compared with chemotherapy alone. These reliable large-scale clinical data indicate that *ERBB2* is a crucial therapeutic target in gastric cancer (31).

In conclusion, our data suggest that *miR-125a-5p* functions as a powerful tumor suppressor and could be a *bona fide* prognostic marker for gastric cancer patients. Furthermore, *miR-125a-5p* mimic alone or in combination with trastuzumab could be a novel therapeutic approach against gastric cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank T. Shimooka, K. Ogata, M. Kasagi, and T. Kawano for their excellent technical assistance.

Grant Support

This work was supported in part by the following grants and foundations: CREST, Japan Science and Technology Agency (JST); Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research: 21679006, 20390360, 20590313, 20591547, 21591644, 21592014, 20790960, 21791297, 21229015, 20659209, and 20012039; NEDO (New Energy and Industrial Technology Development Organization) Technological Development for Chromosome Analysis; The Ministry of Education, Culture, Sports, Science and Technology of Japan for Scientific Research on Priority Areas, Cancer Translational Research Project, Japan.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received August 7, 2010; revised December 6, 2010; accepted December 18, 2010; published online May 2, 2011.

References

- Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009;136:215–33.
- Nicoloso MS, Spizzo R, Shimizu M, Rossi S, Calin GA. MicroRNAs—the micro steering wheel of tumour metastases. *Nat Rev Cancer* 2009;9:293–302.
- Spizzo R, Nicoloso MS, Croce CM, Calin GA. SnapShot: microRNAs in cancer. *Cell* 2009;137:586–e1.
- Guo X, Wu Y, Hartley RS. MicroRNA-125a represses cell growth by targeting HuR in breast cancer. *RNA Biol* 2009;6:575–83.
- Li W, Duan R, Kooy F, Sherman SL, Zhou W, Jin P. Germline mutation of microRNA-125a is associated with breast cancer. *J Med Genet* 2009;46:358–60.
- O'Day E, Lal A. MicroRNAs and their target gene networks in breast cancer. *Breast Cancer Res* 2010;12:201.
- Nam EJ, Yoon H, Kim SW, Kim H, Kim YT, Kim JH, et al. MicroRNA expression profiles in serous ovarian carcinoma. *Clin Cancer Res* 2008;14:2690–5.
- Wang G, Mao W, Zheng S, Ye J. Epidermal growth factor receptor-regulated miR-125a-5p—a metastatic inhibitor of lung cancer. *FEBS J* 2009;276:5571–8.
- Ferretti E, De Smaele E, Po A, Di Marcotullio L, Tosi E, Espinola MS, et al. MicroRNA profiling in human medulloblastoma. *Int J Cancer* 2009;124:568–77.
- Cowden Dahl KD, Dahl R, Kruichak JN, Hudson LG. The epidermal growth factor receptor responsive miR-125a represses mesenchymal morphology in ovarian cancer cells. *Neoplasia* 2009;11:1208–15.
- Park NJ, Zhou H, Elashoff D, Henson BS, Kastratovic DA, Abemayor E, et al. Salivary microRNA: discovery, characterization, and clinical utility for oral cancer detection. *Clin Cancer Res* 2009;15:5473–7.
- Scott GK, Goga A, Bhaumik D, Berger CE, Sullivan CS, Benz CC. Coordinate suppression of *ERBB2* and *ERBB3* by enforced expression of micro-RNA miR-125a or miR-125b. *J Biol Chem* 2007;282:1479–86.
- Gravalos C, Jimeno A. HER2 in gastric cancer: a new prognostic factor and a novel therapeutic target. *Ann Oncol* 2008;19:1523–9.

14. Marx AH, Tharun L, Muth J, Dancau AM, Simon R, Yekebas E, et al. HER-2 amplification is highly homogenous in gastric cancer. *Hum Pathol* 2009;40:769–77.
15. Kim SY, Kim HP, Kim YJ, Oh do Y, Im SA, Lee D, et al. Trastuzumab inhibits the growth of human gastric cancer cell lines with HER2 amplification synergistically with cisplatin. *Int J Oncol* 2008;32:89–95.
16. Matsui Y, Inomata M, Tojigamori M, Sonoda K, Shiraishi N, Kitano S. Suppression of tumor growth in human gastric cancer with HER2 overexpression by an anti-HER2 antibody in a murine model. *Int J Oncol* 2005;27:681–5.
17. Japanese Gastric Cancer Association. Japanese Classification of Gastric Carcinoma—2nd English edition. *Gastric Cancer* 1998;1:10–24.
18. Aihara H, Kawamura YJ, Toyama N, Mori Y, Konishi F, Yamada S, et al. Analysis of the gene-expression profile regarding the progression of human gastric carcinoma. *Surgery* 2002;131:S39–47.
19. Hofmann M, Stoss O, Shi D, Büttner R, van de Vijver M, Kim W, et al. Assessment of a HER2 scoring system for gastric cancer: results from a validation study. *Histopathology* 2008;52:797–805.
20. Rüschoff J, Dietel M, Baretton G, Arbogast S, Walch A, Monges G, et al. HER2 diagnostics in gastric cancer—guideline validation and development of standardized immunohistochemical testing. *Virchows Arch* 2010;457:299–307.
21. Blower PE, Verducci JS, Lin S, Zhou J, Chung JH, Dai Z, et al. MicroRNA expression profiles for the NCI-60 cancer cell panel. *Mol Cancer Ther* 2007;6:1483–91.
22. Yamada Y, Arai T, Gotoda T, Taniguchi H, Oda I, Shirao K, et al. Identification of prognostic biomarkers in gastric cancer using endoscopic biopsy samples. *Cancer Sci* 2008;99:2193–9.
23. Betel D, Wilson M, Gabow A, Marks DS, Sander C. The microRNA.org resource: targets and expression. *Nucleic Acids Res* 2008;36:D149–53.
24. Krek A, Grün D, Poy MN, Wolf R, Rosenberg L, Epstein EJ, et al. Combinatorial microRNA target predictions. *Nat Genet* 2005;37:495–500.
25. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 2005;120:15–20.
26. Guo S, Lu J, Schlanger R, Zhang H, Wang JY, Fox MC, et al. MicroRNA miR-125a controls hematopoietic stem cell number. *Proc Natl Acad Sci U S A* 2010;107:14229–34.
27. Zhang Y, Gao JS, Tang X, Tucker LD, Quesenberry P, Rigoutsos I, et al. MicroRNA 125a and its regulation of the p53 tumor suppressor gene. *FEBS Lett* 2009;583:3725–30.
28. Hsieh AC, Moasser MM. Targeting HER proteins in cancer therapy and the role of the non-target HER3. *Br J Cancer* 2007;97:453–7.
29. Allgayer H, Babic R, Gruetzner KU, Tarabichi A, Schildberg FW, Heiss MM. c-erbB-2 is of independent prognostic relevance in gastric cancer and is associated with the expression of tumor-associated protease systems. *J Clin Oncol* 2000;18:2201–9.
30. Jorgensen JT. Targeted HER2 treatment in advanced gastric cancer. *Oncology* 2010;78:26–33.
31. Bang YJ, Van Cutsem E, Feyereislova A, Chung HC, Shen L, Sawaki A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet* 2010;376:687–97.

Clinical Cancer Research

MicroRNA-125a-5p Is an Independent Prognostic Factor in Gastric Cancer and Inhibits the Proliferation of Human Gastric Cancer Cells in Combination with Trastuzumab

Naohiro Nishida, Koshi Mimori, Muller Fabbri, et al.

Clin Cancer Res 2011;17:2725-2733. Published OnlineFirst January 10, 2011.

Updated version	Access the most recent version of this article at: doi: 10.1158/1078-0432.CCR-10-2132
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2011/05/05/1078-0432.CCR-10-2132.DC1

Cited articles	This article cites 31 articles, 7 of which you can access for free at: http://clincancerres.aacrjournals.org/content/17/9/2725.full#ref-list-1
-----------------------	---

Citing articles	This article has been cited by 7 HighWire-hosted articles. Access the articles at: http://clincancerres.aacrjournals.org/content/17/9/2725.full#related-urls
------------------------	---

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
----------------------	--

Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
-----------------------------------	--

Permissions	To request permission to re-use all or part of this article, use this link http://clincancerres.aacrjournals.org/content/17/9/2725 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.
--------------------	--