Published OnlineFirst October 2, 2012; DOI: 10.1158/1078-0432.CCR-12-2931

Clinical Cancer Research

Correction

Correction: The Use of Quantitative Real-Time Reverse-Transcriptase PCR for 5′ and 3′ Portions of *ALK* Transcripts to Detect *ALK* Rearrangements in Lung Cancers

In this article (Clin Cancer Res 2012;18:4725–32), which was published in the September 1, 2012, issue of *Clinical Cancer Research* (1), the new fusion variants of *ELM4-ALK* in Fig. 4C are incorrect due to a typing error.



Below is the corrected figure.



As a result, the new *EML4-ALK* transcript has a 54 bp deletion, not a 53 bp deletion, in the 5′ portion of the original *EML4* exon 10. The number "53" on page 4728, paragraph 6, should be "54." The authors regret the error.

Reference

1. Wang R, Pan Y, Li C, Hu H, Zhang Y, Li H, et al. The use of quantitative real-time reverse-transcriptase PCR for 5′ and 3′ portions of *ALK* transcripts to detect *ALK* rearrangements in lung cancers. Clin Cancer Res 2012;18:4725–32.

Published OnlineFirst October 2, 2012.
doi: 10.1158/1078-0432.CCR-12-2931

AACR American Association for Cancer Research

Clinical Cancer Research

Correction: The Use of Quantitative Real-Time Reverse-Transcriptase PCR for 5′ and 3′ Portions of *ALK* Transcripts to Detect *ALK* Rearrangements in Lung Cancers

Clin Cancer Res 2012;18:6079. Published OnlineFirst October 2, 2012.