

## BRAF(V600) Inhibitor GSK2118436 Targeted Inhibition of Mutant BRAF in Cancer Patients Does Not Impair Overall Immune Competency

David S. Hong<sup>1</sup>, Luis Vence<sup>2</sup>, Gerald Falchook<sup>1</sup>, Laszlo G. Radvanyi<sup>2</sup>, Chengwen Liu<sup>2</sup>, Vicki Goodman<sup>3</sup>, Jeffery J. Legos<sup>3</sup>, Sam Blackman<sup>3</sup>, Antonio Scarmadio<sup>1</sup>, Razelle Kurzrock<sup>1</sup>, Gregory Lizee<sup>2</sup>, and Patrick Hwu<sup>2</sup>

### Abstract

**Purpose:** An intact immune system likely contributes to the outcome of treatment and may be important for clearance of drug-resistant tumor cells and for prevention of recurrence. Although pharmacologic inhibition of BRAF(V600E) in melanoma patients, which is linked to immune suppression, results in an initial response rate, these responses are typically of limited duration. Combining immunotherapeutic drugs with kinase-targeted agents is an attractive strategy to increase clinical efficacy. Evidence suggesting that mitogen-activated protein kinase pathway activation in tumor cells contributes to immune suppression suggests that the two approaches may be synergistic, provided that BRAF(V600E) inhibitors are nontoxic to immune cells.

**Methods:** To assess effects of mutant BRAF inhibition on systemic immunity, we studied 13 patients with tumors carrying a BRAF mutation who underwent treatment with GSK2118436, a V600 mutant BRAF-specific inhibitor. We carried out peripheral blood immunomonitoring before and following one or two 28-day cycles of treatment.

**Results:** GSK2118436 treatment had no detectable impact on most immune parameters tested, including serum cytokine levels, peripheral blood cell counts, leukocyte subset frequencies, and memory CD4<sup>+</sup> and CD8<sup>+</sup> T-cell recall responses. A slight increase in serum TNF- $\alpha$  over the course of treatment was observed. In addition, three of the four human leukocyte antigen-A2–positive patients experienced a modest increase in circulating tumor antigen–specific CD8<sup>+</sup> T cells following BRAF(V600) inhibitor therapy.

**Conclusions:** GSK2118436 treatment results in no detectable negative impact on existing systemic immunity or the *de novo* generation of tumor-specific T cells. These findings suggest that future trials combining specific BRAF(V600E) inhibition with immunotherapy should not impair immune response. *Clin Cancer Res*; 18(8); 2326–35. ©2012 AACR.

### Introduction

Metastatic melanoma (MM) is one of the deadliest of cancers, with stage IV patients facing a less than 10% five-year survival rate (1–3). The poor survival in the metastatic setting and its increasing annual incidence has created an urgent need for effective new therapies. Two of the most

successful strategies have been to utilize immune-modulating drugs or pharmacologic agents that target oncogenic kinases that constitutively activate the mitogen–activated protein kinase (MAPK) pathway in tumor cells (4, 5). Both approaches have produced a clinical benefit in melanoma patients, and there is an emerging rationale that supports combining these 2 approaches in future clinical trials (6, 7).

Approximately 50% to 60% of melanomas express an activating mutation in the BRAF oncogene at amino acid position 600, with about 85% to 95% of the V600 being V600E mutations and 10% to 15% V600K mutations. These genetic alterations are also found, but to a lesser extent, in colorectal cancer and papillary thyroid carcinoma (8–10). The aberrations result in constitutive activation of the MAPK signaling pathway, which favors tumor cell growth through various mechanisms, including reduced apoptosis, increased metastatic potential, invasiveness, and immune suppression (11–16). New therapies that target the MAPK pathway have produced demonstrable efficacy in the clinic, supporting this approach. Recently, inhibitors targeting BRAF(V600E) have directly induced objective responses in

**Authors' Affiliations:** <sup>1</sup>Department of Investigational Cancer Therapeutics (Phase I Clinical Trials Program), Division of Cancer Medicine; <sup>2</sup>Department of Melanoma Medical Oncology, MD Anderson Cancer Center, Houston, Texas; and <sup>3</sup>GlaxoSmithKline, Collegeville, Pennsylvania

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

D.S. Hong, L. Vence, G. Lizee, and P. Hwu contributed equally to this work.

**Corresponding Author:** David Hong, Department of Investigational Therapeutics (Phase 1 Program)-Unit 455, Division of Cancer Medicine, MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030. Phone: 713-563-5844; Pager: 713-606-3113; Fax: 713-792-0305; E-mail: dshong@mdanderson.org

doi: 10.1158/1078-0432.CCR-11-2515

©2012 American Association for Cancer Research.

### Translational Relevance

The mutant BRAF-specific inhibitor, GSK2118436, is a potent antitumor agent beginning to be widely used in cancer patients harboring BRAF mutations at the valine 600 position (V600). However, little is known on whether this drug negatively affects lymphocyte subsets and memory T-cell responses *in vivo* during therapy. In this study, we measured a number of immune parameters in the peripheral blood of patients during mutant BRAF-specific inhibitor therapy and found no detectable negative impact on existing systemic immunity or the *de novo* generation of tumor-specific T cells. The study findings strongly suggest that future trials combining V600 mutant BRAF inhibition with immunotherapy will not impair immune response as was found with previous mitogen-activated protein kinase inhibitors that were less specific and often resulted in significant detrimental effects on the immune system of cancer patients undergoing therapy.

more than 70% of MM patients whose tumors harbor the V600E mutation (5, 17, 18). Yet in spite of this clinical success, the median duration of response to the most effective of these inhibitors was only 8 months, possibly through BRAF-independent activation of the mitogen-activated protein/extracellular signal-regulated kinase (ERK; MEK) and ERK signaling pathway (19–21).

Immunotherapeutic approaches that activate T cell-mediated immune responses have also met with mixed clinical success, particularly in the melanoma setting (22). These include interleukin-2 (IL-2) therapy, adoptive transfer of expanded tumor-infiltrating lymphocytes (TIL), dendritic cell (DC)-based vaccines, and more recently, antibody-based therapies that target the negative regulatory molecules, cytotoxic T lymphocyte-associated antigen (CTLA)-4 or programmed death receptor (PD)-1 (23–27). Some of these therapies have mediated objective responses and extended survival in MM patients, but disease relapses nevertheless occur in most treated patients.

In light of these clinical findings, future approaches that combine immunotherapeutic drugs with kinase-targeted agents provide an attractive strategy that may potentially increase clinical response rates, with the ultimate goal of improved survival (28). Furthermore, accumulating evidence that MAPK pathway activation in tumor cells contributes to immune suppression suggests that these 2 approaches may be synergistic (16). However, because many immune cells use the MAPK pathway to mediate their own function and survival, they can be sensitive to the effects of MAPK pathway inhibitors. Indeed, previous generations of MAPK inhibitors have produced detrimental effects on systemic immunity, and some therapies have been shown to directly cause lymphopenia in treated patients (29–32). Some of these off-target effects can be attributed to the lack of specificity of the agent and/or the

targeting of wild-type kinases that are essential for immune cell activity.

Newer compounds that specifically target mutated oncogenes like BRAF(V600E) are theoretically less likely to be detrimental to the immune system because the mutation is limited to tumor cells. Laboratory studies have confirmed this notion, as exposure to BRAF(V600E)-specific inhibitors seemed to be much less toxic to human T lymphocytes *in vitro* than an MEK inhibitor (33, 34). However, due to the complexities of the immune system and issues related to pharmacokinetics and bioavailability, it is critical to examine the effects of mutant BRAF inhibition on systemic immunity *in vivo* using more direct *ex vivo* assays on patient blood cells. Here, we report the results of an immune monitoring study in which cancer patients were treated with GSK2118436 (35), a specific inhibitor of mutated BRAF at V600 capable of inhibiting BRAF(V600E), BRAF(V600K), and BRAF(V600G) mutations.

### Materials and Methods

#### Patients

A companion blood collection protocol (local protocol # LAB09-0114) approved by the MD Anderson Cancer Center Institutional Review Board was introduced to patients enrolled on the GlaxoSmithKline study of "A Phase I Open-label, Multi-dose, Dose-escalation Study to Investigate the Safety, Pharmacokinetics and Pharmacodynamics of the BRAF inhibitor GSK2118436 in Subjects with Solid Tumors" (local protocol #2009-0142, NCTNCT00880321). Enrollment on the companion study was optional and only those patients who met eligibility criteria for a targeted therapy trial were approached. Inclusion criteria for the companion protocol included: patients with pathologically confirmed advanced or metastatic cancer (any tumor type); a blood sample adequate to determine baseline (pretreatment) immune response status; scheduled to receive a tyrosine kinase inhibitor regimen; and ability to understand and willingness to sign a written consent document. Exclusion criteria included: immunocompromised patients (i.e., receiving steroid or nonsteroidal anti-inflammatory medication); patients who had a bone marrow transplant at any time before the study, or any other type of transplant within 30 days before day 1 of study treatment; absolute neutrophil counts less than 1,500/mL; uncontrolled intercurrent illness including, but not limited to, ongoing or active infection requiring parenteral antibiotics; known history of HIV1 and HIV2, hepatitis B or hepatitis C; a known hypersensitivity to any of the components or metabolites of the drug products evaluated in this study; and psychologic, familial, sociologic, or geographic conditions that would not permit study follow-up and compliance with the study protocol.

#### Peripheral blood mononuclear cell collection and processing

Blood specimens (60 mL; 6 × 10 mL heparinized tubes) were taken for peripheral blood mononuclear cell (PBMC)

isolation and T-cell number determination. Samples were collected using standard techniques for "Green Top" Vacutainer™ Blood Collection Tubes. PBMCs were isolated with Ficoll-Hypaque gradient, frozen at  $-80^{\circ}\text{C}$  overnight and then transferred to liquid nitrogen. This initial isolation, done by centrifugation (1500 to 1800 RCF, for 20 minutes in a swingout head horizontal rotor) was within 4 hours of blood collection. For analysis, PBMCs were removed from liquid nitrogen and thawed in a  $37^{\circ}\text{C}$  water bath. Thereafter, cells were washed with 20 mL of complete medium (RPMI 1640 plus 10% AB serum), centrifuged at 1,500 rpm for 10 minutes, and suspended in complete medium.

### Multiplex cytokine assay

We measured cytokine levels from sera using a solid phase-based multiplex platform from Meso Scale Discovery (MSD). Each well of the 96-well plate-based assays contained antibodies to GM-CSF, IL-1 $\beta$ , IL-2, IL-12p70, IFN $\gamma$ , IL-6, IL-8, IL-10, TNF- $\alpha$  ("Proinflammatory 9-Plex MULTI-SPOT" 96-well plate kit), as well as VEGF and MCP-1 (both singleplex kits). The MSD platform used in this study consisted of a 96-well plate-based assay with electrochemiluminescence as the basis for detection. Following the capture of the cytokine by the spotted antibody, labeled secondary detection antibodies were bound to the antigen. The detection antibodies were coupled to an electrochemiluminescent label that emitted light when electrochemically stimulated via carbon-coated electrodes in the bottom of the array wells. In addition, a detection buffer that was added before reading the array incorporated reactants that enhanced the electrochemical signal. The resulting signal was read with a charge-coupled device (Sector Imager 2400, MSD).

Calibration curves were prepared in the supplied assay diluent for human serum, with a range of 2,500 to 0.2 pg/mL (Proinflammatory 9-Plex and MCP-1) or 100,000 to 10 pg/mL (VEGF). Arrays were preincubated with 25  $\mu\text{L}$  per well of assay diluent for 30 minutes, with shaking at room temperature. Following the preincubation, 25  $\mu\text{L}$  of sample serum or calibrator was added in duplicate to the appropriate wells. The array was then incubated at room temperature for 2 hours with shaking and washed with Dulbecco's phosphate-buffered saline plus 0.05% Tween 20, and 25 mL of detection antibody reagent. Following a 2-hour room temperature incubation with shaking, the array was washed and detection buffer was added. The assay results were read with an MSD Sector Imager 2400 incorporating the CCD. Sample cytokine concentrations were determined with Discovery Workbench software.

### Recall antigen-specific ELISPOT assay

To test patients' immunocompetence, we measured the ability of CD8 $^{+}$  and CD4 $^{+}$  T cells in PBMCs to react against a series of common recall antigens using a quantitative enzyme-linked immunosorbent spot (ELISPOT) assay measuring the frequency of IFN- $\gamma$ -releasing cells. IFN- $\gamma$  ELISPOT assays were conducted at pretreatment, 3 weeks

posttreatment initiation (at the end of cycle 1), and 6 weeks posttreatment initiation (at the end of cycle 2). Forty-three peptides derived from cytomegalovirus (CMV), Epstein-Barr virus (EBV), and influenza (designated as CEF), presented on a large array of human leukocyte antigen (HLA) class I subtypes covering more than 90% of the human population were used to detect CD8 $^{+}$  T-cell responses against the CEF peptide pool. A similar ELISPOT approach was used to measure patient CD4 $^{+}$  T-cell immunocompetence using a recall antigen response against a chemically inactivated tetanus toxoid (TT), an agent that essentially all humans have been exposed to through vaccination. Phorbol 12-myristate 13-acetate/ionomycin was used as a positive control for IFN- $\gamma$  release. In brief, cryopreserved PBMCs were cultured overnight at  $5 \times 10^5$  cells per well in a 96-well ELISPOT plate for detection of spots with a pair of IFN- $\gamma$ -specific antibodies (Mabtech) in medium containing the 43 CEF peptide pool (each at 2  $\mu\text{g}/\text{mL}$ ), or inactivated TT (4 Lf/mL), or no peptide (negative control). ELISPOTs were developed, dried, and read with a CTL ImmunoSpot ELISPOT reader, using the software programs Image Acquisition 4.5 and Immunospot 5 Professional (Cellular Technology Ltd.). Peptide-specific immune reactivity was determined by subtracting the background spots in wells without peptide. A positive response was defined as peptide-specific spots that were statistically higher (triplicates) than control wells using a 2-tailed *t* test ( $P < 0.05$ ). A zero response was assigned if the peptide-specific wells were not different than control wells. The counts for the 43 CEF peptide pool or TT were presented as the total CEF or tetanus-specific T cells per million PBMCs assessed at each time point, respectively.

### Flow cytometric analysis

Multiparameter flow cytometry analysis of different cell subsets was conducted with PBMCs from patients at different time points. Cells were stained with different antibody panels to study different lymphocyte populations. The LIVE/DEAD Fixable Aqua Dead Cell Stain Kit (Invitrogen) was used in all tubes to exclude dead cells, except for the apoptosis panel. To study dendritic and natural killer (NK) cells, we used the following panel: CD11c-V450, CD303 (BDCA-2)-FITC (FITC, fluorescein isothiocyanate), CD123-PE, CD3/CD14/CD19-PerCP-Cy5.5 (dump channel), CD86-APC, CD56-PE-Cy7, and CD16-APC-C7. For the T-regulatory cell panel, we stained for Foxp3-eFluor450 (eBioscience), CD25-FITC, ICOS-PE (eBioscience), CD4-PerCP-Cy5.5, CD39-APC, CD45RA-PE-Cy7, and CD8-APC-H7. For the apoptosis panel, we stained for CD3-AmCyan, CD19-V450, CD4-FITC, Annexin V-PE, 7-AAD (PerCP-Cy5.5), CD8-APC, CD56-PE-Cy7, and CD16-APC-C7. The frequencies of the different populations were translated into cell numbers (# cells/ $\mu\text{L}$  of blood) using lymphocyte absolute counts (LAC).

All patients were typed for HLA-A2 expression using an anti-HLA-A2-FITC antibody (BD Biosciences). All patients positive for HLA-A2 were tested for the presence of melanoma antigen-specific T cells specific for gp100, MART-1,

**Table 1.** Patient characteristics

Patient # (protocol identifier)	GSK2118436 dose	Sex	Age, y	Tumor histology	BRAF mutation status	Best response and % reduction RECIST (%)
01 (5118)	200 mg b.i.d	F	69	Melanoma	V600E	Not restaged
05 (5119)	200 mg b.i.d	M	52	Melanoma	K601E	PD, 38%
08 (5120)	200 mg b.i.d	M	45	Melanoma	V600E	Unconfirmed PR, -53%
14 (5127)	100 mg t.i.d	F	74	Papillary thyroid	V600E	Confirmed PR, -37%
16 (5128)	100 mg t.i.d	F	66	CRC	V600E	SD
18 (5131)	150 mg b.i.d	M	35	Melanoma	V600E	Unconfirmed PR, -66%
19 (5132)	150 mg b.i.d	F	61	Melanoma	V600E	Confirmed PR, -50%
20 (5133)	200 mg b.i.d	F	76	Melanoma	V600E	PD-new lesion
21 (5134)	200 mg b.i.d	F	62	Melanoma	V600E	SD, -22%
25 (5203)	150 mg b.i.d	M	56	Melanoma	V600E	PR, -54%
34 (5136)	75 mg b.i.d	M	56	Melanoma	V600E	Unconfirmed PR, -33%
35 (5304)	150 mg b.i.d	M	48	CRC	V600E	SD, 1%
40 (5305)	150 mg b.i.d	F	57	Papillary thyroid	V600E	SD, -10%

Abbreviations: CRC, colorectal cancer; PR, partial response; SD, stable disease; PD, progressive disease.

tyrosinase, NY-ESO1, or survivin. The cells were stained with CD62L-V450, CD45RA-FITC, CD3-PE, CD27-PerCP-Cy5.5, CD14/CD16/CD19-PE-Cy7 (dump channel), and CD8-APC-H7. The APC channel was then used to accommodate the class I HLA-A2-restricted gp100 or tyrosinase APC-labeled tetramer (Beckman Coulter), class I HLA-A2-restricted survivin APC-labeled pentamer (Proimmune), or class I HLA-A2-restricted NY-ESO1 or MART-1 APC-labeled dextramer (Immudex). The net tetramer/pentamer/dextramer increase was calculated by subtracting the net baseline frequency (baseline frequency-control tetramer/pentamer/dextramer frequency) from the net frequency at any given time point. All antibodies are from BD Biosciences, unless otherwise indicated, and the acquisition was carried out on a FACS Canto II flow cytometer (BD Biosciences). All analysis was done with the software FlowJo (Tree Star). The frequencies of the multimer<sup>+</sup> CD8 T-cell populations were translated into cell numbers (# cells/mL of blood) using LAC.

### Statistical considerations

Data were summarized descriptively as frequency counts, means, and fold change from baseline. The paired *t* test and/or the Wilcoxon signed-rank test were used to compare changes in quantitative outcomes from baseline to postbaseline assessments (cycle 1 or 2). All statistical tests were 2-sided and *P* < 0.05 was considered statistically significant. Statistical tests were conducted with Prism 4 software.

## Results


### Serum cytokine levels remained largely unaltered during GSK2118436 mutant BRAF inhibitor treatment

Thirteen patients had blood drawn before treatment with the GSK2118436 V600 mutant BRAF-specific inhibitor as well as at the end of each of two 21-day treatment cycles

(cycles 1 and 2). Patient characteristics are listed in Table 1. All patient tumors contained the BRAF(V600E) mutation (GTG to GAG) that changes the encoding amino acid at position 600 from valine (V) to glutamic acid (E) except for 1 patient (05) who had a K601E mutation on exon 15 of the *Raf* gene. Of the 13 patients enrolled, 9 presented with melanoma, 2 had colorectal cancer, and 2 patients had papillary thyroid cancer. Six patients were treated at the 150 mg twice daily dose of GSK2118436, whereas 5 were treated at the 200 mg twice daily dose, and 2 at 100 mg 3 times a day. Six men and 7 women participated in the study, and the median age was 57 years. Similar to other BRAF (V600E)-specific inhibitors, there were significant clinical responses primarily in melanoma patients to drug treatment that will be reported separately.

To detect potential changes in systemic immunity in response to GSK2118436 treatment, we assessed the levels of several proinflammatory cytokines (GM-CSF, IFN- $\gamma$ , IL-2, IL-10, IL-12p70, IL-1 $\beta$ , IL-6, IL-8, and TNF- $\alpha$ ), in addition to MCP-1 and VEGF, in the serum of all patients before and after 1 or 2 cycles of treatment. We found no significant differences over the course of treatment for all cytokines measured (Supplementary Fig. S1), with the exception of TNF- $\alpha$ . As shown in Fig. 1, serum levels of TNF- $\alpha$  increased slightly, during the course of treatment, with mean baseline levels of 9.1 pg/mL rising to 11.2 pg/mL by the end of cycle 1 (*P* = 0.003). The increase in serum TNF- $\alpha$  levels was quite modest for most of the patients, but 1 patient (patient 08) showed a sharp increase in serum TNF- $\alpha$  levels, particularly at the end of treatment cycle 2 (Fig. 1). Interestingly, this patient had a positive blood culture for a bacterial infection and was treated for it during the same period that the rise in TNF- $\alpha$  levels was seen, which may explain this rise. Without this patient, the slight increase in serum TNF- $\alpha$  levels between the basal level and cycle 1 is significant (from *P* = 0.003 to 0.004), and the borderline significant increase





**Figure 1.** TNF- $\alpha$  levels increase through time in the sera of cancer patients treated with the mutant BRAF inhibitor. Sera from the same 13 patients treated with the GSK2118436 V600 mutant BRAF inhibitor were tested for TNF- $\alpha$ , GM-CSF, IFN- $\gamma$ , IL-10, IL-12p70, IL-1b, IL-2, IL-6, and IL-8 throughout the course of the treatment (baseline, cycle 1, and cycle 2) using a multiple cytokine assay (Luminex). TNF- $\alpha$  was the only cytokine to show a significant change (increase) throughout the treatment. The change shown by the other cytokines was not significant (see Supplementary Fig. S1).

between the basal level and cycle 2 becomes significant (from  $P = 0.061$  to  $0.013$ ). Overall, the results indicate that serum cytokine levels remained largely unaffected by the V600 mutant BRAF inhibitor.


#### Mutant BRAF inhibition with GSK2118436 did not significantly alter the cell number (or frequency) of most immune cell subsets in peripheral blood

Because many immune cell types require MAPK pathway signaling for normal function, we analyzed the peripheral blood frequencies of different immune cell

types to determine whether any changes could be observed over the course of GSK2118436 treatment. As shown in Fig. 2, flow cytometric analysis showed that overall cell number of T cells ( $CD4^+$  and  $CD8^+$ ), B cells, NK cells (cytotoxic and cytokine producing), DC (mDC and pDC), monocytes, and regulatory T cells (naïve and effector regulatory T cells) in peripheral blood were not significantly altered following the initiation of mutant BRAF inhibitor therapy. Although fluctuations were observed in individual patients, as a group these changes were not significant and were not likely to be attributed to the effects of drug treatment. The frequency (percentage) of these lymphocyte subpopulations was also unchanged (Supplementary Fig. S2).


#### Mutant BRAF inhibition with GSK2118436 did not significantly impact T cell-mediated recall responses in peripheral blood

To assess the effects of GSK2118436 treatment on functional T cell-mediated immune responses, patient PBMC samples were analyzed with IFN- $\gamma$  ELISPOT assays to measure  $CD8^+$  and  $CD4^+$  T-cell responses against specific recall responses at baseline, 4 weeks and 8 weeks posttreatment initiation (Fig. 3A). Figures 3B and C depict the magnitude and specificity of the  $CD8^+$  and  $CD4^+$  IFN- $\gamma$  responses in response to the pooled 43 CEF peptide library, or inactivated TT, respectively. The results showed that the frequency and functionality of CEF-specific  $CD8^+$  T cells was not significantly altered throughout the course of the treatment (Fig. 3B). Although some patients showed variability in the ratio of IFN- $\gamma$  spots per million PBMCs between cycle 1 or 2 versus baseline, as a group, the changes were not significant ( $P = 0.25$ ). Likewise, the frequency and functionality of TT-specific  $CD4^+$  T cells was not significantly affected over the course of treatment ( $P > 0.2$ ; Fig. 3C). Collectively, these



**Figure 2.** The V600 mutant BRAF inhibitor treatment does not affect the number of  $CD4^+/CD8^+$  T, B, or NK cells in the blood of patients treated. PBMCs of 13 patients with tumors bearing the V600E BRAF mutation were treated with the GSK2118436 V600 mutant BRAF inhibitor for 2 cycles (4 weeks for each cycle). PBMCs before treatment (baseline) and after cycles 1 and 2 were stained with CD3-AmCyan, CD19-FITC, CD8-APC, CD56-PE-Cy7, and CD4-APC-Cy7. The number of  $CD8^+$  T cells (A),  $CD4^+$  T cells (B), B cells (C), or NK cells (D) per microliter of blood did not change throughout the course of treatment. The frequency of these cell types also stayed constant throughout the treatment (Supplementary Fig. S2).

Figure 3. The V600 mutant BRAF inhibitor treatment does not affect CD4<sup>+</sup> and CD8<sup>+</sup> memory T-cell responses to recall antigens. GSK2118436 V600 mutant BRAF inhibitor-treated patients were tested *in vitro* before treatment (baseline) and after cycles 1 and 2 in an IFN- $\gamma$  ELISPOT assay in the presence of a CEF (CMV/EBV/Flu) peptide library containing 43 peptides, chemically inactivated TT, a positive (PMA/ionomycin), or negative controls (no peptide; A). CD8<sup>+</sup> T-cell memory responses against CEF peptides (B) and CD4<sup>+</sup> T-cell memory responses against a chemically inactivated TT protein (C) were measured and represented as fold change from baseline.




results show that treatment with the V600 mutant BRAF inhibitor GSK2118436 has no significant negative impact on recall responses by both CD8<sup>+</sup> and CD4<sup>+</sup> T cells.

#### Mutant BRAF inhibition with GSK2118436 increased the cell number (and frequency) of tumor antigen-specific T cells in a subset of treated patients' peripheral blood

In light of recent reports describing BRAF(V600E)-mediated upregulation of melanocyte differentiation antigens (MDA) expression in melanoma cell lines (33), we were interested to determine whether V600 mutant BRAF inhibition would impact the cell number and frequency of tumor antigen-specific CD8<sup>+</sup> T cells in the peripheral blood of treated patients. To accomplish this, we screened all 13 study patients for expression of HLA-A2 and identified 4 such patients. To monitor the frequencies of tumor antigen-specific CD8<sup>+</sup> T cells, we stained with tetramers, pen-

tamers, or dextramers that were specific for HLA-A2-restricted epitopes derived from gp100, MART-1, tyrosinase, survivin, and NY-ESO1, and quantified their cell number and frequencies by flow cytometric analysis and the absolute lymphocyte count. The results showed that while most of the tumor antigen-specific CD8<sup>+</sup> T cells levels screened were undetectable, 3 of the 4 patients did have detectable melanoma antigen-specific CD8<sup>+</sup> T cells in their blood. Figure 4A shows the different tumor-specific multimer staining that show an increase in frequency with GSK2118436 treatment. The insert in each FACS plot shows the staining of that sample with a control multimer. As shown in Fig. 4B, 2 patients (001 and 014) showed discreet cell number increases in NY-ESO1-specific CD8<sup>+</sup> T cells following GSK2118436 treatment. In both patients, these cells were not detectable before therapy, suggesting that they were generated *de novo*, potentially in response to tumor destruction and subsequent tumor antigen release.



**Figure 4.** The frequency of tumor-associated antigen (TAA)-specific CD8<sup>+</sup> T cells increases in the PBMCs of HLA-A2<sup>+</sup> patients after treatment with V600 mutant BRAF inhibitor. Four of the 13 patients were positive for the HLA-A2 allele, as determined by flow cytometry (patients 001, 014, 021, and 035). Patients were screened with HLA-A2–restricted tetramers (gp100, tyrosinase), dextramers (MART-1, NY-ESO1), or pentamer (survivin). From the 4 HLA-A2<sup>+</sup> patients, 3 had an increase of tumor antigen-specific T cells (A). FACS plots from the 3 patients having an increase of tumor antigen-specific CD8<sup>+</sup> T cells are shown (A) together with a negative control multimer staining (insert in each plot). Two had an increase of the number of NY-ESO1 dextramer<sup>+</sup> CD8<sup>+</sup> T cells [NY-ESO1<sub>157–165</sub> (165V) per milliliter of blood: patients 001 and 014; B]. Patient 001 also showed an increase of the number of MART-1 dextramer<sup>+</sup> (MART-1<sub>26–35</sub>) CD8<sup>+</sup> T cells per milliliter of blood (C). Patient 021 showed an increase of the number of survivin pentamer<sup>+</sup> (survivin<sub>96–104</sub>(96M)) CD8<sup>+</sup> T cells per milliliter of blood (D).

Interestingly, patient 001 also showed a modest increase in MART-1-specific CD8<sup>+</sup> T-cell number between pretreatment baseline levels and the end of cycle 1 (Fig. 4C). In addition, we found moderately amplified CD8<sup>+</sup> T-cell response against survivin in patient 021, when baseline levels were compared with those at the end of cycle 2 (Fig. 4D). Collectively, these results suggest that the *de novo* generation of tumor antigen-specific immunity is not

impaired by the systemic presence of the V600 mutant BRAF-specific inhibitor GSK2118436.

## Discussion

It is now well established that immunotherapeutic interventions can extend survival in a subset of melanoma cancer patients. Such approaches include IL-2 therapy, adoptive T-cell transfer, and antibodies that target T cell coinhibitory

molecules CTLA-4 and PD-1, all of which have been shown to have varying degrees of clinical success (22, 36). However, strong evidence has been accumulating that the immune system can make a critical contribution to additional antitumor responses even in the context of nonimmunotherapeutic treatments (37, 38). For example, different chemotherapeutic agents can either enhance or inhibit antitumor immunity, depending on whether they induce immunogenic or nonimmunogenic forms of tumor cell death, and how harmful the agents are to immune cells *in vivo* (39, 40). The emerging paradigm is that an intact immune system contributes significantly to the outcome of treatment and may be important for clearance of drug-resistant tumor cells and for prevention of recurrence (41).

The emergence of kinase inhibitors that can target oncogenic signaling pathways in tumors and induce tumor growth arrest and/or apoptosis has provided another very promising approach to fighting cancer (42). Early generations of tyrosine kinase inhibitors included dasatinib, sorafenib, and imatinib. Although these agents can induce objective clinical responses in some cancer patients, they also showed severe off-target cytotoxicity to immune cells that resulted in lymphopenia and an increased frequency of pathogen infections (30, 43, 44). These immunosuppressive effects were likely due to the lack of specificity that characterized the early generations of kinase inhibitors that may have resulted in off-target MAPK pathway inhibition compromising the function and survival of lymphocytes and other immune cells (45, 46).

More recent generations of kinase-targeted agents have shown a much higher degree of specificity, resulting in more effective pathway-specific inhibition as well as a greater avidity for the mutated forms of oncogenic kinases found only in tumor cells. Inhibitors that specifically target mutated BRAF(V600E) have shown remarkable efficacy in the clinic and in phase II/III trials, inducing responses in most melanoma patients harboring the V600E mutation (5). The therapeutic window of these specific inhibitors is much higher than previous generations of kinase inhibitors, showing comparatively little toxicity and few off-target effects *in vivo* at doses that are clinically effective (35, 47). However, despite these encouraging results, responses to BRAF(V600E) inhibition are relatively short lived, and disease recurrence of inhibitor-resistant tumors occurs in nearly all treated patients (20).

The development of effective immunotherapeutic interventions and more effective kinase inhibitors have led to the idea of combining these therapies, with the rationale being that because the 2 treatments each use different cytotoxic mechanisms, combination therapies will show additive effects and will increase the duration of responses (28). Interestingly, some studies have shown that constitutive activation of the MAPK pathway by BRAF(V600E) induces the downstream production of immunosuppressive cytokines IL-6, IL-10, and VEGF (16). Furthermore, another study showed that BRAF(V600E) inhibition leads to the upregulation of MDA, leading to increased recognition and killing of tumor cells by MDA-

specific CTL (33). This implied connection between oncogenic BRAF signaling and immune suppression suggests that combination therapies may, in fact, be synergistic, provided that BRAF(V600E) inhibitors are nontoxic to immune cells. Two recent studies are consistent with this notion, both showing that BRAF(V600E) inhibition does not significantly affect T-cell viability, proliferation, or function at physiologically relevant concentrations *in vitro*, in contrast to earlier generations of less specific MAPK pathway inhibitors (33, 34).

One caveat of these *in vitro* studies is that although serum concentrations of BRAF inhibitors can be accurately measured in pharmacokinetic studies in treated patients, much of the drug may not be biologically active due to being bound to serum proteins. Therefore, it is also critical to test immune function in patients exposed to inhibitors, as we have done in this study. By carrying out immune monitoring on BRAF inhibitor-treated patients before and during therapy, we show that GSK2118436 has negligible effects on preexisting immunity and likely does not negatively impact *de novo* antitumor immune responses, and on the contrary, would seem to enhance it, as shown by the increase of tumor antigen-specific T cells. For the most part, serum cytokine levels, immune cell subsets, and memory CD4<sup>+</sup> and CD8<sup>+</sup> T-cell function remained unaffected. In fact, there was a small but significant increase in the serum concentration of TNF- $\alpha$ , and some patients even experienced an increase in melanoma antigen-specific CD8<sup>+</sup> T cells. In addition, the frequency of apoptotic immune cells in the blood was not affected during the course of treatment (data not shown). Although these observations do not provide strong evidence of an enhanced immune response, they do support the notion that specific BRAF(V600E) targeting does not harm the immune system, unlike many other more nonspecific kinase-targeted agents. One weakness to this study was the analysis was only conducted in peripheral blood, and not in pre- and on-treatment tumor biopsies. Additional studies on such biopsy samples or on TIL generated from pre- and on-treatment lesions is warranted.

Because immunotherapy and BRAF inhibition each induce substantial but incomplete clinical responses in melanoma patients, it is hoped that future therapies combining these 2 approaches will result in higher response rates. We found that BRAF(V600E) kinase inhibitors such as GSK2118436 do not adversely affect the frequencies of different lymphocyte and antigen-presenting cell subsets in the blood and do not affect serum cytokine levels. Moreover, we found some evidence of increased melanoma antigen-specific T cells in some patients during therapy. These results suggest that BRAF(V600E) inhibition is well poised as a combinatorial agent with immune therapies to boost antitumor responses. Carrying out immune monitoring studies will be critical to analyze how the 2 therapies work together *in vivo*; in particular, to determine whether such combinations prove to be additive in their effects, which could result in extension of patient survival, or whether they are synergistic, in which case a higher rate of clinical benefit may conceivably be achieved.



### Disclosure of Potential Conflicts of Interest

D.S. Hong is an employee and has received a commercial research grant from GlaxoSmithKline; he has an ownership interest and is also a shareholder of GlaxoSmithKline. G.S. Falchook and R. Kurzrock have received a commercial research grant from GlaxoSmithKline. V.L. Goodman has ownership interest (including patents) from GlaxoSmithKline. J.J. Legos and S.C. Blackman have employment as Director of Cancer Research in GlaxoSmithKline. No potential conflicts of interest were disclosed by the other authors.

### Acknowledgments

The authors thank Joann Aaron for editing the manuscript.

### Grant Support

This work was supported by local protocol #: LAB09-0114. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 29, 2011; revised January 9, 2012; accepted January 23, 2012; published OnlineFirst February 21, 2012.

### References

- Rosenberg SA, Restifo NP, Yang JC, Morgan RA, Dudley ME. Adoptive cell transfer: a clinical path to effective cancer immunotherapy. *Nat Rev Cancer* 2008;8:299–308.
- Sampson JH, Carter JH Jr., Friedman AH, Seigler HF. Demographics, prognosis, and therapy in 702 patients with brain metastases from malignant melanoma. *J Neurosurg* 1998;88:11–20.
- Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *CA Cancer J Clin* 2010;60:277–300.
- Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010;363:711–23.
- Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med* 2010;363:809–19.
- Spagnoli GC, Ebrahimi M, Iezzi G, Mengus C, Zajac P. Contemporary immunotherapy of solid tumors: from tumor-associated antigens to combination treatments. *Curr Opin Drug Discov Devel* 2010;13:184–92.
- Agarwala SS. Novel immunotherapies as potential therapeutic partners for traditional or targeted agents: cytotoxic T-lymphocyte antigen-4 blockade in advanced melanoma. *Melanoma Res* 2010;20:1–10.
- Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the BRAF gene in human cancer. *Nature* 2002;417:949–54.
- Oikonomou E, Pintzas A. Cancer genetics of sporadic colorectal cancer: BRAF and PI3KCA mutations, their impact on signaling and novel targeted therapies. *Anticancer Res* 2006;26:1077–84.
- Lanzilotta SG, Grammatica L, Paradiso A, Simone G. BRAF in papillary thyroid carcinoma. *Cell Oncol* 2007;29:269–77.
- Singh M, Lin J, Hocker TL, Tsao H. Genetics of melanoma tumorigenesis. *Br J Dermatol* 2008;158:15–21.
- Dong J, Phelps RG, Qiao R, Yao S, Benard O, Ronai Z, et al. BRAF oncogenic mutations correlate with progression rather than initiation of human melanoma. *Cancer Res* 2003;63:3883–5.
- Sumimoto H, Miyagishi M, Miyoshi H, Yamagata S, Shimizu A, Taira K, et al. Inhibition of growth and invasive ability of melanoma by inactivation of mutated BRAF with lentivirus-mediated RNA interference. *Oncogene* 2004;23:6031–9.
- Dankort D, Curley DP, Carlidge RA, Nelson B, Karnezis AN, Damsky WE Jr, et al. Braf(V600E) cooperates with Pten loss to induce metastatic melanoma. *Nat Genet* 2009;41:544–52.
- Zheng B, Jeong JH, Asara JM, Yuan YY, Granter SR, Chin L, et al. Oncogenic B-RAF negatively regulates the tumor suppressor LKB1 to promote melanoma cell proliferation. *Mol Cell* 2009;33:237–47.
- Sumimoto H, Imabayashi F, Iwata T, Kawakami Y. The BRAF-MAPK signaling pathway is essential for cancer-immune evasion in human melanoma cells. *J Exp Med* 2006;203:1651–6.
- Dummer R, Robert C, Chapman P, et al. AZD6244 (ARR-142886) vs temozolomide (TMZ) in patients with advanced melanoma: an open-label, randomized, multicenter, phase II study. *J Clin Oncol* 2008;26:9033.
- Flaherty KT, Puzanov J, Sosman J, et al. Phase I study of PLX4032: proof of concept for V600E BRAF mutation as a therapeutic target in human cancer. *J Clin Oncol* 2009;27:9000.
- Nazarian R, Shi H, Wang Q, Kong X, Koya RC, Lee H, et al. Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature* 2010;468:973–7.
- Solitt DB, Rosen N. Resistance to BRAF inhibition in melanomas. *N Engl J Med* 2011;364:772–4.
- Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 2011;364:2507–16.
- Weber J. Immunotherapy for melanoma. *Curr Opin Oncol* 2011;23:163–9.
- Atkins MB, Lotze MT, Dutcher JP, Fisher RI, Weiss G, Margolin K, et al. High-dose recombinant interleukin 2 therapy for patients with metastatic melanoma: analysis of 270 patients treated between 1985 and 1993. *J Clin Oncol* 1999;17:2105–16.
- Rosenberg SA, Dudley ME. Adoptive cell therapy for the treatment of patients with metastatic melanoma. *Curr Opin Immunol* 2009;21:233–40.
- Palucka K, Ueno H, Fay J, Banchereau J. Dendritic cells and immunity against cancer. *J Intern Med* 2011;269:64–73.
- Hodi FS. Overcoming immunological tolerance to melanoma: targeting CTLA-4. *Asia Pac J Clin Oncol* 2010;6 Suppl 1:S16–23.
- Brahmer JR, Drake CG, Wollner I, Powderly JD, Picus J, Sharfman WH, et al. Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates. *J Clin Oncol* 2010;28:3167–75.
- Shada AL, Molhoek KR, Slingluff CL Jr. Interface of signal transduction inhibition and immunotherapy in melanoma. *Cancer J* 2010;16:360–6.
- Fraser CK, Blake SJ, Diener KR, Lyons AB, Brown MP, Hughes TP, et al. Dasatinib inhibits recombinant viral antigen-specific murine CD4+ and CD8+ T-cell responses and NK-cell cytolytic activity *in vitro* and *in vivo*. *Exp Hematol* 2009;37:256–65.
- Weichsel R, Dix C, Wooldridge L, Clement M, Fenton-May A, Sewell AK, et al. Profound inhibition of antigen-specific T-cell effector functions by dasatinib. *Clin Cancer Res* 2008;14:2484–91.
- Dietz AB, Souan L, Knutson GJ, Bulur PA, Litzow MR, Vuk-Pavlovic S. Imatinib mesylate inhibits T-cell proliferation *in vitro* and delayed-type hypersensitivity *in vivo*. *Blood* 2004;104:1094–9.
- Zhao W, Gu YH, Song R, Qu BQ, Xu Q. Sorafenib inhibits activation of human peripheral blood T cells by targeting LCK phosphorylation. *Leukemia* 2008;22:1226–33.
- Boni A, Cogdill AP, Dang P, Udayakumar D, Njauw CN, Sloss CM, et al. Selective BRAFV600E inhibition enhances T-cell recognition of melanoma without affecting lymphocyte function. *Cancer Res* 2010;70:5213–9.
- Comin-Anduix B, Chodon T, Sazegar H, Matsunaga D, Mock S, Jalil J, et al. The oncogenic BRAF kinase inhibitor PLX4032/RG7204 does not affect the viability or function of human lymphocytes across a wide range of concentrations. *Clin Cancer Res* 2010;16:6040–8.
- Kefford R, Arkenau HT, Brown MP, et al. Phase 1/2 study of GSK2118436, a selective inhibitor of oncogenic mutant BRAF kinase in patients with metastatic melanoma and other solid tumors [abstract]. In: Proceedings of the 2010 ASCO Annual Meeting; Chicago, IL: ASCO; 2010. Abstract nr 8503.

36. Jorritsma A, Schumacher TN, Haanen JB. Immunotherapeutic strategies: the melanoma example. *Immunotherapy* 2009;1:679–90.
37. Dhomen N, Da Rocha Dias S, Hayward R, Ogilvie L, Hedley D, Delmas V, et al. Inducible expression of (V600E) Braf using tyrosinase-driven Cre recombinase results in embryonic lethality. *Pigment Cell Melanoma Res* 2010;23:112–20.
38. Green DR, Ferguson T, Zitvogel L, Kroemer G. Immunogenic and tolerogenic cell death. *Nat Rev Immunol* 2009;9:353–63.
39. Kulke MH, Vance EA. *Pneumocystis carinii* pneumonia in patients receiving chemotherapy for breast cancer. *Clin Infect Dis* 1997;25:215–8.
40. Kepp O, Tesniere A, Zitvogel L, Kroemer G. The immunogenicity of tumor cell death. *Curr Opin Oncol* 2009;21:71–6.
41. Zitvogel L, Apetoh L, Ghiringhelli F, Andre F, Tesniere A, Kroemer G. The anticancer immune response: indispensable for therapeutic success? *J Clin Invest* 2008;118:1991–2001.
42. Baker SJ, Reddy EP. Targeted inhibition of kinases in cancer therapy. *Mt Sinai J Med* 2010;77:573–86.
43. Schutz FA, Je Y, Choueiri TK. Hematologic toxicities in cancer patients treated with the multi-tyrosine kinase sorafenib: a meta-analysis of clinical trials. *Crit Rev Oncol Hematol* 2010;80:291–300.
44. Rohon P, Porkka K, Mustjoki S. Immunoprofiling of patients with chronic myeloid leukemia at diagnosis and during tyrosine kinase inhibitor therapy. *Eur J Haematol* 2010;85:387–98.
45. Hartmann JT, Haap M, Kopp HG, Lipp HP. Tyrosine kinase inhibitors - a review on pharmacology, metabolism and side effects. *Curr Drug Metab* 2009;10:470–81.
46. Farber DL. Biochemical signaling pathways for memory T cell recall. *Semin Immunol* 2009;21:84–91.
47. Arkenau HT, Kefford R, Long GV. Targeting BRAF for patients with melanoma. *Br J Cancer* 2010;104:392–8.

## Correction: BRAF(V600) Inhibitor GSK2118436 Targeted Inhibition of Mutant BRAF in Cancer Patients Does Not Impair Overall Immune Competency

In this article (Clin Cancer Res 2012;18:2326–35), which was published in the April 15, 2012, issue of *Clinical Cancer Research* (1), there were several errors in the Disclosure of Potential Conflicts of Interest section. Dr. Hong is not an employee of GlaxoSmithKline, nor does he hold stock or ownership in the company. The author regrets these errors.

### Reference

1. Hong DS, Vence L, Falchook G, Radvanyi LG, Liu C, Goodman V, et al. BRAF(V600) inhibitor GSK2118436 targeted inhibition of mutant BRAF in cancer patients does not impair overall immune competency. Clin Cancer Res 2012;18:2326–35.

Published OnlineFirst May 25, 2012.

doi: 10.1158/1078-0432.CCR-12-1343

©2012 American Association for Cancer Research.

# Clinical Cancer Research

## **BRAF(V600) Inhibitor GSK2118436 Targeted Inhibition of Mutant BRAF in Cancer Patients Does Not Impair Overall Immune Competency**

David S. Hong, Luis Vence, Gerald Falchook, et al.

*Clin Cancer Res* 2012;18:2326-2335. Published OnlineFirst February 21, 2012.

<b>Updated version</b>	Access the most recent version of this article at: doi: <a href="https://doi.org/10.1158/1078-0432.CCR-11-2515">10.1158/1078-0432.CCR-11-2515</a>
<b>Supplementary Material</b>	Access the most recent supplemental material at: <a href="http://clincancerres.aacrjournals.org/content/suppl/2012/02/21/1078-0432.CCR-11-2515.DC1">http://clincancerres.aacrjournals.org/content/suppl/2012/02/21/1078-0432.CCR-11-2515.DC1</a>

<b>Cited articles</b>	This article cites 46 articles, 9 of which you can access for free at: <a href="http://clincancerres.aacrjournals.org/content/18/8/2326.full#ref-list-1">http://clincancerres.aacrjournals.org/content/18/8/2326.full#ref-list-1</a>
<b>Citing articles</b>	This article has been cited by 17 HighWire-hosted articles. Access the articles at: <a href="http://clincancerres.aacrjournals.org/content/18/8/2326.full#related-urls">http://clincancerres.aacrjournals.org/content/18/8/2326.full#related-urls</a>

<b>E-mail alerts</b>	<a href="#">Sign up to receive free email-alerts</a> related to this article or journal.
<b>Reprints and Subscriptions</b>	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at <a href="mailto:pubs@aacr.org">pubs@aacr.org</a> .
<b>Permissions</b>	To request permission to re-use all or part of this article, use this link <a href="http://clincancerres.aacrjournals.org/content/18/8/2326">http://clincancerres.aacrjournals.org/content/18/8/2326</a> . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.