

A Randomized, Multicenter, Placebo-Controlled Clinical Trial of Racotumomab-Alum Vaccine as Switch Maintenance Therapy in Advanced Non-Small Cell Lung Cancer Patients

Sailyn Alfonso¹, Anet Valdés-Zayas², Eduardo R. Santiesteban³, Yoanna I. Flores⁴, Fernando Areces⁴, Maurenis Hernández², Carmen E. Viada², Ivis C. Mendoza⁵, Pedro P. Guerra⁵, Elena García⁵, Ramón A. Ortiz¹, Ana V. de la Torre¹, Meylán Cepeda¹, Kirenía Pérez², Eric Chong², Ana María Hernández², Darien Toledo², Zuyén González², Zaima Mazorra², Tania Crombet², Rolando Pérez², Ana María Vázquez², and Amparo E. Macías²

Abstract

Purpose: Racotumomab-alum is an anti-idiotypic vaccine targeting the NeuGcGM3 tumor-associated ganglioside. This clinical trial was conducted to provide a preliminary estimate of efficacy and safety of racotumomab as switch maintenance for patients with advanced non-small cell lung cancer (NSCLC).

Experimental design: Patients with stage IIIb/IV NSCLC who have at least stable disease after first-line chemotherapy were randomized 1:1 to racotumomab-alum (5 immunizations every 2 weeks and re-immunizations every 4 weeks) or placebo. Treatment was administered beyond progressive disease, until severe performance status worsening or toxicity. At progression, only five patients per group received further anticancer therapy. The primary endpoint was overall survival (OS).

Results: One-hundred and seventy-six patients were randomized to racotumomab-alum ($n = 87$) and placebo ($n = 89$). Median OS was 8.23 and 6.80 months, respectively [HR, 0.63; 95% confidence interval (CI), 0.46–0.87; $P = 0.004$]. Median progression-free survival (PFS) in vaccinated patients was 5.33 versus 3.90 months for placebo (HR, 0.73; 95% CI 0.53–0.99; $P = 0.039$). The most common adverse events in the racotumomab-alum arm were burning and pain at the injection site, bone pain, and asthenia. A high antibody response of IgM and IgG isotype against the NeuGcGM3 ganglioside was obtained. Hyperimmune sera were able to specifically recognize and kill the NeuGcGM3-expressing L1210 cell line. Patients who developed anti-NeuGcGM3 antibodies capable to bind and kill $\geq 30\%$ L1210 cells showed longer median survival times.

Conclusions: Switch maintenance with racotumomab-alum is an effective and a well-tolerated treatment option for patients with advanced NSCLC. *Clin Cancer Res*; 20(14); 3660–71. ©2014 AACR.

Introduction

Lung cancer is the most common cause of cancer-related death worldwide (1). Non-small cell lung cancer (NSCLC) constitutes the predominant histologic type among the lung neoplasias, representing more than 75% of all lung tumors.

A high percentage of the patients are diagnosed at locally advanced stage or with metastatic disease (2). Unfortunately, little progress has been achieved in the overall survival (OS) of patients with advanced NSCLC after first-line chemotherapy, as various chemotherapy combinations have not improved the results obtained with platinum doublets (3).

Recently, the role of maintenance treatment for patients with advanced NSCLC with favorable response [complete response (CR), partial response (PR), or stable disease (SD)] after first-line chemotherapy has been evaluated by an international group of experts. On the basis of the available evidence, the expert panel agreed that maintenance therapy represents a treatment option in advanced NSCLC, in any of its modalities: continuation or switch, the latter with a different chemotherapy or targeted agent (4). Most trials showed significant prolongation of progression-free survival (PFS; refs. 5–14), whereas some studies also showed extension of the OS, particularly after the use of pemetrexed and erlotinib (6, 7, 9, 14, 15).

Authors' Affiliations: ¹Celestino Hernández Robau Hospital, Villa Clara; ²Center of Molecular Immunology; ³José R. López Tabranes Hospital, Matanzas; ⁴National Institute of Oncology and Radiobiology; and ⁵National Coordinating Center of Clinical Trials

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

S. Alfonso and A. Valdés-Zayas contributed equally to this article.

Corresponding Author: Amparo E. Macías, Department of Clinical Investigations, Center of Molecular Immunology; 216 St and 15 Ave., Atabey, Playa, Havana 11600, Cuba. Phone: 53-7-2713146; Fax: 53-7-2720644; E-mail: amparo@cim.sld.cu

doi: 10.1158/1078-0432.CCR-13-1674

©2014 American Association for Cancer Research.

Translational Relevance

In advanced non-small cell lung cancer (NSCLC), there have been few improvements in outcome after first-line chemotherapy as new combinations of cytotoxic drugs have not enhanced the results obtained just with platinum doublets. Thus, the development of new more effective targeted strategies continues to be a priority. Here, we present the clinical and immunologic results from a randomized study of racotumomab-alum vaccine, targeting NeuGcGM3 ganglioside, as switch maintenance therapy in patients with advanced NSCLC. A clinical meaningful improvement in overall survival (OS) and progression-free survival for racotumomab-alum versus placebo is shown. The capacity of IgM anti-NeuGcGM3 antibodies to bind 30% or more of NeuGcGM3-expressing L1210 cells and to kill that percentage of cells was associated with longer OS. To our knowledge, this is the first report showing an association between the active induction of cytotoxic NeuGcGM3-specific antibodies and the clinical outcome of patients with NSCLCs treated with racotumomab-alum vaccine.

Active immunotherapy is one of the most promising fields in cancer research and several strategies have been pursued to design cancer vaccines. One approach to generate an effective immune response against tumor-associated antigens involves the use of anti-idiotypic monoclonal antibodies (Ab2 mAb). Anti-idiotypic antibodies have proved to mimic the nominal antigens and induce antigen-specific responses, even against non-protein tumor-associated antigens like gangliosides (16, 17).

Neu glycolyl (NeuGc)—containing gangliosides are attractive targets for immunotherapy. These glycolipids are not present in the cytoplasmic membrane of humans (18), after a 92-bp deletion in the gene that encodes the cytidine-monophosphate-*N*-acetylneuraminic acid hydroxylase (*cmah*), which catalyzes the conversion of *N*-acetyl to NeuGc sialic acid (19, 20). However, NeuGc—containing gangliosides have been detected in several human tumors, including NSCLC (21–24). NeuGcGM3 is besides a potent immunosuppressive molecule (25).

We previously reported a vaccine preparation composed of a murine anti-idiotypic mAb that mimics NeuGcGM3 ganglioside. This Ab2, named 1E10 and later racotumomab (26), was generated from immunizing BALB/c mice with P3 mAb, an Ab1 antibody that recognizes NeuGc gangliosides (27–30). Racotumomab vaccine was able to generate a strong antitumor activity in mice bearing melanoma or breast carcinomas (26) and to induce antimetastatic effect in 3LL-D122 Lewis Lung carcinoma, a poorly immunogenic and highly metastatic murine model in C57BL/6 mice. Therapeutic effect was associated with an increase of T cells infiltrating the metastases, a reduction of blood vessels, and an increase of apoptotic cells in the lung metastases (31). Three different toxicology studies evaluating acute toxicity,

repeated toxicity, and local tolerance were done. A mutagenic study was also performed. Racotumomab was demonstrated to be safe and non-mutagenic.

Several phase I trials using racotumomab-alum vaccine were conducted in patients with advanced melanoma, breast, and lung cancer. Vaccine was safe and immunogenic as high antibody response to NeuGcGM3 was detected in vaccinated patients (28–30, 32, 33). In addition, NeuGcGM3-specific IFN γ -secreting T cells were detected by ELISPOT in patients with breast cancer vaccinated with racotumomab (34). Although these studies were not designed to evaluate the therapeutic efficacy of the vaccine, a prolonged survival was observed in several patients. In particular, encouraging results were obtained when racotumomab-alum vaccine was evaluated in patients with NSCLC (30). A strong IgM and IgG antibody response against NeuGcGM3 was elicited in patients with advanced NSCLC. Patients' hyperimmune sera were able to specifically recognize and induce cell death of the NeuGcGM3-expressing X63 myeloma cells. Remarkably, patients who developed IgM or IgG antibodies against NeuGcGM3 showed longer median survival time (33).

To assess the efficacy and tolerability of switch maintenance therapy with racotumomab-alum in patients with advanced NSCLC, we conducted this randomized, double-blind, placebo-controlled phase II/III clinical trial.

Patients and Methods

Patient's selection

Eligible patients were 18 years or older, with histologically or cytologically confirmed stage IIIB–IV NSCLC (6th edition AJCC staging system), who had achieved CR, PR, or SD after the standard first-line therapy. Random assignment, 87 patients in the vaccine group and 89 in the placebo group, was performed at least 28 days after completing first-line treatment, but within 2 months. All patients had measurable disease, an Eastern Cooperative Oncology Group performance status (ECOG PS) \leq 2 (35), adequate renal, hepatic, and hematologic functions.

Exclusion criteria included patients who had received immunotherapy or other investigational drugs; patients with known hypersensitivity to any component of the formulation; patients who were pregnant or lactating; patients with uncontrolled chronic diseases, history of severe allergic reactions; patients with brain metastases or other primary neoplastic lesion; patients with active infections, symptomatic congestive heart failure, unstable angina, cardiac arrhythmia or psychiatric disorders; and patients receiving systemic corticosteroids at the time of inclusion and patients with positive serology for hepatitis B and C or HIV.

Study design

This double-blinded, placebo-controlled clinical trial was conducted at 3 hospitals from Villa Clara, Matanzas, and Havana, Cuba. Sample size was estimated considering that 160 patients, or 103 events, would achieve 81% power to

detect a hazard rate of 0.56 when the proportions of survivors in each group are 0.25 and 0.46 at a significance level of 0.05, using a 2-sided log-rank test. About 10% patients' loss was anticipated, and accordingly, the final sample size was 176 patients. The projected accrual period was 3 years. An interim analysis on futility was planned when 52 events occurred. The primary objective was to evaluate the effect of racotumomab-alum vaccine in OS of patients with advanced NSCLC. The secondary objectives were to evaluate PFS, immunologic response, and safety.

Eligible patients were randomized 1:1 to receive racotumomab-alum (group A) or placebo (group B). Randomization was not stratified but it was balanced according to gender, clinical stage, performance status, race, and response to first-line treatment.

Patients were injected intradermally and received 15 doses of 1 mg of racotumomab-alum or placebo. The induction phase consisted of 5 doses administered every 2 weeks. After induction, patients were vaccinated every 4 weeks, for 1 year (10 doses). After treatment was completed (15 doses of vaccine or placebo), the blinding was opened, and only patients in the racotumomab arm continued vaccination every 4 weeks, even beyond progression. Criteria for stopping vaccination included voluntary withdrawal, unacceptable toxicity, or severe worsening of the patient's PS. Patients who discontinued study treatment were followed until death or study termination. Concurrent antitumor therapy was not permitted.

An independent CRO (CENCEC) was responsible for conducting the study. Treatment allocation was done using the automatic MINIM system. To balance both study groups, the investigator provided the information on patient sex, race, clinical stage, PS, and response to first-line treatment before randomization. The trial was blinded and the investigational drug and placebo had the same organoleptic characteristics and were presented in the same primary and secondary containers. A sealed letter enclosing the information of the treatment group was sent together with the investigational product. Information on treatment was disclosed only after completing 15 doses or in case of severe or serious adverse events.

The trial was approved by the local ethic review boards and the Cuban Regulatory Agency. All patients provided written informed consent before study participation. The trial was conducted in accordance to the principles of the Declaration of Helsinki and Good Clinical Practice guidelines. It is registered at the Cuban Registry of Clinical Trials (www.registroclinico.sld.cu; ID: RPCEC00000009).

Efficacy assessments

OS was defined as the time elapsed from randomization to death from any cause. PFS was calculated from the date of randomization up to the date of documented progression or death from any cause.

Patient assessment was performed at baseline and every 4 weeks. Baseline evaluation included medical and smoking history, physical examination, vital signs, PS, complete blood cell count, and blood biochemistry. Toxicity was

evaluated according to the Common Terminology Criteria for Adverse Events (version 3.0), by physical examination and clinical laboratory tests, performed during treatment and follow-up. Tumors were measured by X-rays and CT scans at baseline and then every 3 months. The RECIST 1.0 guideline (36) was used to classify tumor response. After disease progression, patients could receive any subsequent therapy at the discretion of the treating physician.

Anti-idiotypic racotumomab mAb

Active drug was composed of the anti-idiotypic antibody racotumomab, alum hydroxyl gel, mono- and dibasic sodium phosphatase salt, sodium chloride, and water for injection. The placebo had the same composition excluding racotumomab. Racotumomab mAb was purified from mouse ascites, and the aluminum hydroxide-precipitated vaccine was produced according the Good Manufacturing Practice guidelines. The manufacturing process was certified by the Quality Control Department of the Center of Molecular Immunology (CIM).

Ganglioside and tumor cell lines

NeuGcGM3 ganglioside was purified from horse erythrocytes as described previously (37) and was provided by Dr. Carr (CIM, Havana, Cuba). The L1210 murine lymphocytic leukemia cell line, expressing high levels of NeuGcGM3 ganglioside (38, 39), was obtained from the ATCC. L1210 cells deficient of NeuGc sialoconjugates (L1210 *cmah-kd*), were generated at CIM, as previously explained (40). Cells were grown in DMEM/F12 medium (Gibco-BRL) supplemented with 10% of heat-inactivated fetal bovine serum (FBS) (Gibco-BRL), 2 mmol/L L-glutamine, 25 mmol/L HEPES, penicillin (100 µg/mL), and 100 µg/mL of streptomycin (Invitrogen) and maintained at 37°C with 5% CO₂.

Measurement of antibody response

The trial goal was to evaluate the immune response in 30% of the subjects (53 vaccinated and placebo patients). However, 6 of the 53 patients did not complete the vaccine induction period (5 doses) or did not have at least 3 serial serum samples. Consequently, the immunologic assessment was done in a cohort of 47 patients, 24 vaccinated and 23 controls.

The antibody response against racotumomab mAb and the purified ganglioside NeuGcGM3 was determined by a solid-phase ELISA, as formerly reported (28). The inverse of the highest serum dilution giving optical density (OD) values ≥ 0.25 and twice the value of the pre-immune serum was considered as the antibody titer. Assays were performed in triplicate for each sample and the coefficient of variation (CV) was less than 15%. The ODs of the blanks were less than 0.1.

Antibody-binding assay by flow cytometry

The detection of antibodies binding to wild-type and mutated L1210 in patients' sera was performed as reported before (41). L1210 cells were blocked in PBS containing 1% BSA for 20 minutes on ice. Patients' serum samples,

corresponding to pre-immune and hyperimmune sera from 2 to 6 months of treatment, were diluted 1:10 and incubated with 10^5 cells for 30 minutes on ice. After washing with cold PBS, the cells were incubated for 30 minutes on ice, with FITC-conjugated goat anti human IgM or FITC-conjugated goat anti-human IgG antibodies (Jackson ImmunoResearch Laboratories). Percentage of positive stained cells was determined in a FACScan flow cytometer (Becton Dickinson). The FlowJo program (version 5.7.2) was used to analyze a total of 10^4 cells acquired on every FACS assay. Positive patients had a binding percent $\geq 20\%$, after subtracting the pre-immune to the hyperimmune recognition percent. The L1210 *cmah-kd* cell line, deficient of NeuGc sialoconjugates (40), was used as control of the antibody specificity for the NeuGcGM3 ganglioside.

Induction of cell death

The lytic capacity of patients' sera was tested by incubating the 1:5 diluted sera with 10^5 tumor cells in DMEM/F12 medium supplemented with 1% FBS in 5% CO₂ atmosphere at 37°C, for 3 hours (33, 39, 41). Cell death percent was determined by flow cytometry after the addition of propidium iodide (PI; Sigma-Aldrich) at a final concentration of 10 $\mu\text{g/mL}$. The cytotoxicity was considered positive when the percentage of dead cells was $\geq 20\%$, after subtracting the pre-immune to hyperimmune percent.

To determine whether the binding and cytotoxic effects of serum were specifically mediated by anti-NeuGcGM3 antibodies, L1210 *cmah-kd* cells and L1210 cells cultured with 10 $\mu\text{mol/L}$ of D-threo-1-phenyl-2-decanoylamino-3-morpholino-1-propanol (D-threo-PDMP; Matreya, LLC), an inhibitor of glucosylceramide synthetase (40), were used.

Statistical analyses

For the analyses of OS and PFS, median values and 95% confidence intervals (CI) were estimated by Kaplan-Meier methodology, and 2-sided log-rank tests were used to compare treatment groups. A secondary analysis of OS was performed using the Cox proportional hazards regression model to determine the effect of treatment in different subgroups. This analysis included age, sex, ethnic origin, tumor stage, histology, smoking history, race, ECOG, response to first-line treatment, and the immunologic variables. The χ^2 and Fisher tests were used for evaluating distribution differences between groups. Wilcoxon test was used as a nonparametric test for matched-pairs signed-ranks test. Statistical analyses were done with SPSS program (version 17.0).

Results

Patient characteristics

From September 27, 2006, to June 28, 2010, 176 patients were randomized: 87 to racotumomab-alum and 89 to placebo (Supplementary Fig. S1). Thus, 176 patients were included in the intent-to-treat efficacy analysis. One patient with histologically unconfirmed NSCLC randomized to racotumomab-alum group did not receive treatment and was excluded from the safety analysis. Demographic

and baseline characteristics were similar in both groups (Table 1).

All patients received platinum-based regimens as first-line chemotherapy. These regimens included vinblastine or etoposide combined with cisplatin or carboplatin. One hundred and sixty-nine patients (98.3%) received 4 to 6

Table 1. Demographic and baseline characteristics

	Racotumomab-alum (n = 87)	Placebo (n = 89)
Age, y		
≤ 60	38 (43.7%)	41 (46.1%)
≥ 60	49 (56.3%)	48 (53.9%)
Gender		
Female	28 (32.2%)	30 (33.7%)
Male	59 (67.8%)	59 (66.3%)
ECOG PS		
0	40 (46.0%)	40 (44.9%)
1	45 (51.7%)	45 (50.6%)
2	2 (2.3%)	4 (4.5%)
Race		
White	70 (80.5%)	71 (79.8%)
Afro-Caribbean	11 (12.6%)	13 (14.6%)
Other	6 (6.9%)	5 (5.6%)
Smoking history		
Current smoker	16 (18.45%)	19 (21.3%)
Former smoker	67 (77.0%)	65 (73.0%)
Nonsmoker	4 (4.6%)	5 (5.6%)
Tumor histology		
Squamous cell carcinoma	33 (37.9%)	33 (37.1%)
Adenocarcinoma	25 (28.7%)	31 (34.8%)
Large cell carcinoma	18 (20.7%)	14 (15.7%)
NSCLC NOS	11 (12.6%)	11 (12.4%)
Disease stage		
IIIB	48 (55.2%)	51 (57.3%)
IV	39 (44.8%)	38 (42.7%)
First-line treatment		
Chemotherapy	87 (100%)	89 (100%)
Radiotherapy	50 (57.5%)	51 (57.3%)
First-line chemotherapy		
Platinum compounds	87 (100%)	89 (100%)
Cisplatin/Vinblastine	28 (32.2%)	18 (20.2%)
Cisplatin/Etoposide	6 (6.9%)	5 (5.6%)
Carboplatin/Vinblastine	39 (44.8%)	50 (56.2%)
Carboplatin/Etoposide	11 (12.6%)	13 (14.6%)
Other Platinum doublets	3 (3.5%)	3 (3.4%)
Response to first-line treatment		
CR	2 (2.3%)	5 (5.6%)
PR	38 (43.7%)	51 (57.3%)
SD	47 (54.0%)	33 (37.1%)

NOTE: Data are n (%).

Abbreviation: NOS, not otherwise specified.

chemotherapy cycles as first-line treatment and only 3 patients (1.7%) received less than 4 cycles (1 patient from the vaccine group and 2 from the placebo arm). As part of the first-line treatment, 50 vaccinated patients (57.5%) and 51 patients (57.3%) from the placebo group received radiotherapy.

All patients included in the trial achieved SD, PR, or CR after first-line therapy and were randomized between 4 and 8 weeks after completing treatment, with a median of 1.51 months (vaccine) and 1.43 months (placebo). Median time from randomization to start racotumomab or placebo was 0.46 and 0.43 months, respectively.

After interrupting vaccine or placebo, 5 patients per group (5.6% of all enrolled patients), received further anticancer therapy. Because of treatment unavailability, only 1 patient from the placebo arm received docetaxel, one of the recommended drugs for second-line therapy in NSCLC. The remaining 4 patients in the placebo cohort received carboplatin (2 patients) or etoposide (2 patients). On the other

hand, 4 patients from the racotumomab arm received new cycles of carboplatin, whereas 1 subject was treated with cisplatin/vinblastine.

Efficacy

The intent-to-treat median OS was 8.23 months (95% CI, 5.59–10.87) for the racotumomab-alum group versus 6.80 months (95% CI, 5.77–7.83) for the placebo group. This difference was statistically significant (HR, 0.63; 95% CI, 0.46–0.87; $P = 0.004$; Fig. 1A). Ten patients from the racotumomab arm and 2 patients from the placebo cohort were alive at the moment of the analysis and were consequently censored. The 1- and 2-year survival rates were 40.2% and 18.4% for the racotumomab-alum group compared with 22.5% and 6.7% for the placebo group.

Almost all patient subgroups benefited from therapy. Remarkably patients with stage IIIB NSCLC derived a larger benefit than those with stage IV (Supplementary Fig. S2).




Figure 1. Intent-to-treat OS (A) and PFS (B) in patients with advanced NSCLC receiving racotumomab-alum vaccine or placebo.

Vaccinated patients also had a statistically significant advantage in PFS (HR, 0.73; 95% CI, 0.53–0.99; $P = 0.039$; Fig. 1B). Median intent-to-treat PFS was 5.33 months (95% CI, 3.48–7.19) in the racotumomab-alum group versus 3.90 months (95% CI, 3.53–4.27) in the placebo group. Seven patients from the racotumomab arm and 2 from the placebo arm have not progressed at the moment of this report.

Safety

The distribution of treatment-related adverse events in the 2 arms is shown in Table 2. The most common racotumomab-related adverse events (all grades) were burning (41.9%) and pain at the injection site (33.7%), followed by bone pain (18.6%) and asthenia (16.3%). In the placebo group, the most frequently reported adverse events were burning (39.3%) and pain at the injection site (24.7%), as well as bone pain (19.1%).

All but 4 treatment-related adverse events were classified as mild or moderate. Two racotumomab-attributed adverse events consisting of asthenia and bone pain were classified as grade III–IV, whereas 2 patients in the placebo arm developed grade III–IV bone pain and dyspnea. Even though these adverse events were categorized as related with the investigational drug, they might also be associated with the natural course of the lung disease and its metastasis. This needs to be confirmed in a larger study in patients with advanced NSCLCs.

Overall, racotumomab-alum was very well-tolerated and its safety profile was consistent with the previously described (28–30).

The proportion of all adverse events (regardless causality) was not different between treatment groups (Supplementary

Table S1). One-hundred and fifty-eight (90.3%) patients had at least 1 adverse event of any grade (Supplementary Table S2). Thirty-four patients (39.5%) in the racotumomab group and 31 (34.8%) in the placebo arm experienced serious adverse events (SAE). Four events from each group consisting of dyspnea, asthenia, bone pain, and abscess in right upper limb were classified as related to the investigational drug. No patient in the vaccine group and only one (1.1%) patient in the placebo group was withdrawn from treatment because of an adverse event. The proportion of SAEs was not different between both groups. No adverse events leading to death were related to the treatment.

Immune response induced by racotumomab-alum vaccine

All vaccinated patients developed a high antibody response against racotumomab of IgG isotype. The titers ranged from 1:200 to 1:51,200, with a median of 1:6,400 (Supplementary Table S3). No reactivity to racotumomab mAb was detected in the patient's sera from the placebo group, at the lowest serum dilution tested (1:50; data not shown).

To determine whether antibodies induced after vaccination had the same specificity of P3 mAb (Ab1), pre- and hyperimmune serum samples were tested for the recognition of NeuGcGM3 ganglioside. An antibody response against NeuGcGM3 was generated in all vaccinated patients (Supplementary Table S3). Twenty-three patients (95.8%) developed antibodies of both IgG and IgM isotypes that bind to NeuGcGM3 by ELISA, whereas 1 subject (4.2%) elicited only IgM antibodies. None of the patients developed IgG subclass antibodies alone. Titers of up to 1:51,200 and 1:6,400 for IgM and IgG isotypes, respectively, were

Table 2. Treatment-related adverse events recorded in at least 1.5% in either group

Adverse event	Racotumomab-alum (n = 86)		Placebo (n = 89)	
	All grades	Grade III–IV	All grades	Grade III–IV
Burning in injection site	36 (41.9%)	0 (0%)	35 (39.3%)	0 (0%)
Pain in injection site	29 (33.7%)	0 (0%)	22 (24.7%)	0 (0%)
Bone pain	16 (18.6%)	1 (1.1%)	17 (19.1%)	1 (1.1%)
Cough	7 (8.1%)	0 (0%)	11 (12.4%)	0 (0%)
Dyspnea	5 (5.8%)	0 (0%)	5 (5.6%)	1 (1.2%)
Asthenia	14 (16.3%)	1 (1.2%)	10 (11.2%)	0 (0%)
Anorexia	6 (7.0%)	0 (0%)	7 (7.9%)	0 (0%)
Expectoration	1 (1.2%)	0 (0%)	3 (3.4%)	0 (0%)
Induration	10 (11.6%)	0 (0%)	9 (10.1%)	0 (0%)
Headache	8 (9.3%)	0 (0%)	9 (10.1%)	0 (0%)
Pruritus	9 (10.5%)	0 (0%)	5 (5.6%)	0 (0%)
Fever	8 (9.3%)	0 (0%)	12 (13.5%)	0 (0%)
Increased volume in injection site	9 (10.5%)	0 (0%)	3 (3.4%)	0 (0%)
Local erythema	11 (12.8%)	0 (0%)	11 (12.4%)	0 (0%)
Myalgia	5 (5.8%)	0 (0%)	7 (7.9%)	0 (0%)
Arthralgia	5 (5.8%)	0 (0%)	5 (5.6%)	0 (0%)

NOTE: Data are number of patients (%). Adverse events graded by CTCAE version 3.0.

obtained. No reaction was detected when pre-immune patients' sera were tested.

Recognition of tumor cells expressing NeuGcGM3 ganglioside

Next, we evaluated the ability of patient's sera to recognize the NeuGcGM3 ganglioside in the natural context of tumor cell membranes. Hyperimmune sera from 11 of 24 patients (45.8%) showed significant binding of antibodies of both isotypes (IgM and IgG) to L1210 cells, compared with the pre-immune sera ($P < 0.0001$; Wilcoxon test; Supplementary Table S3 and Fig. 2A and B). Seven (29.2%) and 4 patients (16.7%) showed IgM- or IgG-binding antibodies to L1210 cells. Only the hyperimmune sera from 2 patients (8.3%) were not able to recognize the NeuGcGM3-express-

ing tumor cell line (Supplementary Table S3). To confirm that antibodies specifically recognized NeuGcGM3 ganglioside on the cell surface, the binding of patients' serum samples to L1210 *cmah-kd* cells, devoid of Neu5Gc-sialoconjugates, was assessed. As is shown in Fig. 2A, sera that bind the native L1210 cells were not able to recognize L1210 *cmah-kd* cells, suggesting that the antigen recognized is presumably a Neu-glycosylated molecule. Binding of IgM and IgG antibodies to L1210 cells, corresponding to 2 representative patients, is shown in Fig. 2C and D.

Cytotoxic effect of anti-NeuGcGM3 antibodies induced by vaccination with racotumomab-alum

To study whether the antibodies induced in vaccinated patients were capable not only to recognize but also to kill




Figure 2. Binding of pre-immune and hyperimmune sera from 24 NSCLC vaccinated patients to the NeuGcGM3-expressing cell line L1210. Recognition assay was performed by incubating the 1:10 diluted sera with L1210 cells or with L1210 *cmah-kd* cells, devoid of NeuGc sialoconjugates. The reaction was developed with FITC-conjugated anti-human IgM (A) or IgG (B). Bars indicate median values. A significant increment in the median binding to L1210 of IgM and IgG hyperimmune antibodies as compared with the pre-immune sera was observed ($P < 0.0001$; Wilcoxon test). Binding was affected when sera were incubated with L1210 *cmah-kd* cells, deficient of NeuGc sialoconjugates, confirming that the recognized antigen is presumably a NeuGc glycosphingolipid. C and D, flow cytometric assays from 2 representative patients, showing both isotypes' binding to L1210. The numbers represent the percentage of positive cells after subtracting the binding of pre-immune sera.

the NeuGcGM3-expressing cells, patients' sera were incubated with L1210 cells. Cytotoxicity was evaluated by flow cytometry using the PI exclusion assay. In 18 of 24 patients (75%), a significant increase in PI incorporation was observed ($P < 0.0001$; Wilcoxon test; Supplementary Table S3 and Fig. 3A). Sera cytotoxicity was significantly reduced when hyperimmune sera were incubated with L1210 *cmah-kd* or with L1210 cells treated with D-threo-PDMP, suggesting that the antigen recognized in the tumor cell membrane is a Neu-glycosylated glycosphingolipid (Fig. 3A). Histograms of PI-stained cells from 2 representative patients are shown in Fig. 3B.

Immunological response and clinical outcomes of patients


The individual survival data from the 24 vaccinated patients that were sampled for immune response evaluation are shown in Supplementary Table S3. Interestingly, we found that the percentage of IgM recognition (HR, 0.97; 95% CI, 0.95–0.99; $P = 0.040$) and the cytotoxicity against

L1210 cells (HR, 0.98; 95% CI, 0.96–0.99; $P = 0.019$) were statistically associated with survival (Fig. 4). No association was observed when other immunologic variables were analyzed. Then, aiming to find a surrogate of patients' outcome, we looked for the lower recognition and cell death percentage associated with patients' survival. Better survival was found in patients with IgM antibodies able to recognize at least 20% of L1210 cells (19.30 vs. 7.83 months; HR, 0.30; 95% CI, 0.11–0.83; $P = 0.014$) or kill at least 25% of the same NeuGcGM3-positive cells (19.50 vs. 8.23 months; HR, 0.26; 95% CI, 0.10–0.70; $P = 0.005$). The highest differences were detected using 30% recognition (19.50 vs. 8.23 months; HR, 0.26; 95% CI, 0.10–0.73; $P = 0.006$) and 30% cell death cut-off (19.63 vs. 8.23 months; HR, 0.21; 95% CI, 0.08–0.58; $P = 0.003$; Supplementary Table S4).

Discussion

In this randomized, placebo-controlled study, patients with advanced NSCLC who received racotumomab-alum

Figure 3. Cytotoxicity induced by sera from 24 NSCLC vaccinated patients on the NeuGcGM3-expressing L1210 cell line. Cells were incubated for 3 hours at 37°C and 5% CO₂ atmosphere with 1:5 diluted pre-immune (blue line) and hyperimmune (red line) sera. The percentage of cell death was measured by PI incorporation assay. A, percentage of cell death after incubation of pre-immune and hyperimmune sera with the NeuGcGM3-expressing L1210 cells, with L1210 cells transduced with a CMP-Neu5Ac-neuraminic acid hydroxylase gene-specific shRNA (L1210 *cmah-kd*) or with L1210 cells treated with 10 μmol/L D-threo-PDMP, an inhibitor of glycosphingolipid biosynthesis. Bars indicate median values. A significant increase in the cytotoxic activity of hyperimmune sera over L1210 cells was observed. $P < 0.001$; Wilcoxon test. Loss of NeuGc sialoconjugates or treatment with the inhibitor of glucosylceramide synthetase inhibited cytotoxicity, confirming that the antigen recognized is a N-glycosylated glycosphingolipid. B, histograms of PI-stained cells from 2 representative patients. The numbers represent the percentage of positive cells after subtracting the lysis of pre-immune sera.






Figure 4. OS according to functional capacity of the anti-NeuGcGM3 antibodies elicited after racotumomab-alum vaccination. A, survival of patients with IgM antibodies binding $\geq 30\%$ (red line) versus $< 30\%$ (blue line) to L1210 cells. B, survival of patients with cytotoxic antibodies killing 30% or more (red line) L1210 cells versus patients with less than 30% lysis (blue line).

vaccine as switch maintenance after first-line treatment had a significant survival improvement compared with those who received a placebo.

This study is the first to demonstrate the efficacy of a NeuGc-containing ganglioside anti-idiotype vaccine as switch maintenance therapy for advanced NSCLC. It is noteworthy that most studies designed to find the role of maintenance treatment in patients with advanced NSCLC have shown improvement in PFS (5–14), but only a few demonstrated an impact in OS (6, 7, 9, 14, 15). Our vaccine improved both PFS and OS.

The shorter OS seen in our trial can be attributed to the lack of second-line therapy and the time elapsed between completing first-line chemotherapy and trial enrollment. In addition, apart from other studies on maintenance (6, 9), our trial involved patients with a worse PS (≤ 2) and carrying any NSCLC histology.

Several cancer vaccines have been tested for NSCLC, without great success. Belagenpumatucel-L (Lucanix) is an allogeneic whole-cell vaccine with a TGF- $\beta 2$ antisense.

Results from a phase III, placebo-controlled clinical trial demonstrated that vaccination did not improve OS. Some patients, mainly those randomized within 12 weeks of the completion of chemotherapy, did derive some survival benefit (42). L-BLP25 (StimuVax) is a second cancer vaccine in development for advanced NSCLC, which targets aberrantly glycosylated MUC1. Phase III START study recently failed to reach its primary endpoint of improving OS of patients with stage III NSCLC. Only patients who received concurrent chemoradiotherapy before vaccination had a significant survival increase (43). In both cases, new studies should be done to demonstrate vaccine efficacy. On the other hand, preliminary data from a phase III trial evaluating an EGF-based vaccine suggested that vaccination significantly improved survival of patients with advanced NSCLC. Subjects with higher baseline EGF concentration in serum, derived the greatest benefit (44). Final results from this phase III are still pending.

Better survival with racotumomab-alum was not associated with an increase in adverse events as compared with

placebo. Although patients received repeated injections with racotumomab-alum vaccine, the most common adverse events were classified as mild or moderate, confirming our former results on safety (28–30, 32).

Our group has previously reported the induction of an antibody response against NeuGcGM3 ganglioside in patients with breast, melanoma, SCLC, and NSCLC treated with the racotumomab-alum vaccine (28, 29, 32, 33). In the present study, we confirmed the immunogenicity of this anti-idiotypic vaccine. Racotumomab-alum elicited antibodies specific to the NeuGcGM3 ganglioside and most patients developed high titers of both IgM and IgG isotypes. This is a very relevant result taking into account the general lower immunogenicity of carbohydrates (45, 46). This high antibody response against NeuGcGM3 can be explained by the unique properties of NeuGc-containing gangliosides, which are non-self-antigens in humans, as the result of the inactivation of the gene encoding the enzyme responsible for NeuGc conjugate synthesis (19, 20). The selective expression of NeuGc-gangliosides in tumors appears to originate from exogenous sources (47, 48), and various authors have proposed that tumor hypoxia induces the transcription of a sialin transporter that facilitates the incorporation of NeuGc units into tumor gangliosides (48). Recent reports described the unexpected presence of naturally occurring anti-NeuGcGM3 antibodies, with antitumor potential in healthy donors (39). However, the same team found reduced levels of circulating anti-NeuGcGM3 antibodies in nontreated patients with NSCLC, attributed to the inhibition of the specific antibody-secreting B-cell population, or the formation of immune complexes with the NeuGcGM3 released from tumor cells (39). It has been proposed that idiotypic vaccination is able to boost this natural immune response in patients with cancer.

The current clinical study was designed to evaluate the clinical effect of racotumomab vaccine in NSCLC and to define the surrogate value of the antibodies induced after vaccination. In our set, anti-NeuGGM3 antibodies were able to recognize and kill the NeuGcGM3-expressing cell line L1210, confirming the previously reported results (33). More recently, it was proposed that anti-NeuGcGM3 antibodies elicited by racotumomab induced death by an oncotic–necrosis mechanism, independent of complement activation (41). In the current study, we confirmed that the binding and cytotoxicity on L1210 cells was dependent on the presence of NeuGcGM3 in the cell membrane because the hyperimmune sera did not recognize or kill L1210 cells lacking NeuGcGM3 expression.

Several clinical trials using anti-idiotypic vaccines in patients with cancer have reported a correlation between survival and the immune response against the nominal antigen (33, 49). To our knowledge, this is the first report showing an association between the functional capacity of the anti-NeuGcGM3 antibodies elicited after vaccination and the survival. These preliminary results need to be validated in a larger series of patients. Although the mechanism of action of racotumomab seems to be asso-

ciated with the induction of cytotoxic antibodies capable to kill NeuGcGM3-expressing cells, others mechanisms cannot be ruled out. In fact, there are preclinical evidences of the link between tumor-infiltrating T cells and the antimetastatic and pro-apoptotic effects of the vaccine (31). A specific NeuGcGM3 IFN γ response was also found in patients with metastatic breast cancer treated with racotumomab (34). In the forthcoming trials, immunologic studies should be done to define the capacity of the racotumomab to activate the cellular immunity.

In conclusion, the results of the present study support the safety and efficacy of racotumomab-alum as switch maintenance for patients with advanced NSCLC. This study suggests the value of anti-idiotypic immunization to generate vigorous serologic responses in situations where tumor-associated antigens are poorly immunogenic. The impact of using racotumomab as switch maintenance followed by second-line therapy is currently being explored in a new randomized, multinational phase III study (ClinicalTrials.gov identifier: NCT01460472). In this trial, patients with advanced NSCLC who progress on racotumomab may receive pemetrexed, docetaxel, or erlotinib at physician prescription.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: S. Alfonso, C.E. Viada, I.C. Mendoza, P.P. Guerra, D. Toledo, T. Crombet, R. Pérez, A.M. Vázquez, A.E. Macías

Development of methodology: C.E. Viada, I.C. Mendoza, P.P. Guerra, E. García, Z. Mazorra, T. Crombet, A.E. Macías

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): S. Alfonso, A. Valdés-Zayas, E.R. Santiesteban, Y.I. Flores, F. Areces, R.A. Ortiz, A.V. de la Torre, D. Toledo, A.E. Macías

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A. Valdés-Zayas, Y.I. Flores, M. Hernández, C.E. Viada, T. Crombet, A.M. Vázquez, A.E. Macías

Writing, review, and/or revision of the manuscript: S. Alfonso, A. Valdés-Zayas, M. Hernández, C.E. Viada, A.M. Hernández, Z. Mazorra, T. Crombet, A.M. Vázquez, A.E. Macías

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): A.E. Macías

Study supervision: S. Alfonso, I.C. Mendoza, P.P. Guerra, E. García, M. Cepeda, K. Pérez, E. Chong, Z. Mazorra, T. Crombet, A.E. Macías

Conducted clinical trials: E. Chong

Immunological assessments: Z. González

Senior medical research designed, conducted, and interpreted all the clinical and experimental data of the study: A.E. Macías

Acknowledgments

The authors thank all the participating patients and their families and staffs of all the institutions involved in this study. They also thank Dr. Thomas Rothstein for critically reviewing the article.

Grant Support

Financial support for the study was supplied by The Center of Molecular Immunology and Recombio S.L.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 18, 2013; revised March 17, 2014; accepted April 27, 2014; published OnlineFirst May 1, 2014.

References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011;61:69–90.
- Minna JD. Neoplasmas of the lung. In: Kasper AS, Hauser SL, Longo DL, Jameson JL, editors. *Harrison's principles of internal medicine*. New York: McGraw-Hill; 2005. p. 506–16.
- Schiller JH, Harrington D, Belani CP, Langer C, Sandler A, Krook J, et al. Comparison of four chemotherapy regimens for advanced non-small-cell lung cancer. *N Engl J Med* 2002;346:92–8.
- Gridelli C, de Marinis F, Di Maio M, Ardizzone A, Belani CP, Cappuzzo F, et al. Maintenance treatment of advanced non-small-cell lung cancer: results of an international expert panel meeting of the Italian association of thoracic oncology. *Lung Cancer* 2012;76:269–79.
- Brodowicz T, Krzakowski M, Zwitter M, Tzekova V, Ramlau R, Ghilezan N, et al. Cisplatin and gemcitabine first-line chemotherapy followed by maintenance gemcitabine or best supportive care in advanced non-small cell lung cancer: a phase III trial. *Lung Cancer* 2006;52:155–63.
- Ciuleanu T, Brodowicz T, Zielinski C, Kim JH, Krzakowski M, Laack E, et al. Maintenance pemetrexed plus best supportive care versus placebo plus best supportive care for non-small-cell lung cancer: a randomised, double-blind, phase 3 study. *Lancet* 2009;374:1432–40.
- Fidias PM, Dakhil SR, Lyss AP, Loesch DM, Waterhouse DM, Bromund JL, et al. Phase III study of immediate compared with delayed docetaxel after front-line therapy with gemcitabine plus carboplatin in advanced non-small-cell lung cancer. *J Clin Oncol* 2009;27:591–8.
- Miller VA, O'Connor P, Soh C, Kabbinavar FF. A randomized, double-blind, placebo-controlled, phase IIIb trial (ATLAS) comparing bevacizumab (B) therapy with or without erlotinib (E) after completion of chemotherapy with B for first-line treatment of locally advanced, recurrent, or metastatic non-small cell lung cancer (NSCLC). *J Clin Oncol* 27:18s, 2009 (suppl; abstr LBA8002).
- Cappuzzo F, Ciuleanu T, Stelmakh L, Cicens S, Szczesna A, Juhasz E, et al. Erlotinib as maintenance treatment in advanced non-small-cell lung cancer: a multicentre, randomised, placebo-controlled phase 3 study. *Lancet Oncol* 2010;11:521–9.
- Takeda K, Hida T, Sato T, Ando M, Seto T, Satouchi M, et al. Randomized phase III trial of platinum-doublet chemotherapy followed by gefitinib compared with continued platinum-doublet chemotherapy in Japanese patients with advanced non-small-cell lung cancer: results of a west Japan thoracic oncology group trial (WJTOG0203). *J Clin Oncol* 2010;28:753–60.
- Kabbinavar FF, Miller VA, Johnson BE, O'Connor PG, Soh C. Overall survival (OS) in ATLAS, a phase IIIb trial comparing bevacizumab (B) therapy with or without erlotinib (E) after completion of chemotherapy (chemo) with B for first-line treatment of locally advanced, recurrent, or metastatic non-small cell lung cancer (NSCLC). *J Clin Oncol* 28:15s, 2010 (suppl; abstr 7526).
- Zhang L, Shenglin M, Song X, Han B, Cheng Y, Huang C, et al. Efficacy, tolerability, and biomarker analyses from a phase III, randomized, placebo-controlled, parallel group study of gefitinib as maintenance therapy in patients with locally advanced or metastatic non-small cell lung cancer (NSCLC; INFORM; C-TONG 0804). *J Clin Oncol* 29:15s, 2011 (suppl; abstr LBA7511).
- Perol M, Chouaid C, Perol D, Barlesi F, Gervais R, Westeel V, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin-gemcitabine induction chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 2012;30:3516–24.
- Paz-Ares L, de Marinis F, Dediu M, Thomas M, Pujol JL, Bidoli P, et al. PARAMOUNT: Final overall survival (OS) results of the phase III study of maintenance pemetrexed (pem) plus best supportive care (BSC) versus placebo (plb) plus BSC immediately following induction treatment with pemetrexed plus cisplatin (cis) for advanced nonsquamous (NS) non-small cell lung cancer (NSCLC). *J Clin Oncol* 30, 2012 (suppl; abstr LBA7507).
- Belani CP, Brodowicz T, Ciuleanu T, Kim JH, Krzakowski M, Laack E, et al. Maintenance pemetrexed plus best supportive care (BSC) versus placebo plus BSC: a randomized phase III study in advanced non-small cell lung cancer. *J Clin Oncol* 27:18s, 2009 (suppl; abstr CRA8000).
- Foon KA, Sen G, Hutchins L, Kashala OL, Baral R, Banerjee M, et al. Antibody responses in melanoma patients immunized with an anti-idiotypic antibody mimicking disialoganglioside GD2. *Clin Cancer Res* 1998;4:1117–24.
- Chapman PB. Vaccinating against GD3 ganglioside using BEC2 anti-idiotypic monoclonal antibody. *Curr Opin Investig Drugs* 2003;4:710–5.
- Irie A, Suzuki A. CMP-N-Acetylneuraminic acid hydroxylase is exclusively inactive in humans. *Biochem Biophys Res Commun* 1998;248:330–3.
- Irie A, Koyama S, Kozutsumi Y, Kawasaki T, Suzuki A. The molecular basis for the absence of N-glycolylneuraminic acid in humans. *J Biol Chem* 1998;273:15866–71.
- Olson MV, Varki A. Sequencing the chimpanzee genome: insights into human evolution and disease. *Nat Rev Genet* 2003;4:20–8.
- Marquina G, Waki H, Fernandez LE, Kon K, Carr A, Valiente O, et al. Gangliosides expressed in human breast cancer. *Cancer Res* 1996;56:5165–71.
- van Crujisen H, Ruiz MG, van der Valk P, de Grujil TD, Giaccone G. Tissue micro array analysis of ganglioside N-glycolyl GM3 expression and signal transducer and activator of transcription (STAT)-3 activation in relation to dendritic cell infiltration and microvessel density in non-small cell lung cancer. *BMC Cancer* 2009;9:180.
- Blanco R, Rengifo CE, Cedeno M, Frometa M, Rengifo E, Carr A. Immunoreactivity of the 14F7 mAb (raised against N-glycolyl GM3 ganglioside) as a positive prognostic factor in non-small-cell lung cancer. *Patholog Res Int* 2012;2012:235418.
- Hayashi N, Chiba H, Kuronuma K, Go S, Hasegawa Y, Takahashi M, et al. Detection of N-glycosylated gangliosides in non-small-cell lung cancer using GMR8 monoclonal antibody. *Cancer Sci* 2013;104:43–7.
- de Leon J, Fernandez A, Mesa C, Clavel M, Fernandez LE. Role of tumour-associated N-glycosylated variant of GM3 ganglioside in cancer progression: effect over CD4 expression on T cells. *Cancer Immunol Immunother* 2006;55:443–50.
- Vazquez AM, Perez A, Hernandez AM, Macias A, Alfonso M, Bombino G, et al. Syngeneic anti-idiotypic monoclonal antibodies to an anti-NeuGc-containing ganglioside monoclonal antibody. *Hybridoma* 1998;17:527–34.
- Vazquez AM, Alfonso M, Lanne B, Karlsson KA, Carr A, Barroso O, et al. Generation of a murine monoclonal antibody specific for N-glycolylneuraminic acid-containing gangliosides that also recognizes sulfated glycolipids. *Hybridoma* 1995;14:551–6.
- Alfonso M, Diaz A, Hernandez AM, Perez A, Rodriguez E, Bitton R, et al. An anti-idiotypic vaccine elicits a specific response to N-glycolyl sialic acid residues of glycoconjugates in melanoma patients. *J Immunol* 2002;168:2523–9.
- Neninger E, Diaz RM, de la Torre A, Rives R, Diaz A, Saurez G, et al. Active immunotherapy with 1E10 anti-idiotypic vaccine in patients with small cell lung cancer: report of a phase I trial. *Cancer Biol Ther* 2007;6:145–50.
- Alfonso S, Diaz RM, de la Torre A, Santiesteban E, Aguirre F, Perez K, et al. 1E10 anti-idiotypic vaccine in non-small cell lung cancer: experience in stage IIIb/IV patients. *Cancer Biol Ther* 2007;6:1847–52.
- Diaz Y, Gonzalez A, Lopez A, Perez R, Vazquez AM, Montero E. Anti-ganglioside anti-idiotypic monoclonal antibody-based cancer vaccine induces apoptosis and antiangiogenic effect in a metastatic lung carcinoma. *Cancer Immunol Immunother* 2009;58:1117–28.
- Diaz A, Alfonso M, Alonso R, Saurez G, Troche M, Catala M, et al. Immune responses in breast cancer patients immunized with an anti-idiotypic antibody mimicking NeuGc-containing gangliosides. *Clin Immunol* 2003;107:80–9.
- Hernandez AM, Toledo D, Martinez D, Grinan T, Brito V, Macias A, et al. Characterization of the antibody response against NeuGcGM3 ganglioside elicited in non-small cell lung cancer patients immunized with an anti-idiotypic antibody. *J Immunol* 2008;181:6625–34.
- Guthmann MD, Castro MA, Cinat G, Venier C, Koliren L, Bitton RJ, et al. Cellular and humoral immune response to N-Glycolyl-GM3 elicited by prolonged immunotherapy with an anti-idiotypic vaccine in high-risk and metastatic breast cancer patients. *J Immunother* 2006;29:215–23.

35. Oken MM, Creech RH, Tormey DC, Horton J, Davis TE, McFadden ET, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 1982;5:649–55.
36. Therasse P, Arbuuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000;92:205–16.
37. Stults CL, Sweeley CC, Macher BA. Glycosphingolipids: structure, biological source, and properties. *Methods Enzymol* 1989;179:167–214.
38. Roque-Navarro L, Chakrabandhu K, de Leon J, Rodriguez S, Toledo C, Carr A, et al. Anti-ganglioside antibody-induced tumor cell death by loss of membrane integrity. *Mol Cancer Ther* 2008;7:2033–41.
39. Rodriguez-Zhurbenko N, Martinez D, Blanco R, Rondon T, Grinan T, Hernandez AM. Human antibodies reactive to NeuGcGM3 ganglioside have cytotoxic antitumor properties. *Eur J Immunol* 2013;43:826–37.
40. Fernandez-Marrero Y, Hernandez T, Roque-Navarro L, Talavera A, Moreno E, Grinan T, et al. Switching on cytotoxicity by a single mutation at the heavy chain variable region of an anti-ganglioside antibody. *Mol Immunol* 2011;48:1059–67.
41. Hernandez AM, Rodriguez N, Gonzalez JE, Reyes E, Rondon T, Grinan T, et al. Anti-NeuGcGM3 antibodies, actively elicited by idiotypic vaccination in nonsmall cell lung cancer patients, induce tumor cell death by an oncosis-like mechanism. *J Immunol* 2011;186:3735–44.
42. Giaccone G, Bazhenova L, Nemunaitis N, Juhasz E, Ramlau R, van den Heuvel MM, et al. A phase III study of belagenpumatucel-L therapeutic tumor cell vaccine for non-small cell lung cancer. *European Cancer Congress, 2013 (suppl; abstr LBA 2)*.
43. Butts C, Socinski MA, Mitchell P, Thatcher N, Havel L, Krzakowski M, et al. START: A phase III study of L-BLP25 cancer immunotherapy for unresectable stage III non-small cell lung cancer. *J Clin Oncol* 31, 2013 (suppl; abstr 7500).
44. Crombet T, Neninger E, Acosta S, Amador R, Mendoza S, Santiesteban E, et al. EGF-based cancer vaccine for advanced NSCLC: Results from a phase III trial. *J Clin Oncol* 30, 2012 (suppl; abstr 2527).
45. Livingston PO. Augmenting the immunogenicity of carbohydrate tumor antigens. *Semin Cancer Biol* 1995;6:357–66.
46. Yao TJ, Meyers M, Livingston PO, Houghton AN, Chapman PB. Immunization of melanoma patients with BEC2-keyhole limpet hemocyanin plus BCG intradermally followed by intravenous booster immunizations with BEC2 to induce anti-GD3 ganglioside antibodies. *Clin Cancer Res* 1999;5:77–81.
47. Tangvoranuntakul P, Gagneux P, Diaz S, Bardor M, Varki N, Varki A, et al. Human uptake and incorporation of an immunogenic nonhuman dietary sialic acid. *Proc Natl Acad Sci U S A* 2003;100:12045–50.
48. Yin J, Hashimoto A, Izawa M, Miyazaki K, Chen GY, Takematsu H, et al. Hypoxic culture induces expression of sialin, a sialic acid transporter, and cancer-associated gangliosides containing non-human sialic acid on human cancer cells. *Cancer Res* 2006;66:2937–45.
49. Birebent B, Mitchell E, Akis N, Li W, Somasundaram R, Purev E, et al. Monoclonal anti-idiotypic antibody mimicking the gastrointestinal carcinoma-associated epitope CO17-1A elicits antigen-specific humoral and cellular immune responses in colorectal cancer patients. *Vaccine* 2003;21:1601–12.

Clinical Cancer Research

A Randomized, Multicenter, Placebo-Controlled Clinical Trial of Racotumomab-Alum Vaccine as Switch Maintenance Therapy in Advanced Non–Small Cell Lung Cancer Patients

Sailyn Alfonso, Anet Valdés-Zayas, Eduardo R. Santiesteban, et al.

Clin Cancer Res 2014;20:3660-3671. Published OnlineFirst May 1, 2014.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-13-1674](https://doi.org/10.1158/1078-0432.CCR-13-1674)

Supplementary Material Access the most recent supplemental material at:
<http://clincancerres.aacrjournals.org/content/suppl/2014/05/01/1078-0432.CCR-13-1674.DC1>

Cited articles This article cites 40 articles, 13 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/20/14/3660.full#ref-list-1>

Citing articles This article has been cited by 8 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/20/14/3660.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/20/14/3660>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.