

# Identification of a Three-Biomarker Panel in Urine for Early Detection of Pancreatic Adenocarcinoma

Tomasz P. Radon<sup>1</sup>, Nathalie J. Massat<sup>2</sup>, Richard Jones<sup>3</sup>, Wasfi Alrawashdeh<sup>1</sup>, Laurent Dumartin<sup>1</sup>, Darren Ennis<sup>1</sup>, Stephen W. Duffy<sup>2</sup>, Hemant M. Kocher<sup>4</sup>, Stephen P. Pereira<sup>5</sup>, Luisa Guarner (posthumous)<sup>6</sup>, Cristiane Murta-Nascimento<sup>7</sup>, Francisco X. Real<sup>8</sup>, Núria Malats<sup>8</sup>, John Neoptolemos<sup>9</sup>, Eithne Costello<sup>9</sup>, William Greenhalf<sup>9</sup>, Nick R. Lemoine<sup>1</sup>, and Tatjana Crnogorac-Jurcevic<sup>1</sup>

## Abstract

**Purpose:** Noninvasive biomarkers for early detection of pancreatic ductal adenocarcinoma (PDAC) are currently not available. Here, we aimed to identify a set of urine proteins able to distinguish patients with early-stage PDAC from healthy individuals.

**Experimental design:** Proteomes of 18 urine samples from healthy controls, chronic pancreatitis, and patients with PDAC (six/group) were assayed using GeLC/MS/MS analysis. The selected biomarkers were subsequently validated with ELISA assays using multiple logistic regression applied to a training dataset in a multicenter cohort comprising 488 urine samples.

**Results:** LYVE-1, REG1A, and TFF1 were selected as candidate biomarkers. When comparing PDAC ( $n = 192$ ) with healthy ( $n = 87$ ) urine specimens, the resulting areas under the receiver-operating characteristic curves (AUC) of the panel were 0.89 [95% confidence interval (CI), 0.84–0.94] in the training (70% of the

data) and 0.92 (95% CI, 0.86–0.98) in the validation (30% of the data) datasets. When comparing PDAC stage I–II ( $n = 71$ ) with healthy urine specimens, the panel achieved AUCs of 0.90 (95% CI, 0.84–0.96) and 0.93 (95% CI, 0.84–1.00) in the training and validation datasets, respectively. In PDAC stage I–II and healthy samples with matching plasma CA19.9, the panel achieved a higher AUC of 0.97 (95% CI, 0.94–0.99) than CA19.9 (AUC = 0.88; 95% CI, 0.81–0.95,  $P = 0.005$ ). Adding plasma CA19.9 to the panel increased the AUC from 0.97 (95% CI, 0.94–0.99) to 0.99 (95% CI, 0.97–1.00,  $P = 0.04$ ), but did not improve the comparison of stage I–IIA PDAC ( $n = 17$ ) with healthy urine.

**Conclusions:** We have established a novel, three-protein biomarker panel that is able to detect patients with early-stage pancreatic cancer in urine specimens. *Clin Cancer Res*; 21(15):3512–21. ©2015 AACR.

## Introduction

Pancreatic ductal adenocarcinoma (PDAC) is one of the rare cancers for which no significant improvements in diagnosis and therapy have been made in the last 30 years. Despite considerable progress in our understanding of the disease at the molecular level, novel findings have not yet translated into clinical benefit, and the majority of patients are still faced with a grim median

survival of 5 to 6 months. With over 38,000 PDAC-related deaths in the United States and over 40,000 in Europe in 2013, this malignancy is currently the fourth leading cause of cancer-related death, but predicted to become the second by 2030 (1–3).

These worrying figures would change significantly with improved tool(s) for early detection, as 5-year survival approaching 70% has been reported after incidental diagnosis of stage I PDAC tumors, when they were still confined to the pancreas with a size <2 cm (4–6). Detection at an early stage is also crucial given the poor efficacy of current therapies for metastatic disease, when potentially curative surgery is no longer feasible.

Timely detection of PDAC is, however, hampered by several factors: lack of specific clinical symptoms in the early stage of the disease, insufficient sensitivity of current imaging modalities and, despite intensive efforts, lack of accurate body fluid–based biomarkers of early-stage disease (for review, see 7). Early-stage PDAC is also difficult to differentiate from chronic pancreatitis (CP), a benign inflammatory disease of the pancreas and one of the risk factors for PDAC (8). Serum CA19.9, the only PDAC biomarker in widespread clinical use, suffers from false-negative results in patients with Lewis-negative genotype, low sensitivity (79%–81%) in symptomatic patients, and its levels may be elevated in various other benign and malignant pancreatic and hepatobiliary diseases, as well as in unrelated cystic and inflammatory diseases (for review, see 9).

After a successful proof-of-concept study where we showed that the protein signatures of CP and PDAC can be identified in urine

<sup>1</sup>Centre for Molecular Oncology, Barts Cancer Institute, Queen Mary University of London, London, United Kingdom. <sup>2</sup>Centre for Cancer Prevention, Wolfson Institute of Preventive Medicine, Queen Mary University of London, London, United Kingdom. <sup>3</sup>MS Bioworks, LLC, Ann Arbor, Michigan. <sup>4</sup>Centre for Tumour Biology, Barts Cancer Institute, Queen Mary University of London, London, United Kingdom. <sup>5</sup>Institute for Liver and Digestive Health, University College London, London, United Kingdom. <sup>6</sup>Hospital General Universitari Vall Hebron, Barcelona, Spain. <sup>7</sup>Hospital del Mar — Parc de Salut Mar, Barcelona, Spain. <sup>8</sup>Centro Nacional de Investigaciones Oncológicas (CNIO), Madrid, Spain. <sup>9</sup>The NIHR Liverpool Pancreas Biomedical Research Unit, Liverpool, United Kingdom.

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

**Corresponding Author:** Tatjana Crnogorac-Jurcevic, Barts Cancer Institute, Queen Mary University of London, John Vane Science Centre, Charterhouse Square, London, EC1M 6BQ, UK. Phone: 44-0-20-7882-3554; Fax: 44-0-20-7882-3884; E-mail: t.c.jurcevic@qmul.ac.uk

**doi:** 10.1158/1078-0432.CCR-14-2467

©2015 American Association for Cancer Research.

### Translational Relevance

Currently, 80% of patients diagnosed with pancreatic adenocarcinoma present with locally invasive and/or metastatic disease, resulting in a poor 5-year survival of <5%. The development of a diagnostic tool for early detection of patients with this malignancy may significantly impact their prognosis. We established a panel of three urine biomarkers that can distinguish patients with early-stage disease from healthy people, which could enable completely noninvasive and inexpensive screening of patients at high risk of developing pancreatic adenocarcinoma.

(10), we here describe the development of a three-biomarker panel that can detect early-stage PDAC completely noninvasively, through analysis of urine samples.

## Materials and Methods

### Clinical specimens

Healthy, CP, and PDAC ( $n = 6$ /each group) urine specimens were obtained from the Royal London Hospital (RLH) and used for the discovery phase. For validation purposes, a total of 371 urine specimens [87 healthy, 92 CP, and 192 PDAC urine samples

from RLH and University College London (jointly referred to as "LON"), the Department of Surgery, Liverpool University ("LIV"), and the CNIO Madrid, Spain ("SPA")] were assayed. Demographics and clinical characteristics of patients and healthy participants included in the study are shown in Table 1. Additional 117 urine samples from patients with other benign and malignant hepatobiliary pathologies [33 from patients with intraductal papillary mucinous tumors (IPMN) without associated adenocarcinoma, 18 from pancreatic neuroendocrine tumors (NET), 16 from duodenal cancers, 26 from ampullary cancers, and 24 from patients with cholangiocarcinomas (CHL)] were obtained from LIV (demographic details are provided in Supplementary Fig. S5A). Matching plasma samples for measuring CA19.9 were available from RLH and LIV. Healthy individuals had no known pancreatic conditions or malignancies, and all samples were derived from individuals with no history of renal diseases. Dipstick test analysis (Bayer multistix SG 08935414) was also performed to exclude potential bilirubinemia, proteinuria, bacterial contamination, and hematuria. The specimens in all participating centers were collected using the same standard operating procedures: clean-catch, midstream urine was collected, frozen within 2 hours of collection, and stored at  $-80^{\circ}\text{C}$  until utilized; samples were collected before any surgery or chemotherapeutic treatment. All samples were collected with full ethical approval from the involved centers and with informed consent from all individuals who donated urine/blood samples.

**Table 1.** Demographics and clinical characteristics of the healthy and patient cohorts

	Normal			CP			PDAC			Stage/n plasma <sup>a</sup>
	Cases (n)	Gender	Age range (median)	Cases (n)	Gender	Age range (median)	Cases (n)	Gender	Age range (median)	
LON	87	M = 46 F = 41	28–87 (55)	45	M = 32 F = 13	29–82 (53)	60	M = 38 F = 22	29–82 (64)	I = 4/4 IIA = 1/1 IIB = 13/13 III = 33/30 IV = 6/5 U = 3/3
Plasma (CA19.9)	28	M = 16 F = 12	28–67 (46)	19	M = 14 F = 5	29–74 (54)	56	M = 34 F = 22	29–82 (64)	
LIV	0	N/A	N/A	41	M = 25 F = 16	29–82 (51)	91	M = 53 F = 38	39–83 (68)	I = 3/3 IIA = 8/8 IIB = 42/42 III = 38/38 IV = 0/0 U = 0/0
Plasma (CA19.9)	0	N/A	N/A	31	M = 17 F = 14	37–73 (51)	91	M = 53 F = 38	39–83 (68)	
SPA	0	N/A	N/A	6	M = 4 F = 2	54–68 (57)	41	M = 23 F = 18	43–94 (72)	I = 0/NA II = 0/NA III = 0/NA IV = 0/NA U = 41/NA
Plasma (CA19.9)	0	N/A	N/A	0	N/A	N/A	0	N/A	N/A	
Total	87	M = 46 F = 41	28–87 (55)	92	M = 61 F = 31	29–82 (54)	192	M = 114 F = 78	29–94 (68)	I = 7/7 IIA = 9/9 IIB = 55/55 III = 71/68 IV = 6/5 U = 44/3
Plasma (CA19.9)	28	M = 16 F = 12	28–67 (46)	50	M = 31 F = 19	29–74 (53)	147	M = 87 F = 60	29–83 (67)	

NOTE: Total urine = 371; total plasma = 147.

<sup>a</sup> Number of cases with plasma CA19.9.

### GeLC-MS/MS (SDS-PAGE-liquid chromatography-tandem mass spectrometry) analysis of urine proteomes

Six urine samples (three males and three females) for each group (healthy, CP, and PDAC; in total 18 samples) were utilized: healthy males/females age 45, 50, 60/44, 45, 54 years; CP males/females age 46, 48, 51/47, 69, 74 years; PDAC males/females age 44, 74, 84/71, 73, 77 years; male PDAC stage all IIB/female two IB, one IIA. All urine samples were desalted and concentrated as described previously (10). Twenty microgram of each preprocessed pool of three samples per group was separated in duplicate on 4%–12% mini-gels (Invitrogen); female and male urine specimens were analyzed separately. The gels were stained with Colloidal Coomassie, and each sample lane cut using a grid into 40 equally sized slices. Gel slices were digested robotically with trypsin and resultant peptides analyzed by nano LC/MS/MS using a nanoAcquity (Waters) interfaced to a LTQ Orbitrap XL tandem mass spectrometer (ThermoFisher). Product ion data were searched against the human IPI protein database using Mascot, and subsequently parsed into the Scaffold software (Proteome Software) for collation into nonredundant protein lists. Reversed database searching was used to assess false discovery rates, the target protein FDR being <0.5% per sample. A semiquantitative assessment of relative protein abundance between PDAC, CP, and healthy samples was obtained using the spectral counting approach (11).

### Urine biomarkers and CA19.9 measurements

Total protein concentration in urine specimens was determined by Bradford assay (Coomassie Protein Assay Reagent; Pierce). The quantitative ELISA determination of human LYVE-1 (Cat# SEB049Hu; Uscn Life Science Inc.) and human TFF1 (Cat# ELH-LYVE1-001; RayBiotech Inc.) was performed according to the manufacturer's instructions; human REG1A levels were initially assessed in our laboratory, and afterwards by BioVendor Analytical Testing Service (BioVendor - Laboratorní medicína a.s). Calibration curves were prepared using purified standards for each protein assessed. Curve fitting was accomplished by a four-parameter logistic regression following the manufacturer's instructions. The limits of detection and the coefficient of variation (CV) for each of the ELISA assays were as follows: LYVE-1, 8.19 pg/mL, intraassay CV, 9%, interassay CV, 12%; TFF1, 0.037 ng/mL, intraassay CV, 9%, interassay CV, 12%; REG1A, 0.094 ng/mL, intraassay CV, 9%, interassay CV, 20%; REG1B, 3.13 pg/mL, intraassay CV, 3.9%, interassay CV, 2.7%. Urine creatinine was measured by the Jaffé method using the Roche Cobas 8000 system (Roche Diagnostics) and plasma CA19.9 using a Roche Modular E170 instrument according to the routine protocols at the Clinical Biochemistry Laboratory, RLH (London, UK).

### Tissue microarrays and immunohistochemistry

The details of the tissue microarray and scoring procedure used in evaluating the expression of the biomarkers were described previously (12). Immunohistochemistry (IHC) was performed with anti-REG1A (Abcam; Rabbit polyclonal, ab47099, 1:100 dilution), anti-TFF1 (Abcam; Rabbit polyclonal, ab50806, 1:100 dilution), and anti-LYVE1 (Acris; Rabbit polyclonal, DP3500PS, 1:100 dilution) antibodies using the Ventana Discovery system, according to standard protocols (sCC1, 1-hour incubation).

### Statistical analysis

To identify potential urine biomarkers from the MS data, the statistical analysis was performed on the normalized data (based on the sum of spectral counts/sample) using Arraytrack software (<http://edkb.fda.gov/webstart/arraytrack>) by the  $t$  test. The data were further filtered according to both  $P$  values and fold change between any two sample groups, separately in males and females.

The concentrations of the selected proteins (LYVE1, REG1A, and TFF1) obtained by ELISA assays were compared using non-parametric Kruskal–Wallis tests. To adjust for inflated type I error due to multiple testing between patients with PDAC and control groups (healthy and CP), i.e., 27 comparisons, the significance threshold for  $P$  values was adjusted using the conservative Bonferroni correction, i.e., a threshold of  $\alpha(0.05)/27 = 0.0018$  was used to define a significant result at the 5% level after adjustment for multiplicity.

Correlation between the three biomarkers was assessed using the Pearson correlation coefficient.

Each individual biomarker and the panel were investigated for their ability to discriminate between patients with PDAC (all stages, or early stages I–II) and control samples (healthy and CP) using ROC analysis and a hold-out approach. For each comparison, 70% of the subjects in the patient and control datasets were randomly selected for inclusion in the training dataset. Logistic regression was then applied. All protein concentration data were natural log-transformed and mean-centered before regression analysis. In individual biomarker analyses, creatinine-normalized data were used to correct for the urine dilution factor; for the panel analysis, the model included the three biomarkers (before creatinine normalization) and was adjusted for creatinine and age (as the median age of patients with PDAC was higher than that of healthy and CP individuals, Table 1), i.e., 5-parameter model. Separate models were applied to the training datasets for the comparison of PDAC all stages versus healthy, PDAC stages I–II versus healthy, PDAC all stages versus CP and PDAC stages I–II versus CP. ROC curves were generated for each of the above regression models; the AUC, and the sensitivity (SN) and specificity (SP) at the "optimal" cut-point for discrimination between groups were obtained. The optimal cut-point corresponded to the point closest to the top-left part of the plot in the ROC plane (coordinates 0,1) with optimal SN and SP according to the following criterion:

$$\min((1 - \text{sensitivities})^2 + (1 - \text{specificities})^2),$$

as calculated by the "ci.threshold" procedure of the R "pROC" package (13). This approach has been shown to have good performance in the estimation of the optimal cut-point of a biomarker (14).

The rest of the subjects (30%) formed independent datasets which were used for model validation. For the primary analysis (all PDAC versus healthy), 49 PDAC and 28 healthy samples give more than 90% power to detect a standardized difference of 1.0 (i.e., a difference between PDAC and healthy samples of at least one standard deviation) using a one-sided test.

Validation was performed by classifying each sample in the validation dataset according to the logistic regression model developed based on the training dataset, and comparing this classification with the actual diagnosis, hence deriving a new ROC curve. The optimal cut-points computed for the training sets were used to derive the SN and SP of the validation dataset. Confidence intervals (CI, 95%) for AUCs were derived based on

the DeLong' asymptotically exact method to evaluate the uncertainty of an AUC (15); SN and SP 95% CI were derived using nonparametric stratified resampling with the percentile method (2,000 bootstrap replicates) as described by (16). AUCs were compared using the DeLong's 1-sided test for correlated/paired AUCs (15).

For exploratory analyses, ROC curves were derived for the comparison of PDAC stage I–IIA versus healthy or CP based on logistic regression modeling using all available samples.

ROC curve analyses were performed in R version 2.13.0 (The R Foundation for Statistical Computing, <http://www.r-project.org/foundation>) using procedures from the Epi (14), pROC (13), and ROCr (17) packages.

## Results

### Urine proteomes

We undertook an in-depth proteomics analysis by GeLC/MS/MS of 18 urine specimens derived from PDAC, CP, and healthy individuals (6 per group, three males, three females; Supplementary Fig. S1A). This analysis resulted in the identification of around 1,500 (1,198 in male and 1,061 in female urine) nonredundant proteins (Supplementary Table S1). These proteins originated from all cellular compartments and were mapped using IPA (Ingenuity pathway analysis, <http://www.ingenuity.com/>) to a number of cellular functions and diseases (Supplementary Fig. S1B and S1C, respectively), confirming that urine provides a rich source of diverse proteins with respect to their origin and functional roles.

Our MS analysis was performed separately on urine samples from male and female subjects. We noticed considerable gender-specific differences: of 997 proteins identified in healthy urine samples, 398 (40%) were unique to those found in samples from males, 118 (12%) were unique to females, and 481 (48%) were common to both.

Three proteins commonly deregulated in both males and females (LYVE1, REG1A, and TFF1) were selected for further evaluation based on the statistics (*t* test, *P* values <0.05; fold change >1.5), interrogation of Pancreatic Expression Database (<http://www.pancreasexpression.org/>; ref. 18), and additional literature search for previous knowledge on the potential candidates, and also on the availability of commercial ELISA assays. While REG1B in our proteomics data appeared to be slightly better candidate, only REG1A ELISA assay was available commercially at the time. However, when REG1B ELISA became available, we tested a subset of urine samples and similar results were obtained (see later). The presence of our three selected biomarkers as full-size proteins: 35 kDa for LYVE1, 19 kDa for REG1A, and 9 kDa for TFF1 in urine specimens was confirmed by Western blot (data not shown).

### Biomarker panel in detecting PDAC

The selected biomarkers were subsequently assessed using ELISA assays on 371 urine samples collected from three centers: London and Liverpool, UK, and Madrid, Spain. Demographics and clinical characteristics of patients and healthy participants included in the study are shown in Table 1.

### PDAC stage I–IV versus healthy

The ELISA analysis showed significantly higher urine concentrations for each of the candidate biomarkers in the urine of patients with PDAC (*n* = 192) when compared with healthy

samples (*n* = 87, all with *P* < 0.0001, Fig. 1). Of note, REG1B and REG1A ELISA assays produced similar results (Fig. 1).


In PDAC, LYVE1, REG1A, and TFF1 were positively correlated with each other, whereas in healthy samples, only LYVE1 and REG1A were correlated (Supplementary Fig. S2).

The diagnostic performance of LYVE1, REG1A, and TFF1 was established using ROC curve analysis (Fig. 2). We first assessed their individual performance in discriminating between PDAC stages I–IV and healthy urine samples in a training dataset (70% of the samples, *n* = 143 and *n* = 59, respectively). Individual (creatinine-normalized) urine biomarkers were able to discriminate between the two groups with AUC values of 0.851 (95% CI, 0.801–0.902) for LYVE1, 0.823 (95% CI, 0.766–0.879) for REG1A, and 0.686 (95% CI, 0.606–0.765) for TFF1, with respective SN of 76.9% (95% CI, 69.3–83.2), 62.2% (95% CI, 53.8–69.9), and 72.7% (95% CI, 65.0–79.7), and respective SP of 88.1% (95% CI, 79.6–96.6), 94.9% (95% CI, 88.1–100.0), and 59.3% (95% CI, 47.5–71.2; Fig. 2A and C). The three biomarkers were then combined into a panel adjusted for creatinine and age (Fig. 2B). The results of the logistic regression model underlying the ROC analysis in the training and validation (30% of the samples, PDAC *n* = 49, healthy *n* = 28) datasets are shown in Fig. 2B and C. The panel achieved SN >75% and SP >85% for AUCs of 0.891 (95% CI, 0.847–0.935) and 0.921 (95% CI, 0.863–0.978) in the training and validation datasets, respectively, thus showing better performance than any of the individual biomarkers.

### PDAC early stages versus healthy

Next, we assessed the performance of our biomarkers in discriminating early-stage cancers from healthy individuals. Tumor staging information was available for 148 (77%) of the PDAC patients. The concentrations of each of the biomarkers were significantly increased in later stages (stage III–IV, *n* = 77, all *P* < 0.0001), in stages I–II (*n* = 71, all *P* < 0.0001), and in stages I–IIA (locally invasive disease without lymph node metastases, *n* = 16, *P* < 0.0001, except for REG1A *P* = 0.007) compared with healthy people (*n* = 87; Fig. 3). The performance of the individual markers and the panel in discriminating between PDAC stage I–II from healthy urine samples was first assessed in a new training dataset (70% of the samples; PDAC stage I–II *n* = 56 and healthy *n* = 61, respectively). A new 5-parameter model was built using this training dataset and validated using the rest of the data (30% of the samples; PDAC stage I–II *n* = 15, healthy *n* = 26; Fig. 4A and B). The panel achieved AUCs of 0.900 (95% CI, 0.843–0.957) and 0.926 (95% CI, 0.843–1.000) in the training and validation datasets, respectively (Fig. 4C). Therefore, our urine biomarker panel can differentiate early PDAC from healthy samples with high accuracy.

As an exploratory analysis, we then selected the urine samples from individuals for which matched plasma samples were available so CA19.9 values could be obtained. The ROC curves were derived for plasma CA19.9 (as a categorical variable with a cutoff at clinically established threshold of 37 U/mL), the panel, and a combination of the panel and CA19.9. For the comparison of PDAC stage I–II (*n* = 71) versus healthy (*n* = 28) samples, we obtained AUCs of 0.880 (95% CI, 0.947–0.999) for CA19.9, and 0.973 (95% CI, 0.947–0.999) for the panel, which was significantly greater than plasma CA19.9 alone (*P* = 0.005). The addition of plasma CA19.9 to the panel significantly increased the AUC to 0.991 (95% CI, 0.979–1.000, *P* = 0.04; Fig. 5A and C). When PDAC stages I–IIA (*n* = 16) were compared with healthy



**Figure 1.** Urine concentration of the candidate protein biomarkers. A, scatter dot plots of LYVE1, REG1A, REG1B, and TFF1 protein concentration (creatinine-normalized) analyzed by ELISA in healthy, CP, and PDAC patients' urine. Top bars: Kruskal-Wallis test; \*\*\*\*,  $P < 0.0001$ ; \*\*\*,  $P < 0.001$ . B, statistical summary. Median and interquartile range (IQR) of raw/creatinine-normalized data for the biomarkers, median, and IQR of urine creatinine (mmol/L), as well as plasma CA19.9 (U/mL) by sample groups are shown.

samples, AUCs were 0.839 (95% CI, 0.719–0.959) for CA19.9, and 0.971 (95% CI, 0.929–1.000) for the panel ( $P = 0.006$ ). The addition of plasma CA19.9 to the panel did not result in any improvement (AUC = 0.969; 95% CI, 0.924–1.000,  $P = 0.7$ ; Fig. 5B and C).


**Biomarker panel in differentiating PDAC from CP**

**PDAC stage I–IV versus CP.** Urine concentration for all three biomarkers was higher in PDAC ( $n = 192$ ) compared with CP samples ( $n = 92$ ,  $P < 0.0001$  for LYVE1 and REG1A and  $P = 0.0002$  for TFF1; Fig. 1), and as for PDAC, the biomarker concentrations were positively correlated with each other in the CP data (Supplementary Fig. S2). In the training dataset (PDAC  $n = 143$ , CP  $n = 62$ ), LYVE1 and REG1A were able to discriminate between the two groups with SN of 77% to 78% and SP of 66% to 69% [respective AUC values of 0.775 (95% CI, 0.704–0.846) and 0.722 (95% CI, 0.643–0.801; Supplementary Fig. S3)], while the SP of TFF1 only reached 50% for a similar SN. Combining the three biomarkers into a panel only improved marginally the performance of LYVE1 and REG1A alone as assessed in the training (AUC = 0.815; 95% CI, 0.752–0.878) and validation (PDAC  $n = 49$ , CP  $n = 30$ ; AUC = 0.839; 95% CI, 0.751–0.928) datasets.

**PDAC early stages versus CP.** Biomarker urine concentrations were significantly increased in stage I–II PDAC ( $n = 71$ ) compared with CP ( $n = 87$ , with  $P < 0.0001$  for LYVE1 and REG1A and  $P = 0.0001$  for TFF1, data not shown). The panel achieved high SN (>85%) in both the training (PDAC stage I–II  $n = 56$ , CP  $n = 66$ ) and validation (PDAC stage I–II  $n = 15$ , CP  $n = 26$ ) datasets, but relatively low SP (66.7% and 50%), similar to the SP observed for individual biomarkers, with respective AUCs of 0.831 (95% CI, 0.762–0.901) and 0.846 (95% CI, 0.730–0.963; Supplementary Fig. S3D–S3F).

As before, we explored the panel in combination with plasma CA19.9. For the comparison of PDAC stage I–II ( $n = 71$ ) versus CP ( $n = 50$ ) samples, the ROC curves showed AUCs of 0.775 (95% CI, 0.699–0.852) for CA19.9, 0.830 (95% CI, 0.759–0.902) for the panel ( $P = 0.1$ ), and 0.885 (95% CI, 0.825–0.945) for the panel in combination with CA19.9 ( $P = 0.01$  for superiority over the panel alone; Supplementary Fig. S4A and S4C). In the comparison of PDAC stage I–IIA ( $n = 16$ ) versus CP, the ROC curves showed AUCs of 0.735 (95% CI, 0.609–0.861) for CA19.9, 0.871 (95% CI, 0.770–0.972) for the panel ( $P = 0.004$  for superiority over plasma CA19.9), and 0.866 (95% CI, 0.749–0.984) for the combination ( $P = 0.6$ ; Supplementary Fig. S4B and S4C).

**Figure 2.** Diagnostic performance of urine biomarkers in discriminating patients with PDAC from healthy controls. A, ROC curves of PDAC ( $n = 143$ ) versus healthy ( $n = 59$ ) subjects for individual creatinine-normalized urine biomarkers in the training set (70% of the data). B, ROC curves of PDAC versus healthy for the panel in the training set and in the independent validation set (30% of the data: PDAC  $n = 49$ , healthy  $n = 28$ ); C, summary table. SN and SP in the validation set are derived for optimal cutpoint determined in the training dataset. cnorm, creatinine-normalized; creat, creatinine.



Therefore, the panel performed better in differentiating stage I–IIA from CP than CA19.9.


**Biomarker expression in urine of other hepatobiliary pathologies**

We explored the expression of our biomarkers in urine specimens collected from patients with several other benign or malignant hepatobiliary pathologies and compared it to the expression in patients with early-stage PDAC (Supplementary Fig. S5). Levels of LYVE1 tended to be higher in PDAC stage I–II samples compared with IPMNs, ampullary cancer, and pancreatic NETs specimens, whereas REG1A levels were only increased in comparison with IPMNs. Plasma CA19.9 levels were higher in PDACs stage I–II compared with pancreatic NETs and duodenal cancer samples. This might suggest a potential utility for LYVE1 and REG1A in


distinguishing other benign or malignant hepatobiliary pathologies from early-stage PDACs.

**Tissue origin of the three biomarkers**

Having demonstrated a good performance of the panel in differentiating early cancer patients from healthy individuals, we next sought to establish the expression of our biomarkers in pancreatic tissue. We performed IHC using in-house constructed PDAC tissue microarrays. A strong expression of REG1A was seen in histologically normal adjacent acinar cells, but the staining was also seen in 44 of 60 tumors (73%; Supplementary Fig. S6A). TFF1 was absent in normal pancreas, but was expressed in 43 of 60 (72%) PDACs (Supplementary Fig. S6B). Although no LYVE1 expression was seen in any of the cancer cells, it was seen in lymphatic vessels in eight PDAC tissues (Supplementary



**Figure 3.** Urine concentration of the three biomarkers in different stages of PDAC. Scatter dot plots of urine LYVE1, REG1A, TFF1 protein concentration (creatinine-normalized) in urine samples of healthy ( $n = 87$ ) and patients with PDAC at different stages of disease development (I–IIA  $n = 16$ , I–II  $n = 71$ , III–IV  $n = 77$ ). Bars indicate median and IQR values. Top bars: Kruskal–Wallis test; \*\*\*\*,  $P < 0.0001$ .



**Figure 4.** Diagnostic performance of urine biomarkers in discriminating early patients with PDAC from healthy individuals. A, ROC curves of stage I-II PDAC ( $n = 56$ ) versus healthy ( $n = 61$ ) subjects for individual urine biomarkers in the training set (70% of the data). B, ROC curves of stage I-II PDAC versus healthy for the panel in the training set and in the independent validation set (30% of the data; PDAC  $n = 15$ , healthy  $n = 26$ ); C, summary table. SN and SP in the validation set are derived for optimal cutpoint determined in the training dataset. cnorm, creatinine-normalized; creat, creatinine.

Fig. S6C). The lack of LYVE1 immunoreactivity in the remaining cases, however, might be due to underrepresentation of relevant peripheral regions in our tissue microarray.

Next, we measured the levels of all three biomarkers in urine samples from 7 patients with PDAC for whom samples were collected before and after surgery (Supplementary Fig. S6D). In all patients, levels of LYVE1 and REG1A decreased after surgery, and this was also seen in 6 of 7 patients for TFF1 (except for patient 2, where the first postsurgical urine sample was collected 4 months after the procedure), likely due to substantial loss of tumor mass after surgery.


## Discussion

PDAC is one of the most challenging cancers to detect; the majority of patients thus present at an advanced stage of the disease. Hence, less than 20% of patients with PDAC undergo potentially curative surgery, whereas the remainder can only be offered palliative treatment.

Here, we describe a three-biomarker urine panel that discriminates patients with early-stage PDAC from healthy subjects with high accuracy. We chose to develop a diagnostic test based on urine specimens, as this body fluid has several advantages over blood: it is far less complex, provides an "inert" and stable matrix for analysis, and can be repeatedly and noninvasively sampled in sufficient volumes. So far, more than 2,300 proteins have been detected in urine (19), of which at least a third are of a systemic origin (20). As an ultrafiltrate of blood, it can be expected that at least some of the biomarkers will be found in higher concentration in urine than in blood.

Interestingly, when exploring the proteomes of healthy male and female urine samples, similarly as indicated previously (21), we observed large differences, indicating the need for the systematic exploration of gender-specific differences in biofluids.

Of the three biomarkers in our panel, REG1A and TFF1 have already been associated with PDAC. The **REG1A** gene product belongs to a family of REG (regenerating) glycoproteins, which are expressed in pancreatic acinar cells, and acts as both autocrine and paracrine growth factor (22, 23). The REG protein(s) are expressed during islet cell regeneration (24, 25), but are also potentially associated with differentiation and maintenance of an exocrine phenotype (26, 27). REG1 (nonspecified) expression has been reported previously in about 25% of PDACs ( $n = 20$ ; ref. 28), whereas we observed strong expression of REG1A in 73% of the cases examined here. REG1A and REG1B proteins are almost 90% identical (29) and are difficult to distinguish. REG1A levels have previously been observed in the plasma of mice with PanIN (Pancreatic intraepithelial neoplasia) lesions compared with control mice (30), and REG1B has recently been reported as one of the four candidate serological biomarkers that can improve the performance of CA19.9 in PDAC (31). Both proteins were able to significantly differentiate healthy from PDAC urine samples, demonstrating thus for the first time their high discriminatory power in urine specimens. TFF1 belongs to a family of gastrointestinal secretory peptides, which interact with mucins and are expressed at increased levels during reconstitution and repair of mucosal injury. They protect epithelial cells from apoptotic death and increase their motility, but also play similar pivotal roles in cancer cells, and are thus involved in the development and progression of various cancer types (32, 33). In PDAC, TFF1 has been reported in both sporadic (34, 35) and familial PanINs (36), and it has been associated with early stages (I and II) of the disease and disease without lymph node involvement, i.e., stage IIA (37). This expression mimics the pattern seen in our urine samples and confirms the biologic importance of TFF1 as an early diagnostic biomarker. Transcripts of all three members of



**Figure 5.**

Exploratory comparison of plasma CA19.9 and the urine biomarker panel in discriminating early patients with PDAC from healthy individuals. A, ROC curves of the biomarker panel with corresponding plasma CA19.9 alone and in combination comparing healthy urine ( $n = 28$ ), and urine samples from PDAC stages I-II,  $n = 71$ , and I-IIA,  $n = 16$  (B). C, summary table. SN and SP in the validation set were derived for optimal cutpoint determined in the training dataset.

Biomarkers	AUC (95% CI)	% SN (95% CI) at optimal cutpoint	% SP (95% CI) at optimal cutpoint
<b>PDAC I-II vs. healthy</b>			
Plasma CA19.9 (37 U/mL)	0.880 (0.814–0.945)	83.1 (74.6–91.6) <sup>^</sup>	92.9 (82.1–100.0) <sup>^</sup>
Panel	0.973 (0.947–0.999) <sup>+</sup>	93.0 (85.9–98.6)	92.9 (82.1–100.0)
Panel + Plasma CA19.9 (37 U/mL)	0.991 (0.979–1.000) <sup>\$</sup>	94.4 (88.7–98.6)	100.0 (100.0–100.0)
<b>PDAC I-IIA vs. healthy</b>			
Plasma CA19.9 (37 U/mL)	0.839 (0.719–0.959)	75.0 (56.3–93.8)	92.9 (82.1–100.0)
Panel	0.971 (0.929–1.000) <sup>++</sup>	87.5 (68.8–100.0)	96.4 (89.3–100.0)
Panel + Plasma CA19.9 (37 U/mL)	0.969 (0.924–1.000) <sup>\$\$</sup>	87.5 (68.8–100.0)	96.4 (89.3–100.0)

<sup>^</sup> Optimal cutpoint for CA19.9 is 37 U/mL

<sup>+</sup> DeLong's 1-sided test for correlated/paired AUCs to assess whether the urine panel gives a significantly greater AUC compared with plasma CA19.9 alone used as a dichotomous biomarker (0.973 vs. 0.880),  $P = 0.005$

<sup>\$</sup> DeLong's 1-sided test for correlated/paired AUCs to assess whether the addition of plasma CA19.9 used as a dichotomous biomarker significantly increase the AUC over the urine panel alone (0.991 vs. 0.973),  $P = 0.04$

<sup>++</sup> DeLong's 1-sided test for correlated/paired AUCs to assess whether the urine panel gives a significantly greater AUC compared with plasma CA19.9 alone used as a dichotomous biomarker (0.971 vs. 0.839),  $P = 0.006$

<sup>\$\$</sup> DeLong's 1-sided test for correlated/paired AUCs to assess whether the addition of plasma CA19.9 used as a dichotomous biomarker significantly increase the AUC over the urine panel alone (0.969 vs. 0.971),  $P = 0.7$

the family, and in particular TFF3, have been observed in the urinary tract of healthy subjects, but only TFF2 peptide was excreted in the urine (38). The increased TFF1 levels in urine samples from our cancer patients and its sharp decrease after surgical removal of the tumor further suggest that TFF1 protein originates from PDAC. LYVE1 (lymphatic vessel endothelial hyaluronan receptor) binds hyaluronan (HA), an extracellular matrix mucopolysaccharide, and transports it across the lymphatic vessel wall, particularly in the lymph nodes, a site of HA degradation (39). Most studies utilize LYVE1 as a marker of lymphatic endothelium in the context of lymphangiogenesis; however, markedly different findings regarding active lymphangiogenesis across different tumor types exist (40). No functional lymphangiogenesis is seen intratumorally in PDAC (41, 42), which may explain the infrequent presence of LYVE1 immunoreactive lymphatic vessels in our IHC analysis. It is also possible that LYVE1 expression was scarce due to the underrepresentation of extratumoral tissues within our case series, similarly to the findings of Carreira and colleagues, who reported a complete absence of LYVE1-positive lymphatic vessels in 25 hepatocellular carcinomas and 17 metastatic adenocarcinomas, but detected LYVE1 in the regions further away from the tumor nodules (43). LYVE1 showed potential also for differentiating IPMN, ampullary cancer, and NET from

early-stage PDAC samples; however, its role in PDAC still needs to be established.

When combined, REG1A, TFF1, and LYVE1 form a powerful urinary panel that can detect patients with stage I-II PDAC, with over 90% accuracy. To our knowledge, a panel with such performance has not been reported as yet. Our exploratory analyses also suggest that when combined with CA19.9, accuracy may be increased, which may prove important in light of recent finding that serum CA19.9 is upregulated up to 2 years before PDAC diagnosis (44). In addition, the panel may prove useful in discriminating patient in stages I-IIA from healthy ones, although this will need to be confirmed in a larger, independent study.

The strength of our study is manifested in multiple ways: the three biomarkers were discovered by MS analysis, validated using an independent technique (ELISA assays), and examined in urine specimens that originated from three different centers. This may facilitate the successful "portability" of our biomarker panel in future follow-up studies.

The diagnostic performance of our biomarker panel now needs to be further validated: our healthy controls were younger on average than our cancer patients; an older control group would thus be more relevant. In addition, further comparison of the performance of urine markers with CA19.9 is needed. Finally, it is now essential to establish if/how early in the latency period our



panel can detect PDAC, as soon as such a longitudinally collected urine cohort becomes available.

The high-risk groups for developing PDAC represent the priority cohort for screening strategies. These include pancreatic cancer families (FPC) with at least two affected first-degree relatives and individuals with hereditary syndromes with known underlying gene abnormalities, such as hereditary intestinal polyposis syndrome Peutz-Jeghers (*STK11/LKB1*), FAMMM, familial atypical multiple mole and melanoma (*p16/CDKN2A*), and hereditary pancreatitis (*PRSS1*, *SPINK1*; for a recent review, see 45). Although we have previously reported deregulation of *TFF1* and several *REG* genes in PanINs from FPC tissues (36), it is now critical to test *LYVE1*, and the panel as a whole, directly on urine samples collected from such high-risk individuals. This would best be performed within the framework of already established surveillance programs (46) in research setting. A detailed recommendation on management of individuals at high risk was recently reported by the Cancer of the Pancreas Screening (CAPS) consortium (47). We propose to include a urine test based on our three biomarkers into the predefined algorithms of the surveillance protocols to ultimately validate its performance. Ease of sampling and repeated testing using the urinary panel might also help shed some light on currently somewhat conflicting (but not mutually exclusive) data on the timescale of PDAC progression (48, 49).

It would also be worthwhile to explore the use of our urinary panel in clinical decision-making in individuals with various environmental exposures (smoking, obesity, new onset type II diabetes) which carry an increased risk of developing PDAC that could be predicted in primary care using models such as QCancer (Pancreas; ref. 50) or derived from the UK THIN database (51), but for which a consensus on screening has yet to be reached. Similarly, as shown in our study, urinary panel testing may prove useful and possibly superior to plasma CA19.9, in identifying cancer patients among patients with CP, another risk factor for developing PDAC (8). This performance should be further strengthened with additional marker(s) that could be selected from our proteomics analysis.

Being completely noninvasive and inexpensive, this urine screening test could, upon further validation, and when coupled with timely surgical intervention, lead to a much improved outcome in patients with high risk of developing PDAC.

## References

- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin* 2013;63:11–30.
- Malvezzi M, Bertuccio P, Levi F, La Vecchia C, Negri E. European cancer mortality predictions for the year 2013. *Ann Oncol* 2013;24:792–800.
- Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* 2014;74:2913–21.
- Tsuchiya R, Noda T, Harada N, Miyamoto T, Tomioka T, Yamamoto K, et al. Collective review of small carcinomas of the pancreas. *Ann Surg* 1986;203:77–81.
- Furukawa H, Okada S, Saisho H, Ariyama J, Karasawa E, Nakaizumi A, et al. Clinicopathologic features of small pancreatic adenocarcinoma. A collective study. *Cancer* 1996;78:986–90.
- Shimizu Y, Yasui K, Matsueda K, Yanagisawa A, Yamao K. Small carcinoma of the pancreas is curable: new computed tomography finding, pathological study and postoperative results from a single institute. *J Gastroenterol Hepatol* 2005;20:1591–4.
- Kaur S, Baine MJ, Jain M, Sasson AR, Batra SK. Early diagnosis of pancreatic cancer: challenges and new developments. *Biomark Med* 2012;6:597–612.
- Malka D, Hammel P, Maire F, Rufat P, Madeira I, Pessione F, et al. Risk of pancreatic adenocarcinoma in chronic pancreatitis. *Gut* 2002;51:849–52.
- Ballehaninna UK, Chamberlain RS. Serum CA 19-9 as a biomarker for pancreatic cancer—A comprehensive review. *Indian J Surg Oncol* 2011;2:88–100.
- Weeks ME, Hariharan D, Petronijevic L, Radon TP, Whiteman HJ, Kocher HM, et al. Analysis of the urine proteome in patients with pancreatic ductal adenocarcinoma. *Proteom Clin Appl* 2008;2:1047–57.
- Liu H, Sadygov RG, Yates JR 3rd. A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal Chem* 2004;76:4193–201.
- Ene-Obong A, Clear AJ, Watt J, Wang J, Fatah R, Riches JC, et al. Activated pancreatic stellate cells sequester CD8+ T cells to reduce their infiltration of the juxtatumoral compartment of pancreatic ductal adenocarcinoma. *Gastroenterology* 2013;145:1121–32.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Authors' Contributions

**Conception and design:** H.M. Kocher, J. Neoptolemos, N.R. Lemoine, T. Crnogorac-Jurcevic

**Development of methodology:** N.J. Massat, W. Alrawashdeh, J. Neoptolemos

**Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.):** T.P. Radon, R. Jones, D. Ennis, H.M. Kocher, S.P. Pereira, L. Guamer, C. Murta-Nascimento, N. Malats, J. Neoptolemos, W. Greenhalf

**Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis):** N.J. Massat, R. Jones, W. Alrawashdeh, S.W. Duffy, N. Malats, J. Neoptolemos, E. Costello, T. Crnogorac-Jurcevic

**Writing, review, and/or revision of the manuscript:** N.J. Massat, R. Jones, W. Alrawashdeh, L. Dumartin, S.W. Duffy, H.M. Kocher, S.P. Pereira, F.X. Real, N. Malats, J. Neoptolemos, E. Costello, W. Greenhalf, N.R. Lemoine, T. Crnogorac-Jurcevic

**Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases):** T.P. Radon, W. Alrawashdeh, L. Dumartin, D. Ennis, L. Guamer, J. Neoptolemos

**Study supervision:** J. Neoptolemos

**Other (provided samples for study):** F.X. Real

**Other (provision of clinical samples and data):** J. Neoptolemos

## Acknowledgments

The authors thank Chipso Chitsenga for help with sample collection. They are indebted to all the patients and healthy donors, without whom this study would not be possible. They also acknowledge the advice from Bendix Carstensen on the use of his Epi package.

## Grant Support

The study was funded by Pancreatic Cancer Research Fund (PCRF) and Queen Mary Innovation (to T. Crnogorac-Jurcevic), Cancer Research UK award 18975 (to N.R. Lemoine and T. Crnogorac-Jurcevic), The NIHR Liverpool Pancreas Biomedical Research Unit (to J. Neoptolemos, E. Costello, W. Greenhalf), The NIHR UCLH Biomedical Research Centre and NIH P01 CA084203 (to S.P. Pereira), and #PI09-02102, #PI12-00815, RD12/0036/0034, and RD12/0036/0050 from Instituto de Salud Carlos III, Madrid, Spain (to F.X. Real and N. Malats). H.M. Kocher, S.P. Pereira, F.X. Real, J. Neoptolemos, E. Costello, W. Greenhalf, T. Crnogorac-Jurcevic, and N. Malats are members of an EU Pancreas - BM1204 COST Action (<http://www.eupancreas.com>).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 30, 2014; revised March 27, 2015; accepted March 28, 2015; published online August 3, 2015.

13. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, et al. pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* 2011;12:77.
14. Rota M, Antolini L. Finding the optimal cut-point for Gaussian and Gamma distributed biomarkers. *Comput Stat Data An* 2014;69:1–14.
15. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837–45.
16. Carpenter J, Bithell J. Bootstrap confidence intervals: when, which, what? A practical guide for medical statisticians. *Stat Med* 2000;19:1141–64.
17. Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCr: visualizing classifier performance in R. *Bioinformatics* 2005;21:3940–1.
18. Dayem Ullah AZ, Cutts RJ, Ghetia M, Gadaleta E, Hahn SA, Crnogorac-Jurcevic T, et al. The pancreatic expression database: recent extensions and updates. *Nucl Acids Res* 2014;42:D944–9.
19. Kentsis A, Monigatti F, Dorff K, Campagne F, Bachur R, Steen H. Urine proteomics for profiling of human disease using high accuracy mass spectrometry. *Proteom Clin Appl* 2009;3:1052–61.
20. Jia L, Zhang L, Shao C, Song E, Sun W, Li M, et al. An attempt to understand kidney's protein handling function by comparing plasma and urine proteomes. *PLoS One* 2009;4:e5146.
21. Liu X, Shao C, Wei L, Duan J, Wu S, Li X, et al. An individual urinary proteome analysis in normal human beings to define the minimal sample number to represent the normal urinary proteome. *Proteome Sci* 2012;10:70.
22. Wu H, Zheng Q, Rengabhashyam P, Zenilman ME. A brief history of pancreatic reg: implications as to its clinical importance. *Einstein Q J Biol Med* 2000;17:178.
23. Zhang YW, Ding LS, Lai MD. Reg gene family and human diseases. *World J Gastroenterol* 2003;9:2635–41.
24. Watanabe T, Yonemura Y, Yonekura H, Suzuki Y, Miyashita H, Sugiyama K, et al. Pancreatic beta-cell replication and amelioration of surgical diabetes by Reg protein. *Proc Natl Acad Sci U S A* 1994;91:3589–92.
25. Parikh A, Stephan AF, Tzanakakis ES. Regenerating proteins and their expression, regulation and signaling. *Biomol Concepts* 2012;3:57–70.
26. Sanchez D, Mueller CM, Zenilman ME. Pancreatic regenerating gene I and acinar cell differentiation: influence on cellular lineage. *Pancreas* 2009;38:572–7.
27. Zenilman ME, Perfetti R, Swinson K, Magnuson T, Shuldiner AR. Pancreatic regeneration (reg) gene expression in a rat model of islet hyperplasia. *Surgery* 1996;119:576–84.
28. Kimura N, Yonekura H, Okamoto H, Nagura H. Expression of human regenerating gene mRNA and its product in normal and neoplastic human pancreas. *Cancer* 1992;70:1857–63.
29. Sanchez D, Figarella C, Marchand-Pinatel S, Bruneau N, Guy-Crotte O. Preferential expression of reg I beta gene in human adult pancreas. *Biochem Biophys Res Commun* 2001;284:729–37.
30. Faca VM, Song KS, Wang H, Zhang Q, Krasnoselsky AL, Newcomb LF, et al. A mouse to human search for plasma proteome changes associated with pancreatic tumor development. *PLoS Med* 2008;5:e123.
31. Makawita S, Dimitromanolakis A, Soosaipillai A, Soleas I, Chan A, Gallinger S, et al. Validation of four candidate pancreatic cancer serological biomarkers that improve the performance of CA19.9. *BMC Cancer* 2013;13:404.
32. Emami S, Rodrigues S, Rodrigue CM, Le Floch N, Rivat C, Attoub S, et al. Trefoil factor family (TFF) peptides and cancer progression. *Peptides* 2004;25:885–98.
33. Perry JK, Kannan N, Grandison PM, Mitchell MD, Lobie PE. Are trefoil factors oncogenic? *Trends Endocrinol Metab* 2008;19:74–81.
34. Arumugam T, Brandt W, Ramachandran V, Moore TT, Wang H, May FE, et al. Trefoil factor 1 stimulates both pancreatic cancer and stellate cells and increases metastasis. *Pancreas* 2011;40:815–22.
35. Prasad NB, Biankin AV, Fukushima N, Maitra A, Dhara S, Elkahloun AG, et al. Gene expression profiles in pancreatic intraepithelial neoplasia reflect the effects of Hedgehog signaling on pancreatic ductal epithelial cells. *Cancer Res* 2005;65:1619–26.
36. Crnogorac-Jurcevic T, Chelala C, Barry S, Harada T, Bhakta V, Lattimore S, et al. Molecular analysis of precursor lesions in familial pancreatic cancer. *PLoS One* 2013;8:e54830.
37. Sagol O, Tuna B, Coker A, Karademir S, Obuz F, Astarcioğlu H, et al. Immunohistochemical detection of pS2 protein and heat shock protein-70 in pancreatic adenocarcinomas. Relationship with disease extent and patient survival. *Pathol Res Pract* 2002;198:77–84.
38. Rinnert M, Hinz M, Buhtz P, Reiher F, Lessel W, Hoffmann W. Synthesis and localization of trefoil factor family (TFF) peptides in the human urinary tract and TFF2 excretion into the urine. *Cell Tissue Res* 2010;339:639–47.
39. Jackson DG. Immunological functions of hyaluronan and its receptors in the lymphatics. *Immunol Rev* 2009;230:216–31.
40. Jackson DG, Prevo R, Clasper S, Banerji S. LYVE-1, the lymphatic system and tumor lymphangiogenesis. *Trends Immunol* 2001;22:317–21.
41. Sipos B, Kojima M, Tiemann K, Klapper W, Kruse ML, Kalthoff H, et al. Lymphatic spread of ductal pancreatic adenocarcinoma is independent of lymphangiogenesis. *J Pathol* 2005;207:301–12.
42. Schneider M, Buchler P, Giese N, Giese T, Wilting J, Buchler MW, et al. Role of lymphangiogenesis and lymphangiogenic factors during pancreatic cancer progression and lymphatic spread. *Int J Oncol* 2006;28:883–90.
43. Mouta Carreira C, Nasser SM, di Tomaso E, Padera TP, Boucher Y, Tomarev SI, et al. LYVE-1 is not restricted to the lymph vessels: expression in normal liver blood sinusoids and down-regulation in human liver cancer and cirrhosis. *Cancer Res* 2001;61:8079–84.
44. O'Brien DP, Sandanayake NS, Jenkinson C, Gentry-Maharaj A, Apostolidou S, Fourkala EO, et al. Serum CA19–9 is significantly upregulated up to 2 years before diagnosis with pancreatic cancer: implications for early disease detection. *Clin Cancer Res* 2014;21:622–31.
45. Templeton AW, Brentnall TA. Screening and surgical outcomes of familial pancreatic cancer. *Surg Clin North Am* 2013;93:629–45.
46. Wada K, Takaori K, Traverso LW, Hruban RH, Furukawa T, Brentnall TA, et al. Clinical importance of Familial Pancreatic Cancer Registry in Japan: a report from kick-off meeting at International Symposium on Pancreas Cancer 2012. *J Hepatobiliary Pancreat Sci* 2013;20:557–66.
47. Canto MI, Harinck F, Hruban RH, Offerhaus GJ, Poley JW, Kamel I, et al. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. *Gut* 2013;62:339–47.
48. Yachida S, Jones S, Bozic I, Antal T, Leary R, Fu B, et al. Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature* 2010;467:1114–7.
49. Yu J, Blackford AL, Dal Molin M, Wolfgang CL, Goggins M. Time to progression of pancreatic ductal adenocarcinoma from low-to-high tumour stages. *Gut* 2015;0:1–7.
50. Hippisley-Cox J, Coupland C. Identifying patients with suspected pancreatic cancer in primary care: derivation and validation of an algorithm. *Br J Gen Pract* 2012;62:e38–45.
51. Keane MG, Horsfall L, Rait G, Pereira SP. A case-control study comparing the incidence of early symptoms in pancreatic and biliary tract cancer. *BMJ Open* 2014;4:e005720.

# Clinical Cancer Research

## Identification of a Three-Biomarker Panel in Urine for Early Detection of Pancreatic Adenocarcinoma

Tomasz P. Radon, Nathalie J. Massat, Richard Jones, et al.

*Clin Cancer Res* 2015;21:3512-3521.

<b>Updated version</b>	Access the most recent version of this article at: <a href="http://clincancerres.aacrjournals.org/content/21/15/3512">http://clincancerres.aacrjournals.org/content/21/15/3512</a>
<b>Supplementary Material</b>	Access the most recent supplemental material at: <a href="http://clincancerres.aacrjournals.org/content/suppl/2015/07/23/21.15.3512.DC1">http://clincancerres.aacrjournals.org/content/suppl/2015/07/23/21.15.3512.DC1</a>

<b>Cited articles</b>	This article cites 51 articles, 8 of which you can access for free at: <a href="http://clincancerres.aacrjournals.org/content/21/15/3512.full#ref-list-1">http://clincancerres.aacrjournals.org/content/21/15/3512.full#ref-list-1</a>
<b>Citing articles</b>	This article has been cited by 7 HighWire-hosted articles. Access the articles at: <a href="http://clincancerres.aacrjournals.org/content/21/15/3512.full#related-urls">http://clincancerres.aacrjournals.org/content/21/15/3512.full#related-urls</a>

<b>E-mail alerts</b>	<a href="#">Sign up to receive free email-alerts</a> related to this article or journal.
<b>Reprints and Subscriptions</b>	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at <a href="mailto:pubs@aacr.org">pubs@aacr.org</a> .
<b>Permissions</b>	To request permission to re-use all or part of this article, use this link <a href="http://clincancerres.aacrjournals.org/content/21/15/3512">http://clincancerres.aacrjournals.org/content/21/15/3512</a> . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.