

# Prognostic Value of $\alpha 6\beta 4$ Integrin Expression in Breast Carcinomas Is Affected by Laminin Production from Tumor Cells<sup>1</sup>

Elda Tagliabue, Cristina Ghirelli,  
Paolo Squicciarini, Pierangela Aiello,  
Maria Ines Colnaghi,<sup>2</sup> and Sylvie Ménard

Divisions of Experimental Oncology E [E. T., C. G., P. A., M. I. C., S. M.] and Surgical Oncology C [P. S.], Istituto Nazionale Tumori, 20133 Milan, Italy

## ABSTRACT

Immunocytochemical analysis of breast carcinoma specimens for  $\alpha 6\beta 4$  integrin expression and other different pathobiological markers revealed  $\beta 4$  integrin subunit expression in 36 of 80 cases analyzed and a significant association only with  $\alpha 6$  integrin subunit expression ( $P < 0.01$ ) and laminin production ( $P = 0.01$ ) by tumor cells. Survival analysis indicated that  $\beta 4$  and  $\alpha 6$  expression are associated with poor prognosis ( $P = 0.02$ ), whereas laminin production showed only borderline association ( $P = 0.06$ ). However, analysis of disease outcome in relation to expression of both  $\alpha 6\beta 4$  and laminin indicated best outcomes for patients with tumors producing laminin but not expressing  $\alpha 6\beta 4$  integrin, whereas worst outcomes were observed for  $\alpha 6\beta 4$ - and laminin-positive tumors, indicating that  $\alpha 6\beta 4$  expression was associated with prognosis, mainly in the laminin-producing tumor subset. These data indicate that the prognostic value of  $\alpha 6\beta 4$  integrin expression is affected by laminin production from tumor cells and suggest that interaction between these two molecules mediates distinct signals that are important for tumor progression.

## INTRODUCTION

The  $\alpha 6\beta 4$  integrin, corresponding to the TSP-180 tumor-associated complex that was described in mouse mammary tumor cells and human carcinoma cell lines (1-3), is the receptor for various laminin isoforms (4, 5) and is mainly expressed in cell types that interact with laminin, *i.e.*, epithelial (6), endothelial (7, 8), and Schwann cells (9). Moreover,  $\alpha 6\beta 4$ , by virtue of a specific tyrosine activation motif in the  $\beta 4$  tail, plays a crucial role in the assembly of hemidesmosomes (10), the junctions connecting the basal cells of stratified and complex epithelia to the basement membrane (11). The basal polarization of this molecular complex

in normal epithelia results in its gradual loss in various solid tumors (12-16). The role of the  $\alpha 6\beta 4$  integrin in tumor progression is still unclear. Indeed, several studies indicate that  $\alpha 6\beta 4$  is overexpressed in squamous carcinomas of lung, skin, oral cavity, and cervix (14, 15, 17, 18), whereas it is down-regulated in adenocarcinomas derived from epithelial tissues that normally express high levels of this integrin, *e.g.*, adenocarcinomas of breast and prostate (19-21). Recent *in vitro* studies indicate that  $\alpha 6\beta 4$ , which does not itself have an intracellular catalytic domain, is physically and functionally associated with cytoplasmic tyrosine kinases (7, 22) and that laminin binding to this integrin activates the associated kinase and consequently tyrosine phosphorylation of the  $\beta 4$  subunit cytoplasmic domain. Tyrosine phosphorylation motifs in the  $\beta 4$  tail, distinct from those involved in the assembly of hemidesmosomes, mediate the recruitment of Shc/Grb2 and may link the integrin to ras signaling pathways (10). Because the mitogen-activated protein kinase pathway activated by ras has been implicated in controlling proliferation or differentiation depending on the cellular context (23), the linkage of  $\alpha 6\beta 4$  to ras may explain the effects of laminins on morphogenesis and tumor progression. Recently,  $\alpha 6\beta 4$  overexpression in rectal carcinoma cells was found to induce p21<sup>WAF1/CIP1</sup> expression with consequent cell growth arrest but also with increased invasiveness (24). These contrasting results support the hypothesis that  $\alpha 6\beta 4$  is linked to a growth suppression pathway and that laminin binding reverses this connection and couples the integrin through Shc and Grb2 to ras with consequent cell proliferation and/or invasion. On the basis of these *in vivo* and *in vitro* studies, we analyzed breast carcinomas for expression of the  $\alpha 6\beta 4$  integrin and other pathobiological markers and the prognostic value of the  $\alpha 6\beta 4$  in relation to the laminin produced by the tumor cells.

## PATIENTS AND METHODS

**Patients.** The study included 80 patients who were surgically treated at Istituto Nazionale Tumori (Milan, Italy) from 1988 to 1990: 51% showed no lymph node involvement and 49% presented a tumor that was  $\leq 2$  cm. Surgical treatment consisted of radical or modified radical mastectomy and axillary dissection. Histologically node-positive patients received post-surgical adjuvant treatment.

**Immunocytochemical Staining.** Frozen sections fixed in acetone were stained by the immunoperoxidase method using the avidin-biotin complex kit (Vector Laboratories, Burlingame, CA), and reactivity was detected with the following MAb<sup>3</sup>: DO7, directed against p53 protein (Ylem-Novocastra, New Castle, United Kingdom); clone 124, directed against bcl-2 product (DAKO A/S, Glostrup, Denmark); MGR1, directed against the EGFR (25); MGR2, directed against p185<sup>HER2</sup> (26); MAR6, directed against

Received 7/22/97; revised 11/14/97; accepted 11/21/97.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was partially supported by a grant from the Associazione Italiana Ricerca sul Cancro/Fondazione Italiana Ricerca sul Cancro.


<sup>2</sup> To whom requests for reprints should be addressed, at Oncologia Sperimentale E, Istituto Nazionale per lo Studio e la Cura dei Tumori, Via G. Venezian 1, 20133 Milan, Italy. Phone: 39-2-2390566; Fax: 39-2-2362692; E-mail: colnaghi@istitutotumori.mi.it.

<sup>3</sup> The abbreviations used are: MAb, monoclonal antibody; EGFR, epidermal growth factor receptor.

**Table 1** Analysis of association between  $\beta 4$  integrin subunit expression and other evaluated parameters

Parameter	Category	No. of $\beta 4$ + cases/no. of tested cases	%	<i>P</i>
Size	$\leq 2$ cm	23/45	51	<0.01
	>2 cm	12/34	35	
Node	+	21/44	48	
	-	15/36	42	
Grade	I-II	28/59	47	
	III	8/20	40	
ER <sup>a</sup>	+	26/62	42	
	-	7/12	58	
PGR	+	21/49	43	
	-	12/25	48	
LI	+	18/35	55	
	-	7/15	47	
p185 <sup>HER2</sup>	+	13/29	45	
	-	21/47	45	
EGFR	+	8/17	47	
	-	26/59	42	
p53	+	8/11	73	
	-	28/65	43	
bcl-2	+	21/40	52	
	-	10/27	37	
$\alpha 6$	+	32/34	94	
	-	3/45	7	
LN	+	18/27	67	0.01
	-	14/40	35	

<sup>a</sup> ER, estrogen receptor; PGR, progesterone receptor; LI, lymphoid infiltration; LN, laminin.



**Fig. 1** Immunocytochemical analysis of frozen sections of breast carcinoma with Mab against  $\beta 4$  (a) or polyclonal serum against laminin (b).

the  $\alpha 6$  integrin chain (12); and 3E1, directed against the  $\beta 4$  integrin chain (Telios, San Diego, CA). Rabbit serum directed against human placental laminin (Telios) was used.

Tumors were considered positive when more than 20% of cells were reactive with the Mab tested.

**Statistical Analysis.** Differences in frequencies were analyzed by the  $\chi^2$  test with Yates correction. The survival curve

**Table 2** Analysis of relapses and deaths according to different pathobiological parameters

Parameter	No. of cases	% of relapse	<i>P</i>	% of death	<i>P</i>
Node	+	44	0.14	16	0.01
	-	36		19	
Size	$\leq 2$ cm	45	0.03	4	0.2
	>2 cm	24		42	
p165 <sup>HER2</sup>	-	46	0.15	7	0.3
	+	30		37	
$\alpha 6$	-	45	0.7	2	0.02
	+	35		34	
$\beta 4$	-	44	0.3	2	0.02
	+	36		33	
Laminin	-	46	0.2	4	0.06
	+	30		20	

**Table 3** Analysis of relapses and deaths according to laminin production and  $\alpha 6\beta 4$  expression

Laminin production	$\alpha 6\beta 4$ expression	No. of cases	% relapse	% death	<i>P</i>
+	+	19	32	26	<0.01
	-	11	0	0	
-	+	16	37	6	NS <sup>a</sup>
	-	30	33	3	


<sup>a</sup> NS, not significant.

was calculated from the date of surgery, using death from cancer only (overall survival) and recurrence (disease-free survival) as end points. Survival rates were calculated using the actuarial life table method, considering the subgroups identified by the variables examined, and compared using the log-rank test.

## RESULTS

Expression of the  $\beta 4$  integrin subunit was detected in 36 of the 80 cases (45%). The expression of  $\beta 4$  integrin subunit was then analyzed in relation to clinical and pathological parameters, such as tumor size, nodal status, degree of tumor differentiation, lymphoid infiltration, and a panel of biological markers associated with tumor progression.  $\beta 4$  expression was found to be strongly associated with  $\alpha 6$  integrin subunit expression ( $P < 0.01$ ) and laminin production by tumor cells ( $P = 0.01$ ), whereas, except for a borderline correlation with p53 positivity ( $P = 0.06$ ),  $\beta 4$  expression was not related to any of tumor progression markers tested (Table 1). All cases expressing the  $\alpha 6\beta 4$  integrin showed pericellular staining, with a positivity ranging from 50 to 100% of

**Fig. 2** Disease-free (A) and overall (B) survival rates of patients with breast carcinomas according to  $\beta 4$  expression and laminin production. ●,  $\alpha 6\beta 4$ -negative, laminin-positive cases; ■,  $\alpha 6\beta 4$ -positive, laminin-positive cases; ▼,  $\alpha 6\beta 4$ -negative, laminin-negative cases; ▲,  $\alpha 6\beta 4$ -positive, laminin-negative cases. A, ● versus ■,  $P = 0.04$ ; ● versus ▲,  $P = 0.03$ . B, ● versus ■,  $P = 0.07$ .



tumor cells (Fig. 1a), whereas tumors considered laminin positive showed specific staining at the membrane and/or the cytoplasmic level on the majority of tumor cells (Fig. 1b).

The frequency of relapses and deaths for tumor was then analyzed according to pathological and biological parameters. The median follow-up of these cases was 7 years. Significant correlations between relapses and tumor size ( $P = 0.03$ ) and between deaths and nodal status ( $P = 0.01$ ) were found. The expressions of both  $\alpha 6$  and  $\beta 4$  were significantly associated with death ( $P = 0.02$ ), whereas laminin production was only borderline significant ( $P = 0.06$ ; Table 2). Analysis of disease outcome in relation to expression of  $\alpha 6\beta 4$  and laminin revealed significant differences in relapse and survival (Table 3). The best outcome was observed for patients with tumors producing laminin but not expressing  $\alpha 6\beta 4$  integrin, whereas the worst outcome was found for  $\alpha 6\beta 4$ -positive, laminin-positive tumors. The other two groups, both negative for laminin production, showed a relapse rate comparable to that of  $\alpha 6\beta 4$ -positive, laminin-positive tumors, but they showed a lower death frequency (Fig. 2).

Analysis of these four tumor subsets for pathobiological features, including nodal status, tumor size, and other markers, such as p185<sup>HER2</sup> and p53, revealed no significant differences. Indeed, each subset presented as follows: 80% of cases had tumor sizes of  $\leq 2$  cm, grades I–II, and hormone receptor positivity; 50% of cases had lymph node infiltration and p185<sup>HER2</sup> and bcl-2 expression; and 20% of cases expressed EGFR. p53 positivity was higher in  $\alpha 6\beta 4$ -positive tumors (20%) than it was in  $\alpha 6\beta 4$ -negative tumors (10%), independent of laminin production by tumor cells.

## DISCUSSION

The present data show that breast carcinomas producing laminin also frequently express the  $\beta 4$  integrin subunit in association with the  $\alpha 6$  subunit. The finding that  $\beta 4$  is consistently expressed in association with  $\alpha 6$  is in keeping with observations in normal tissues, where coexpressed  $\alpha 6$  and  $\beta 4$  always associate together, independent of the presence of  $\beta 1$  subunit (13, 15). This suggests that the frequency of breast carcinomas expressing  $\alpha 6\beta 1$  is very low and that this integrin has only marginal relevance in tumor progression. Expression of the  $\alpha 6\beta 4$  integrin is maintained mainly in tumors producing laminin, raising the possibility of an autocrine regulatory mechanism in which laminin produced by tumor cells inter-

feres with  $\alpha 6\beta 4$  expression and/or redistribution. Because the antiserum used in the present study is directed against human placenta laminin, we actually analyzed the production by tumor cells of all of the different laminin isoforms presented in placenta, including kalinin/laminin 5, which is also recognized by  $\alpha 6\beta 4$  (5) and is detected only in invading cells of the tumors (27, 28). However, the antilaminin staining observed in our series was distributed throughout the tumor areas with no apparent correlation to invasive malignant cells. Preliminary data indicate that only the tumors stained with the antilaminin polyclonal antibody were also positive with an antilaminin 5 MAb, but the staining was confined to a very small proportion of tumor cells. These findings suggest a coordinated regulation of laminin isoform production in breast carcinomas.

$\alpha 6\beta 4$  expression in breast tumor cells, unlike that in ovary carcinomas (12), was found to be independent of tumor grade or other prognostic factors but was nevertheless significantly associated with the worst patient outcome ( $P = 0.02$ ). This is in keeping with the finding that restoration of  $\alpha 6\beta 4$  receptor expression in a  $\beta 4$ -deficient rectal carcinoma cell line facilitates its ability to invade both matrigel and collagen I matrices (29). Moreover, high-level expression of  $\alpha 6\beta 4$  on mammary carcinoma cells was found to contribute to liver metastasis formation by mediating adhesion of the cancer cells to hepatocytes (30). Interestingly, 4 of the 5 liver metastases observed thus far in our series derived from  $\alpha 6\beta 4$ -positive primary tumors. However, our findings point to the crucial role of the  $\alpha 6\beta 4$  integrin in tumor progression only when it can interact with its appropriate substrate, supporting the hypothesis that laminin(s) also plays an active role in tumor progression. Indeed, although the patients with laminin-positive,  $\beta 4$ -positive tumors showed a recurrence rate similar to that of patients with laminin-negative tumors, they clearly had a higher death frequency (80% of relapse patients) than did the laminin-negative group (12% of relapse patients). The absence of significant differences between the different group of tumor patients in overall survival are most likely due to the low number of events thus far. It has been hypothesized that ligand-bound and unbound  $\alpha 6\beta 4$  integrins activate distinct intracellular pathways (31). Indeed, laminin-induced binding of the  $\alpha 6\beta 4$  receptor was found to induce tyrosine phosphorylation of the  $\beta 4$  subunit followed by recruitment of Shc/Grb2, thus linking the integrin activation to the ras pathway of proliferation. The lack of aggressiveness by laminin-positive,  $\alpha 6\beta 4$ -neg-

ative tumors may reflect the binding of the adhesion molecule to other laminin receptors that mediate adhesion rather than proliferation and/or migration. The slight increase in relapses seen in patients with laminin-negative,  $\alpha 6 \beta 4$ -positive tumors suggests that  $\alpha 6 \beta 4$  integrin in these cases interacts with laminin produced by the host, including these patients in the category of the laminin-positive,  $\alpha 6 \beta 4$ -positive cases.

Consistent with previous analyses indicating a correlation between  $\alpha 6$  expression and reduced survival (32), our data emphasize the importance of the interaction of laminin with  $\alpha 6 \beta 4$  rather than  $\alpha 6 \beta 4$  expression *per se* in influencing breast carcinoma progression. Specific up- and down-regulation of the expression of laminins and integrins by differentiating agents and cytokines, respectively, may help in modulating tumor aggressiveness.

## ACKNOWLEDGMENTS

We thank Laura Mameli for manuscript preparation and Mario Azzini for photographic reproduction.

## REFERENCES

- Kennel, S. J., Foote, L. J., Falcioni, R., Sonnenberg, A., Stringer, C. D., Crouse, C., and Hemler, M. E. Analysis of the tumor-associated antigen TSP-180: identity with  $\alpha 6 \beta 4$  in the integrin superfamily. *J. Biol. Chem.*, *264*: 15515–15521, 1989.
- Hemler, M. E., Crouse, C., and Sonnenberg, A. Association of the VLA  $\alpha 6$  subunit with a novel protein: a possible alternative to the common VLA b1 subunit on certain cell lines. *J. Biol. Chem.*, *264*: 6529–6535, 1989.
- Sonnenberg, A. F., Hogervorst, F., Osterop, A., and Veltman, F. E. M. Identification and characterization of a novel antigen complex on mouse mammary tumor cells using a monoclonal antibody against platelet glycoprotein I<sub>c</sub>. *J. Biol. Chem.*, *263*: 14030–14038, 1988.
- Lee, E. C., Lotz, M. M., Steele, G. D., Jr., and Mercurio, A. M. The integrin  $\alpha 6 \beta 4$  is a laminin receptor. *J. Cell Biol.*, *117*: 671–678, 1992.
- Niessen, C. M., Hogervorst, F., Jaspars, L. H., De Melker, A. A., Delwel, G. O., Hulsman, E. H. M., Kuikman, I., and Sonnenberg, A. The  $\alpha 6 \beta 4$  integrin is a receptor for both laminin and kalinin. *Exp. Cell Res.*, *211*: 360–367, 1994.
- Kajiji, S., Tamura, R. N., and Quaranta, V. A novel integrin ( $\alpha E \beta 4$ ) from human epithelial cells suggests a fourth family of integrin adhesion receptors. *EMBO J.*, *8*: 673–680, 1989.
- Kennel, S. J., Godfrey, V., Chang, L. Y., Lankfork, T. K., Foote, L. J., and Makkinje, A. The integrin  $\beta 4$  subunit is displayed on a restricted subset of endothelium in mice. *J. Cell Sci.*, *101*: 145–150, 1992.
- Klein, S., Giancotti, F. G., Presta, M., Alberta, S. M., Buck, C. A., and Rifkin, D. B. Basic fibroblast growth factor modulates integrin expression in microvascular endothelial cells. *Mol. Biol. Cell.*, *4*: 973–982, 1993.
- Sonnenberg, A., and Linders, C. J. T. The  $\alpha 6 \beta 1$  (VLA-6) and  $\alpha 6 \beta 4$  protein complexes: tissue distribution and biochemical properties. *J. Cell Sci.*, *96*: 207–217, 1990.
- Mainiero, F., Pepe, A., Wary, K. K., Spinardi, L., Mohammadi, M., Schlessinger, J., and Giancotti, F. G. Signal transduction by the  $\alpha 6 \beta 4$  integrin: distinct  $\beta 4$  subunit sites mediate recruitment of Shc/Grb2 and association with the cytoskeleton of hemidesmosomes. *EMBO J.*, *14*: 4470–4481, 1995.
- Legan, P. K., Collins, J. E., and Garrod, D. R. The molecular biology of desmosomes and hemidesmosomes: "what's in a name?". *BioEssays*, *14*: 385–393, 1992.
- Bottini, C., Miotti, S., Fiorucci, S., Facheris, P., Ménard, S., and Colnaghi, M. I. Polarization of the  $\alpha 6 \beta 4$  integrin in ovarian carcinomas. *Int. J. Cancer*, *54*: 261–267, 1993.
- Natali, P. G., Nicotra, M. R., Bigotti, A., and De Martino, C. Localization of the  $\alpha 6$  and  $\beta 4$  integrin subunits in normal human non-lymphoid tissues. *J. Cell Sci.*, *103*: 1243–1247, 1992.
- Mariani-Costantini, R., Falcioni, R., Battista, P., Zupi, G., Kennel, S. J., Colasante, A., Venturo, I., Curcio, C. G., and Sacchi, A. Integrin ( $\alpha 6$ ,  $\beta 4$ ) expression in human lung cancer as monitored by specific monoclonal antibodies. *Cancer Res.*, *50*: 6107–6112, 1990.
- Kimmel, K. A., and Carey, T. E. Altered expression in squamous carcinoma cells of an orientation restricted epithelial antigen detected by monoclonal antibody A9. *Cancer Res.*, *46*: 3614–3623, 1986.
- Van Waes, C., Kozarsky, K. F., Warren, A. B., Kidd, L., Paugh, D., Liebert, M., and Carey, T. E. The A9 antigen associated with aggressive human squamous carcinoma is structurally and functionally similar to the newly defined integrin  $\alpha 6 \beta 4$ . *Cancer Res.*, *51*: 2395–2402, 1991.
- Carico, E., French, D., Bucci, B., Falcioni, R., Vecchione, A., and Mariani-Costantini, R. Integrin  $\beta 4$  expression in the neoplastic progression of cervical epithelium. *Gynecol. Oncol.*, *49*: 61–66, 1993.
- Wolf, G. T., Carey, T. E., Schmaltz, S. P., McClatchey, K. D., Poor Glaser, L., Hayashida, D. J. S., and Hsu, S. Altered antigen expression predicts outcome in squamous cell carcinomas of the head and neck. *J. Natl. Cancer Inst.*, *82*: 1566–1572, 1990.
- Knox, J. D., Cress, A. E., Clark, V., Manriquez, L., Affinito, K.-S., Dalkin, B. L., and Nagle, R. B. Differential expression of extracellular matrix molecules and the  $\alpha 6$ -integrins in the normal and neoplastic prostate. *Am. J. Pathol.*, *145*: 167–174, 1994.
- Koukoulis, G. K., Virtanen, I., Korhonen, M., Laitinen, L., Quaranta, V., and Gould, V. E. Immunohistochemical localization of integrins in the normal, hyperplastic, and neoplastic breast: correlations with their functions as receptors and cell adhesion molecules. *Am. J. Pathol.*, *139*: 787–799, 1991.
- Natali, P. G., Nicotra, M. R., Botti, C., Mottolese, M., Bigotti, A., and Segatto, O. Changes in expression of  $\alpha 6 \beta 4$  integrin heterodimer in primary and metastatic breast cancer. *Br. J. Cancer*, *66*: 318–322, 1992.
- Weiss, A., and Littman, D. R. Signal transduction by lymphocyte antigen receptors. *Cell*, *76*: 263–274, 1994.
- Cowley, S., Paterson, H., Kemp, P., and Marshall, C. J. Activation of MAP kinase is necessary and sufficient for PC12 differentiation and for transformation of NIH 3T3 cells. *Cell*, *77*: 841–852, 1994.
- Clark, E. A., and Brugge, J. S. Integrins and signal transduction pathways: the road taken. *Science (Washington DC)*, *268*: 233–239, 1995.
- Pellegrini, R., Centis, F., Martignone, S., Mastroianni, A., Tagliabue, E., Tosi, E., Ménard, S., and Colnaghi, M. I. Characterization of a monoclonal antibody directed against the epidermal growth factor receptor binding site. *Cancer Immunol. Immunother.*, *34*: 37–42, 1991.
- Tagliabue, E., Centis, F., Campiglio, M., Mastroianni, A., Martignone, S., Pellegrini, R., Casalini, P., Lanzi, C., Ménard, S., and Colnaghi, M. I. Selection of monoclonal antibodies which induce internalization and phosphorylation of p185<sup>HER2</sup> and growth inhibition of cells with HER2/neu gene amplification. *Int. J. Cancer*, *47*: 933–937, 1991.
- Pyke, C., Romer, J., Kallunki, P., Lund, L. R., Ralfkiaer, E., Dano, K., and Tryggvason, K. The  $\gamma 2$  chain of kalinin/laminin 5 is preferentially expressed in invading malignant cells in human cancers. *Am. J. Pathol.*, *145*: 782–791, 1994.
- Pyke, C., Sale, S., Ralfkiaer, E., Romer, J., Dano, K., and Tryggvason, K. Laminin-5 is a marker of invading cancer cells in some human carcinomas and is coexpressed with the receptor for urokinase plasminogen activator in budding cancer cells in colon adenocarcinomas. *Cancer Res.*, *55*: 4132–4139, 1995.
- Chao, C., Lotz, M. M., Clarke, A. C., and Mercurio, A. M. A function for the integrin  $\alpha 6 \beta 4$  in the invasive properties of colorectal carcinoma cells. *Cancer Res.*, *56*: 4811–4819, 1996.
- Kemperman, H., Wijnands, Y., De Rijk, D., and Roos, E. The integrin  $\alpha 6 \beta 4$  on TA3/Ha mammary carcinoma cells is involved in adhesion to hepatocytes. *Cancer Res.*, *53*: 3611–3617, 1993.
- Giancotti, F. G. Signal transduction by the  $\alpha 6 \beta 4$  integrin: charting the path between laminin binding and nuclear events. *J. Cell Sci.*, *109*: 1165–1172, 1996.
- Friedrichs, K., Ruiz, P., Franke, F., Gille, I., Terpe, H.-J., and Imhof, B. A. High expression level of  $\alpha 6$  integrin in human breast carcinoma is correlated with reduced survival. *Cancer Res.*, *55*: 901–906, 1995.

# Clinical Cancer Research

## Prognostic value of alpha 6 beta 4 integrin expression in breast carcinomas is affected by laminin production from tumor cells.

E Tagliabue, C Ghirelli, P Squicciarini, et al.

*Clin Cancer Res* 1998;4:407-410.

**Updated version** Access the most recent version of this article at:  
<http://clincancerres.aacrjournals.org/content/4/2/407>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://clincancerres.aacrjournals.org/content/4/2/407>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.