

Antitumor Activity of Sequential Treatment with Topotecan and Anti-Epidermal Growth Factor Receptor Monoclonal Antibody C225¹

Fortunato Ciardiello,² Roberto Bianco, Vincenzo Damiano, Sonya De Lorenzo, Stefano Pepe, Sabino De Placido, Zhen Fan, John Mendelsohn, A. Raffaele Bianco, and Giampaolo Tortora

Cattedra di Oncologia Medica, Dipartimento di Endocrinologia e Oncologia Molecolare e Clinica, Facoltà di Medicina e Chirurgia, Università degli Studi di Napoli Federico II, Naples, Italy [F. C., R. B., V. D., S. D. L., S. P., S. D. P., A. R. B., G. T.], and University of Texas M. D. Anderson Cancer Center, Houston, Texas [Z. F., J. M.]

ABSTRACT

Epidermal growth factor (EGF)-related proteins such as transforming growth factor α (TGF- α) control cancer cell growth through autocrine and paracrine pathways. Overexpression of TGF- α and/or its receptor (EGFR) has been associated with a more aggressive disease and a poor prognosis. The blockade of EGFR activation has been proposed as a target for anticancer therapy. Monoclonal antibody (MAb) C225 is an anti-EGFR humanized chimeric mouse MAb that is presently in Phase II clinical trials in cancer patients. Previous studies have suggested the potentiation of the antitumor activity of certain cytotoxic drugs, such as cisplatin and doxorubicin, in human cancer cell lines by treatment with anti-EGFR antibodies. We have evaluated in human ovarian, breast, and colon cancer cell lines, which express functional EGFR, the antiproliferative activity of MAb C225 in combination with topotecan, a cytotoxic drug that specifically inhibits topoisomerase I and that has shown antitumor activity in these malignancies. A dose-dependent supraadditive increase of growth inhibition *in vitro* was observed when cancer cells were treated with topotecan and MAb C225 in a sequential schedule. In this respect, the cooperativity quotient, defined as the ratio between the actual growth inhibition obtained by treatment with topotecan followed by MAb C225 and the sum of the growth inhibition

achieved by each agent, ranged from 1.2 to 3, depending on drug concentration and cancer cell line. Treatment with MAb C225 also markedly enhanced apoptotic cell death induced by topotecan. For example, in GEO colon cancer cells, 5 nM topotecan, followed by 0.5 μ g/ml MAb C225, induced apoptosis in 45% cells as compared with untreated cells (6%) or to 5 nM topotecan-treated cells (22%). Treatment of mice bearing established human GEO colon cancer xenografts with topotecan or with MAb C225 determined a transient inhibition of tumor growth because GEO tumors resumed the growth rate of untreated tumors at the end of the treatment period. In contrast, an almost complete tumor regression was observed in all mice treated with the two agents in combination. This determined a prolonged life span of the mice that was significantly different as compared with controls ($P < 0.001$), to MAb C225-treated group ($P < 0.001$), or to the topotecan-treated group ($P < 0.001$). All mice of the topotecan plus MAb C225 group were the only animals alive 14 weeks after tumor cell injection. Furthermore, 20% of mice in this group were still alive after 19 weeks. The combined treatment with MAb C225 and topotecan was well tolerated by mice with no signs of acute or delayed toxicity. These results provide a rationale for the evaluation of the anticancer activity of the combination of topoisomerase I inhibitors and anti-EGFR blocking MAbs in clinical trials.

INTRODUCTION

Medical treatment of cancer is presently based on cytotoxic drugs that determine cell killing by acting on cellular targets that are generally common to both cancer and normal cells. Therapeutic regimens generally consist of a combination of two or more non-cross-resistant drugs. More recently, the increase in dose of various cytotoxic drugs to supertoxic levels obtained with the use of hematopoietic cell rescue has been tested and shown effective in some malignancies. However, several human cancer types are relatively insensitive or become resistant to the treatment with cytotoxic drugs.

The increasing understanding of the molecular mechanisms of neoplastic transformation and progression has prompted the search for biological agents interfering with the intracellular targets involved in this process. Several of these agents have recently entered clinical evaluation in cancer patients. In this respect, peptide growth factors are involved in regulating normal and cancer epithelial cell proliferation and differentiation (1, 2). EGF³-related growth factors, such as TGF- α , are potent

Received 11/11/98; revised 1/19/99; accepted 1/20/99.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This study was supported by a grant from the Associazione Italiana per la Ricerca sul Cancro and from the Consiglio Nazionale delle Ricerche Target Project on Biotechnologies.

² To whom requests for reprints should be addressed, at Cattedra di Oncologia Medica, Dipartimento di Endocrinologia e Oncologia Molecolare e Clinica, Università degli Studi di Napoli Federico II, Via S. Pansini, 5-80131 Naples, Italy. Phone: 39-081-7462061; Fax: 39-081-7462066.

³ The abbreviations used are: EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; TGF- α , transforming growth factor α ; MAb, monoclonal antibody.

mitogens for several human epithelial cell types including breast, colon, ovary, kidney, prostate, and lung and have been implicated in human cancer development and progression through autocrine and paracrine pathways (3). TGF- α binds to the extracellular domain of the EGFR and activates its intracellular tyrosine kinase domain (3). Enhanced expression of TGF- α and/or EGFR has been detected in the majority of human carcinomas (3) and has been associated with a poor prognosis in several human tumor types, such as breast cancer (4). For these reasons, the blockade of the TGF- α -EGFR autocrine pathway by using anti-EGFR blocking MAbs, recombinant proteins containing TGF- α or EGF fused to toxins, or EGFR-specific tyrosine-kinase inhibitors has been proposed as a potential therapeutic modality (5–12). Several blocking anti-EGFR MAbs that inhibit the *in vitro* and *in vivo* growth of human cancer cell lines that express TGF- α and EGFR have been generated (13–16). Among these, MAb 528 and MAb 225 are two mouse MAbs that have been extensively characterized for their biological and pharmacological properties and represent the first anti-EGFR blocking agents that have entered clinical evaluation in cancer patients (5, 6). MAb 528 and MAb 225 bind to the EGFR with affinity similar to TGF- α , compete with ligand for receptor binding, and block ligand-induced activation of EGFR tyrosine kinase (13, 14). To avoid human anti-mouse antibody production that could interfere with the therapeutic efficacy of repeated administrations of mouse MAbs, a chimeric human-mouse MAb 225 (MAb C225), which contains the human IgG1 constant region, has recently been developed and purified for clinical use and is in Phase II trials in patients with advanced cancer (17, 18).

In recent years, there has been a growing interest in combining conventional chemotherapeutic agents with biological agents that selectively inhibit key intracellular targets involved in the process of neoplastic transformation. In this respect, it has been shown that the combined treatment of mice bearing well-established human tumor xenografts with blocking anti-EGFR MAbs and with some cytotoxic drugs, such as doxorubicin and cisplatin, significantly increases the antitumor activity of these drugs (6, 19, 20). In the present study, we have investigated whether MAb C225 has any cooperative effect with topotecan in a series of human cancer cell lines *in vitro* and in nude mice bearing human GEO colon cancer xenografts. We have selected the cytotoxic agent topotecan, a novel promising chemotherapeutic agent that selectively inhibits the nuclear enzyme topoisomerase I, because it has shown antitumor activity in a variety of human malignancies, including ovarian cancer, small cell lung cancer, breast cancer, and colorectal cancer, in which EGFR is generally overexpressed (21).

MATERIALS AND METHODS

Materials. MAb C225 is a human-mouse chimeric anti-EGFR IgG1 class MAb, the biochemical and biological characteristics of which have been described previously (17). MAb C225 was kindly provided by Dr. H. Waksal (ImClone Systems, New York, NY). The topoisomerase I inhibitor topotecan was a generous gift of SmithKline Beecham Italia (Milan, Italy).

Cell Lines. GEO human colon cancer, OVCAR-3 human ovarian cancer, and ZR-75-1 human breast cancer cell lines

were obtained from the American Type Culture Collection (Rockville, MD). 1A9 and their paclitaxel-resistant derivative PTX-10 and PTX-22 human ovarian cancer cell lines (22) were kindly provided by Dr. A. Fojo (National Cancer Institute, Bethesda, MD). The p53 status of the cancer cell lines is the following: wild-type gene, GEO,⁴ ZR-75-1 (23); point mutated gene, OVCAR-3, G to A in codon 248 (23); and unknown, 1A9, PTX-10, PTX-22. All cell lines were maintained in DMEM supplemented with 10% heat-inactivated fetal bovine serum, 20 mM HEPES (pH 7.4), 100 IU/ml penicillin, 100 μ g/ml streptomycin, and 4 mM glutamine (ICN, Irvine, UK) in a humidified atmosphere of 95% air and 5% CO₂ at 37°C.

Growth in Soft Agar. Cells (10⁴ cells/well) were suspended in 0.5 ml of 0.3% Difco Noble agar (Difco, Detroit, MI) supplemented with complete culture medium. This suspension was layered over 0.5 ml of 0.8% agar-medium base layer in 24-multiwell cluster dishes (Becton Dickinson, Lincoln Park, NJ) and treated with different concentrations of MAb C225 and/or topotecan. After 10–14 days, the cells were stained with nitro blue tetrazolium (Sigma Chemical Co., St. Louis, MO), and colonies larger than 0.05 mm were counted as described previously (24).

Apoptosis. To evaluate the induction of apoptosis, 10⁵ cells were plated in complete medium in 60-mm tissue culture dishes (Becton Dickinson) and treated every day for a total of 3 days with different concentrations of MAb C225 and/or topotecan. For the combined treatment, cells were treated every day for 3 days with the indicated concentration of topotecan and then every day for 3 days with the indicated concentration of MAb C225. After 7 days, both adherent and detached cells were harvested. Flow cytometric analysis of apoptotic cell death was performed on cell pellet fixed in 70% ethanol, washed in PBS, and mixed with RNase (Sigma) and propidium iodide (Sigma) solution as reported previously (25). DNA content was analyzed by a FACScan flow-cytometer (Becton Dickinson, San Jose, CA) coupled with a Hewlett Packard computer, and the percentage of apoptotic cells was calculated by gating the hypodiploid region on the DNA content histogram using the LYSYS software (Becton Dickinson) as reported previously (25).

GEO Xenografts in Nude Mice. Five- to 6-week-old female Balb/cAnNCrIBR athymic (*nu+/nu+*) mice were purchased from Charles River Laboratories (Milan, Italy). The research protocol was approved, and mice were maintained in accordance with institutional guidelines of the University of Naples Animal Care and Use Committee. Mice were acclimated to the University of Naples Medical School Animal Facility for one week before being injected with cancer cells. Mice were injected s.c. with 10⁷ GEO cells that had been resuspended in 200 μ l of Matrigel (Collaborative Biomedical Products, Bedford, MA). After 7 days, when well-established tumors of ~0.2 cm³ in diameter were detected, 10 mice/group were treated i.p. with topotecan alone (2 mg/kg/dose, twice weekly on days 1 and 2 for 2 weeks), with MAb C225 alone (0.25 mg/dose, twice weekly on days 3 and 6 for 5 weeks), or with both drugs. Tumor size was measured using the formula $\pi/6 \times \text{larger diameter} \times (\text{smaller diameter})^2$, as reported previously (26). The exper-

⁴ Unpublished observations.




Fig. 1 Dose-dependent growth inhibitory effects of MAb C225 (A) or of topotecan (B) on the soft agar growth of human cancer cell lines. Data represent the average of three different experiments, each performed in triplicate; bars, SD.

iment was repeated three times with a total of 30 mice for each group.

Statistical Analysis. The Student's *t* test (27) and the Mantel-Cox log-rank test (28) were used to evaluate the statistical significance of the results. All *P*s represent two-sided tests of statistical significance. All analyses were performed with the BMDP New System statistical package version 1.0 for Microsoft Windows (BMDP Statistical Software, Los Angeles, CA).

RESULTS

As shown in Fig. 1, we first evaluated the effect of topotecan or MAb C225 treatment on the soft agar growth of various human epithelial cancer cell lines. For this purpose, we selected GEO colon cancer, ZR-75-1 breast cancer, and OVCAR-3 and 1A9 ovarian cancer cell lines. We also tested two paclitaxel-resistant derivatives of 1A9 cells that possess specific point mutations in the β -tubulin gene (22). All these human cancer

cell lines have functional EGFR, ranging from \sim 20,000 (ZR-75-1) to 40,000 (GEO), 100,000 (1A9, PTX-10, and PTX-22), 150,000 (OVCAR-3) EGF binding sites/cell (24, 29).⁴ Treatment with the anti-EGFR blocking MAb C225 determined a dose-dependent inhibition of colony formation in soft agar at doses ranging between 0.25 and 10 μ g/ml in all cancer cell lines tested (Fig. 1A). Similarly, a dose-dependent, growth-inhibitory effect was observed when the different cancer cell lines were treated with the topoisomerase I-selective drug topotecan at doses between 1 and 100 nM (Fig. 1B). To determine whether the combined treatment with topotecan and MAb C225 could enhance the antiproliferative effect of single-agent treatment, the six cancer cell lines were treated with different combinations of the two compounds in a sequential schedule of treatment with topotecan followed by MAb C225. As shown in Figs. 2 and 3, a supraadditive growth-inhibitory effect was observed at all doses of MAb C225 and topotecan tested. Furthermore, when lower doses of these agents were used, the antiproliferative effect was clearly cooperative in all cell lines examined. For example, in GEO cells, treatment with 5 nM topotecan or with 0.25 μ g/ml MAb C225 determined \sim 10% growth inhibition, whereas the sequential treatment caused a 50% inhibition of colony formation in soft agar (Fig. 2). The cooperativity quotient of this treatment, defined as the ratio between the actual growth inhibition obtained with topotecan followed by MAb C225 and the sum of the growth inhibition achieved by each agent, was approximately 2.5.

The supraadditive antiproliferative effects of the combination of topotecan and MAb C225 described above could be dependent on the sequence of treatment. We, therefore, tested an inverted treatment schedule in which cancer cells were treated first with MAb C225, then with the cytotoxic drug topotecan. In ZR-75-1, OVCAR-3, and GEO cancer cells pretreatment with either 0.25 or 0.5 μ g/ml MAb C225 reduced the growth-inhibitory effect of topotecan at all doses tested (data not shown). Similar interference in growth inhibition was observed when treatment with topotecan and MAb C225 was concomitant (data not shown). These results suggest that treatment with the anti-EGFR blocking MAb should follow the cytotoxic drug in the sequence of treatment.

We next determined whether the cooperative growth-inhibitory effect of topotecan and MAb C225 could involve induction of programmed cell death in cancer cells. GEO and OVCAR-3 cells were treated with different concentrations of topotecan or MAb C225 for 3 days or with 5 nM topotecan for 3 days followed by 0.5 μ g/ml MAb C225 for 3 days. The percentage of apoptotic cells was determined on day 7 from the beginning of treatment (Fig. 4). MAb C225 treatment alone did not induce apoptosis in either GEO or OVCAR-3 cells. Topotecan treatment increased in a dose-dependent manner apoptotic cell death both in GEO and OVCAR-3 cells. The sequential treatment with the two agents determined a marked induction of apoptosis in both cell lines. In fact, 5 nM topotecan, followed by 0.5 μ g/ml MAb C225, induced apoptosis in 45% of GEO cells compared with untreated cells (6%) or to 5 nM topotecan-treated cells (22%; Fig. 4A). A similar potentiation of apoptosis was observed in OVCAR-3 cells (Fig. 4B). These results suggest that the cooperative antiproliferative activity of topotecan and MAb C225 is accompanied by induction of programmed cell death.




Fig. 2 Growth-inhibitory effects of the sequential treatment with topotecan and MAb C225 on the soft agar growth of GEO, ZR-75-1, and OVCAR-3 cancer cells. GEO: topotecan, 1 nM (a, e, and i), 5 nM (b, f, and j), 10 nM (c, g, and k), 25 nM (d, h, and l). MAb C225: 0.25 µg/ml (a–d), 0.5 µg/ml (e–h), 1 µg/ml (i–l). ZR-75-1: topotecan, 1 nM (a, e, and i), 5 nM (b, f, and j), 10 nM (c, g, and k), 25 nM (d, h, and l). MAb C225: 0.25 µg/ml (a–d), 0.5 µg/ml (e–h), 1 µg/ml (i–l). OVCAR-3: topotecan, 1 nM (a, e, and i), 5 nM (b, f, and j), 10 nM (c, g, and k), 25 nM (d, h, and l). MAb C225: 0.25 µg/ml (a–d), 0.5 µg/ml (e–h), 1 µg/ml (i–l). Cells were plated in soft agar as described in “Materials and Methods” and treated sequentially with topotecan on days 1 and 2 and with MAb C225 on days 3 and 5 or with each agent alone. Colonies were counted after 10–14 days. Data are expressed as the percentage of growth inhibition as compared with the growth of untreated control cells. For each pair of columns, the height of the columns on the left represents the sum of the effect of each agent and, therefore, the expected growth inhibition if their effect is additive when used in combination. The total height of the columns on the right indicates the observed growth inhibition when the agents are used in combination. The difference between the heights of the paired columns reflects the magnitude of cooperativity on growth inhibition. Data represent the average of three different experiments, each performed in duplicate; bars, SD.

We next evaluated whether the cooperative effect observed *in vitro* could be obtained also *in vivo*. We have previously studied the effect of different novel therapeutic agents including MAb C225 in nude mice bearing GEO colon cancer xenografts (26). GEO cells form moderately differentiated adenocarcino-




Fig. 3 Growth-inhibitory effects of the sequential treatment with topotecan and MAb C225 on the soft agar growth of 1A9, PTX-10, and PTX-22 cancer cells. Topotecan: 1 nM (a and e), 5 nM (b and f), 10 nM (c and g), 25 nM (d and h). MAb C225: 0.5 µg/ml (a–d), 1 µg/ml (e–h). Cells were plated in soft agar as described in “Materials and Methods” and treated sequentially with topotecan on days 1 and 2 and with MAb C225 on days 3 and 5 or with each agent alone. Colonies were counted after 10–14 days. Data are expressed as percentage of growth inhibition as compared with the growth of untreated control cells. For each pair of columns, the height of the columns on the left represents the sum of the effect of each agent and, therefore, the expected growth inhibition if their effect is additive when used in combination. The total height of the columns on the right indicates the observed growth inhibition when the agents are used in combination. The difference between the heights of the paired columns reflects the magnitude of cooperativity on growth inhibition. Data represent the average of three different experiments, each performed in duplicate; bars, SD.

mas that express the EGFR and various specific ligands when injected s.c. in immunodeficient mice. Therefore, we used this model to investigate the antitumor activity of MAb C225 and topotecan combination. GEO cells were injected s.c. into the dorsal flank of nude mice. When established GEO tumors of ~0.2 cm³ were detectable, mice were treated i.p. with topotecan, 2 mg/kg/dose, on days 1 and 2 of each week for a total of 2 weeks or with 0.25 mg/dose MAb C225, twice weekly on days 3 and 6 of each week for a total of 5 weeks, or with both compounds. Tumor size was measured weekly, and mice sur-




Fig. 4 Induction of apoptosis by treatment with MAb C225 and/or topotecan in GEO (A) or OVCAR-3 (B) cancer cells. Cells were treated with: topotecan, 1 (column 1), 5 (column 2), 10 (column 3), and 25 (column 4) nM each day for 3 days; MAb C225, 0.25 (column 5), 0.5 (column 6), 1 (column 7) $\mu\text{g/ml}$ each day for 3 days; topotecan, 5 nM each day for 3 days followed by 0.5 $\mu\text{g/ml}$ MAb C225, each day for 3 days (column 8). C, untreated control cells. Flow cytometric analysis of DNA fragmentation was performed 7 days after the beginning of treatment. Data represent the average of triplicate determinations; bars, SD.

vival was monitored. As illustrated in Fig. 5, mice treatment with each agent inhibited tumor growth *in vivo* as compared with control untreated mice. However, shortly after the end of the 2 weeks of treatment with topotecan or of the 5 weeks of treatment with MAb C225, GEO tumors resumed the growth rate of untreated tumors. When the two agents were used in combination, an almost complete regression of tumors in all mice was observed. Furthermore, GEO tumor growth was significantly delayed in mice that received MAb C225 plus topotecan treatment. In fact, tumors grew very slowly for ~50–70 days following the end of treatment when they resumed a growth rate similar to controls (Fig. 5).

As shown in Fig. 6, GEO tumors reached a size not compatible with normal life in all untreated mice within 4–5 weeks. A small but significant increase in mice survival was observed in the group treated with topotecan alone ($P < 0.02$). MAb C225 treatment determined a longer survival as compared with the control group ($P < 0.001$) and to the topotecan treated-




Fig. 5 Antitumor activity of topotecan and MAb C225 on established GEO human colon carcinoma xenografts. Mice were injected s.c. in the dorsal flank with 10^7 GEO cells as described in “Materials and Methods.” After 7 days (average tumor size, 0.2 cm^3), mice were treated i.p. with topotecan alone (2 mg/kg/dose, twice weekly on days 1 and 2 of each week for 2 weeks), or with MAb C225 alone (0.25 mg/dose, twice weekly on days 3 and 6 of each week for 5 weeks), or with both drugs with the same sequential schedule. Each group consisted of 10 mice. The experiment was repeated for three times. Data represent the average on a total of 30 mice for each group; bars, SD. Student’s *t* test was used to compare tumor sizes among different treatment groups at day 29 after GEO cell injection. MAb C225 versus control ($P < 0.001$); topotecan versus control ($P < 0.001$); topotecan followed by MAb C225 versus MAb C225 ($P < 0.001$); topotecan followed by MAb C225 versus topotecan ($P < 0.001$).

group ($P < 0.001$). The delayed GEO tumor growth in the MAb C225 plus topotecan-treated group was accompanied by a prolonged life span of the mice that was significantly different as compared with controls ($P < 0.001$), to MAb C225-treated group ($P < 0.001$), or to topotecan-treated group ($P < 0.001$). In fact, all mice of this group were the only animals alive 14 weeks after tumor cell injection (Fig. 6). Furthermore, 20% of mice in this group were still alive after 19 weeks. The combined treatment with MAb C225 and topotecan was well tolerated by mice, because no weight loss or other signs of acute or delayed toxicity were observed.

DISCUSSION

The possibility of combining conventional cytotoxic drugs with novel agents that specifically interfere with key pathways controlling cancer cell survival, proliferation, invasion, and/or metastatic spreading has generated a wide interest in the past few years. In fact, this could be a promising therapeutic approach for several reasons: (a) because the cellular targets for these agents and their mechanism(s) of action are different from those of cytotoxic drugs, it is possible their combination with chemotherapy without cross-resistance; and (b) alterations in the expression and/or the activity of genes that regulate mitogenic signals not only can directly cause perturbation of cell growth but also may affect the sensitivity of cancer cells to conventional chemotherapy and radiation therapy (30). EGFR overexpression has been generally found in human cancer cell lines that are




Fig. 6 Effects of topotecan and/or MAb C225 treatment on the survival of GEO tumor-bearing mice. The experimental design is described in the legend to Fig. 5. ●, Control; ○, topotecan; ▼, MAb C225; ▽, topotecan followed by MAb C225. Thirty mice/group were monitored for survival. Differences in animal survival among groups were evaluated using the Mantel-Cox log-rank test. Mice survival was significantly different between the MAb C225-treated group and the control group ($P < 0.001$); the MAb C225-treated group and the topotecan-treated group ($P < 0.001$); the topotecan-treated group and the control group ($P < 0.02$); the topotecan plus MAb C225-treated group and the control group ($P < 0.001$); the topotecan plus MAb C225-treated group and the MAb C225-treated group, ($P < 0.001$); and the topotecan plus MAb C225-treated group and the topotecan-treated group, ($P < 0.001$).

resistant to different cytotoxic drugs (31–33). In this respect, it has been postulated that EGFR overexpression would serve as a cell survival response to counteract apoptotic signaling in cancer cells exposed to a cytotoxic drug (33).

The results of the present study demonstrate that treatment with the anti-EGFR MAb C225 increases the topoisomerase I inhibitor topotecan-induced antiproliferative effects in several human epithelial cancer cell lines that express functional EGFR. In fact, blockade of EGFR after treatment with topotecan determines growth inhibition *in vitro* and *in vivo* that is not simply additive. In established GEO human colon carcinoma xenografts, treatment with each agent transiently inhibits tumor growth because tumors resume the growth rate of untreated control after cessation of therapy. In contrast, the sequential treatment with topotecan and MAb C225 determines tumor regression in all mice. In this group of mice, tumors grow very slowly for approximately 8–9 weeks following the end of treatment when they finally resume a growth rate similar to controls. This effect is accompanied by a significant benefit in animal survival in the group treated with both agents as compared with the groups treated with a single agent. In fact, 100% of mice treated with topotecan and MAb C225 are still alive 14 weeks after cancer cell injection, whereas all mice died within 6 or 10 weeks in the groups treated with topotecan or with MAb C225, respectively. This study also shows that the combined treatment with topotecan and MAb C225 is well tolerated by mice without signs of toxicity, suggesting that additional cycles of topotecan followed by MAb C225 could be given in an attempt to increase mice survival and possibly to obtain eradication of established tumors.

The growth-inhibitory effect of the combined treatment with topotecan and MAb C225 is schedule dependent because a clear cooperative antiproliferative activity is observed only when cancer cells are first treated with topotecan and then with MAb C225. This can be due to the effects on cell cycle determined by treatment with each agent. Topotecan is effective in killing cells mostly during the S and G₂ phases of the cell cycle (21), whereas the anti-EGFR blocking antibody MAb C225 determines a cytostatic growth-inhibitory effect with G₁ arrest (34, 35). Therefore, pretreatment with MAb C225 could render cells less responsive to topotecan. On the other hand, a possible mechanism by which MAb C225 exerts a supraadditive antiproliferative effect when administered following topotecan could be by increasing programmed cell death. We have observed a significant potentiation of apoptosis in both GEO and OVCAR-3 cells treated with topotecan and MAb C225 in a sequential schedule. In this context, the blockade of the EGFR-mediated signaling could render cancer cells unable to overcome the genotoxic damage induced by topotecan inhibition of topoisomerase I and, therefore, could render them more susceptible to enter programmed cell death.

The results of the present study extend those of previous reports that have shown a cooperative antitumor activity of some cytotoxic drugs such as cisplatin, doxorubicin, or paclitaxel with various blocking antibodies generated against the EGFR or the closely related type I tyrosine kinase receptor *c-erbB-2* (15, 19, 20, 36–40). Therefore, treatment with agents that are able to selectively inhibit biochemical pathways controlling mitogenic signaling, DNA integrity, and/or cell cycle progression may cooperate with cytotoxic drugs by rendering cancer cells more susceptible to irreparable damage. In fact, an increase in apoptosis and growth inhibition induced by several drugs such as topoisomerase II inhibitors, taxanes, and platinum derivatives has been demonstrated in a large series of human cancer cell lines for agents that selectively inhibit type I protein kinase A, an intracellular serine-threonine kinase that acts down-stream to various mitogenic signals including the activated EGFR (41–43). Similarly, inhibition of *c-raf* expression by an antisense oligonucleotide enhances the antitumor activity of cisplatin, doxorubicin, or mitomycin C in human cancer tumor xenografts in nude mice (44). Taken together, these studies support the hypothesis that it is possible to enhance antitumor activity by treatment with maximum tolerated doses of cytotoxic drugs and specific inhibitors of signal transduction pathways instead of increasing chemotherapy doses to super-toxic levels that require complex medical support, including hematopoietic rescue (30). Because Phase I studies have shown that MAb C225 can be given to cancer patients with weekly repeated *i.v.* injections at doses that are well tolerated and that saturate the EGFR (45), our study provides a rationale for translating the combination of topotecan and MAb C225 into clinical trials. In this respect, preliminary data have shown the feasibility of a treatment with cisplatin and MAb C225 in patients with advanced head and neck or lung carcinomas (18). Similarly, a Phase II study has shown recently that recombinant humanized anti-*c-erbB-2* MAb (Herceptin) could increase the antitumor activity of cisplatin in metastatic breast cancer patients overexpressing *c-erbB-2* (46). Finally, a preliminary report of a randomized Phase III trial has demonstrated recently

that addition of Herceptin to first-line chemotherapy of advanced breast cancer patients with paclitaxel or doxorubicin increases the anticancer activity of chemotherapy alone (47).

REFERENCES

- Sporn, M. B., and Roberts, A. B. Autocrine secretion: ten years later. *Ann. Intern. Med.*, 117: 408–414, 1992.
- Aaronson, S. A. Growth factors and cancer. *Science (Washington DC)*, 254: 1146–1153, 1991.
- Salomon, D. S., Brandt, R., Ciardiello, F., and Normanno, N. Epidermal growth factor-related peptides and their receptors in human malignancies. *Crit. Rev. Oncol. Hematol.*, 19: 183–232, 1995.
- Fox, S. B., Smith, K., Hollyer, J., Greenall, M., Hastrich, D., and Harris, A. L. The epidermal growth factor receptor as a prognostic marker: results of 370 patients and review of 3009 patients. *Breast Cancer Res. Treat.*, 29: 41–49, 1994.
- Fan, Z., and Mendelsohn, J. Therapeutic application of anti-growth factor receptor antibodies. *Curr. Opin. Oncol.*, 10: 67–73, 1998.
- Baselga, J., and Mendelsohn, J. Type I receptor tyrosine kinases as targets for therapy in breast cancer. *J. Mammary Gland Biol. Neopl.*, 2: 165–174, 1997.
- Masui, H., Kamrath, H., Apell, G., Houston, L. L., and Mendelsohn, J. Cytotoxicity against human tumor cells mediated by the conjugate of anti-epidermal growth factor receptor monoclonal antibody to recombinant ricin A chain. *Cancer Res.*, 49: 3482–3488, 1989.
- Arteaga, C. L., Hurd, S. D., Dugger, T. C., Winnier, A. R., and Robertson, J. B. Epidermal growth factor receptors in human breast carcinoma cells: a potential selective target for transforming growth factor α -*Pseudomonas* exotoxin 40 fusion protein. *Cancer Res.*, 54: 4703–4709, 1994.
- Kirk, J., Carmichael, J., Stratford, I. J., and Harris, A. L. Selective toxicity of TGF- α -PE40 to EGFR-positive cell lines: selective protection of low EGFR-expressing cell lines by EGF. *Br. J. Cancer*, 69: 988–994, 1994.
- Lorimer, I. A. J., Wikstrand, C. J., Batra, S. K., Bigner, D. D., and Pastan, I. Immunotoxins that target an oncogenic mutant epidermal growth factor receptor expressed in human tumors. *Clin. Cancer Res.*, 1: 859–864, 1995.
- Schmidt, M., and Wels, W. Targeted inhibition of tumor cell growth by a bispecific single-chain toxin containing an antibody domain and TGF α . *Br. J. Cancer*, 74: 853–862, 1996.
- Buchdunger, E., Mett, H., Trinks, U., Regenass, U., Muller, M., Meyer, T., Beilstein, P., Wirz, B., Schneider, P., Traxler, P., and Lydon, N. 4,5-Bis(4-fluoroanilino)phthalimide: a selective inhibitor of the epidermal growth factor receptor signal transduction pathway with potent *in vivo* antitumor activity. *Clin. Cancer Res.*, 1: 813–821, 1995.
- Gill, G. N., Kawamoto, T., Cochet, C., Lee, A., Sato, J. D., Masui, H., MacLeod, C., and Mendelsohn, J. Monoclonal anti-epidermal growth factor receptor antibodies which are inhibitors of epidermal growth factor binding and antagonists of epidermal growth factor-stimulated tyrosine protein kinase activity. *J. Biol. Chem.*, 259: 7755–7760, 1984.
- Masui, H., Kawamoto, T., Sato, J. D., Wolf, B., Sato, G., and Mendelsohn, J. Growth inhibition of human tumor cells in athymic mice by anti-epidermal growth factor receptor monoclonal antibodies. *Cancer Res.*, 44: 1002–1007, 1984.
- Aboud-Pirak, E., Hurwitz, E., Pirak, M. E., Bellot, F., Schlessinger, J., and Sela, M. Efficacy of antibodies to epidermal growth factor receptor against KB carcinoma *in vitro* and in nude mice. *J. Natl. Cancer Inst.*, 80: 1605–1611, 1988.
- Modjtahedi, H., Eccles, S., Box, G., Styles, J., and Dean, C. Immunotherapy of human tumour xenografts overexpressing the EGF receptor with rat antibodies that block growth factor-receptor interaction. *Br. J. Cancer*, 67: 254–261, 1993.
- Goldstein, N. I., Prewett, M., Zuklys K., Rockwell, P., and Mendelsohn, J. Biological efficacy of a chimeric antibody to the epidermal growth factor receptor in a human tumor xenograft model. *Clin. Cancer Res.*, 1: 1311–1318, 1995.
- Falcey, J., Pfister, D., Cohen, R., Cooper, M., Bowden, C., Burtness, B., Mendelsohn, J., and Waksal, H. A study of anti-epidermal growth factor receptor (EGFr) monoclonal antibody C225 and cisplatin in patients (pts) with head and neck or lung carcinomas. *Proc. Am. Soc. Clin. Oncol.*, 16: 1364, 1997.
- Fan, Z., Baselga, J., Masui, H., and Mendelsohn, J. Antitumor effect of anti-epidermal growth factor receptor monoclonal antibodies plus *cis*-diamminedichloroplatinum on well established A431 cell xenografts. *Cancer Res.*, 53: 4637–4642, 1992.
- Baselga, J., Norton, L., Masui, H., Pandiella, A., Coplan, K., Miller, W. H., and Mendelsohn, J. Antitumor effects of doxorubicin in combination with anti-epidermal growth factor receptor monoclonal antibodies. *J. Natl. Cancer Inst.*, 85: 1327–1333, 1993.
- Carmichael, J., and Ozols, R. F. Topotecan, an active new antineoplastic agent: review and current status. *Exp. Opin. Invest. Drugs*, 6: 593–608, 1997.
- Giannakakou, P., Sackett, D. L., Kang, Y.-K., Zhan, Z., Buters, J. T. M., Fojo, T., and Poruchynsky, M. S. Paclitaxel-resistant human ovarian cancer cells have mutant β -tubulins that exhibit impaired paclitaxel-driven polymerization. *J. Biol. Chem.*, 272: 17118–17125, 1997.
- O'Connor, P. M., Jackman, J., Bae, I., Myers, T. G., Fan, S., Mutoh, M., Scudiero, D. A., Monks, A., Sausville, E. A., Weinstein, J. N., Friend, S., Fornace, A. J., and Kohn, K. W. Characterization of the p53 tumor suppressor pathway in cell lines of the National Cancer Institute anticancer drug screen and correlations with the growth-inhibitory potency of 123 anticancer agents. *Cancer Res.*, 57: 4285–4300, 1997.
- Ciardiello, F., Damiano, V., Bianco, C., di Isernia, G., Ruggiero, A., Caraglia, M., Tagliaferri, P., Baselga, J., Mendelsohn, J., Bianco, A. R., and Tortora, G. Cooperative antiproliferative effects of 8-Cl-cAMP and 528 anti-epidermal growth factor receptor monoclonal antibody on human cancer cells. *Clin. Cancer Res.*, 1: 161–167, 1995.
- Nicoletti, I., Migliorati, G., Pagliacci, M. C., Grignani, F., and Riccardi, C. A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J. Immunol. Methods*, 139: 271–279, 1991.
- Ciardiello, F., Damiano, V., Bianco, R., Bianco, C., Fontanini, G., De Laurentis, M., De Placido, S., Mendelsohn, J., Bianco, A. R., and Tortora, G. Antitumor activity of combined blockade of epidermal growth factor receptor and protein kinase A. *J. Natl. Cancer Inst.*, 88: 1770–1776, 1996.
- Fleiss, J. L. *Statistical Methods for Rates and Proportions*, Ed. 2. New York: Wiley, 1981.
- Mantel, N. Evaluation of survival data and two new rank order statistics arising in its consideration. *Cancer Chem. Rep.*, 50: 163–170, 1966.
- Wosikowski, K., Schuurhuis, D., Johnson, K., Paull, K. D., Myers, T. G., Weinstein, J. N., and Bates, S. E. Identification of epidermal growth factor receptor and c-erbB2 pathway inhibitors by correlation with gene expression patterns. *J. Natl. Cancer Inst.*, 89: 1505–1515, 1997.
- Mendelsohn, J., and Fan, Z. Epidermal growth factor receptor family and chemosensitization. *J. Natl. Cancer Inst.*, 89: 341–343, 1997.
- Vickers, P. J., Dickson, R. B., Shoemaker, R., and Cowan, K. H. A multidrug resistant MCF-7 human breast cancer cell line which exhibits cross-resistance to antiestrogens and hormone-independent tumor growth *in vivo*. *Mol. Endocrinol.*, 2: 886–892, 1988.
- Meyers, M. B., Shen, W. P. V., Spengler, B. A., Ciccarone, V., O'Brien, J. P., Donner, D. B., Furth, M. E., and Biedler, J. L. Increased epidermal growth factor receptor in multidrug-resistant human neuroblastoma cells. *J. Cell. Biochem.*, 38: 87–97, 1988.
- Wosikowski, K., Schuurhuis, D., Kops, G. J. P. L., Saceda, M., and Bates, S. E. Altered gene expression in drug-resistant human breast cancer cells. *Clin. Cancer Res.*, 2: 2405–2414, 1997.
- Peng, D., Fan, Z., Lu, Y., DeBlasio, T., Scher, H., and Mendelsohn, J. Anti-epidermal growth factor receptor monoclonal antibody 225 up-

- regulates p27^{KIP1} and induces G₁ arrest in prostatic cancer cell line DU145. *Cancer Res.*, 56: 3666–3669, 1996.
35. Wu, X., Rubin, M., Fan, Z., DeBlasio, T., Soos, T., Koff, A., and Mendelsohn, J. Involvement of p27^{KIP1} in G₁ arrest mediated by anti-epidermal growth factor receptor monoclonal antibody. *Oncogene*, 12: 1397–1403, 1996.
36. Hancock, M. C., Langton, B. C., Chan, T., Toy, P., Monahan, J. J., Mischak, R. P., and Shaver, L. K. A monoclonal antibody against the c-erbB-2 protein enhances the cytotoxicity of *cis*-diamminedichloroplatinum against human breast and ovarian tumor cell lines. *Cancer Res.*, 51: 4574–4580, 1991.
37. Arteaga, C. L., Winnier, A. R., Poirer, M. C., Lopez-Larraz, D. M., Shaver, L. K., Hurd, S. D., and Stewart, S. J. p185c-erbB-2 signaling enhances cisplatin-induced cytotoxicity in human breast carcinoma cells: association between an oncogenic receptor tyrosine kinase and drug-induced DNA repair. *Cancer Res.*, 54: 3758–2765, 1994.
38. Pietras, R. J., Fendly, B. M., Chazin, V. R., Pegram, M. D., Howell, S. B., and Slamon, D. J. Antibody to HER-2/neu receptor blocks DNA repair after cisplatin in human breast and ovarian cancer cells. *Oncogene*, 9: 1829–1838, 1994.
39. Baselga, J., Norton, L., Albanell, J., Kim, Y.-M., and Mendelsohn, J. Recombinant humanized anti-HER2 antibody (Herceptin) enhances the antitumor activity of paclitaxel and doxorubicin against HER2/*neu* overexpressing human breast cancer xenografts. *Cancer Res.*, 58: 2825–2831, 1998.
40. Pietras, R. J., Pegram, M. D., Finn, R. S., Maneval, D. A., and Slamon, D. J. Remission of human breast cancer xenografts on therapy with humanized monoclonal antibody to HER-2 receptor and DNA-reactive drugs. *Oncogene*, 17: 2235–2249, 1998.
41. Tortora, G., di Isernia, G., Sandomenico, C., Bianco, R., Pomatico, G., Pepe, S., Bianco, A. R., and Ciardiello, F. Synergistic inhibition of growth and induction of apoptosis by 8-Chloro-cAMP and paclitaxel or cisplatin in human cancer cells. *Cancer Res.*, 57: 5107–5111, 1997.
42. Tortora, G., Caputo, R., Damiano, V., Bianco, R., Pepe, S., Bianco, A. R., Jiang, Z., Agrawal, S., and Ciardiello, F. Synergistic inhibition of human cancer cell growth by cytotoxic drugs and mixed backbone antisense oligonucleotide targeting protein kinase A. *Proc. Natl. Acad. Sci. USA*, 94: 12586–12591, 1997.
43. Ciardiello, F., and Tortora, G. Interactions between the epidermal growth factor receptor and type I protein kinase A: biological significance and therapeutic implications. *Clin. Cancer Res.*, 4: 821–828, 1998.
44. Geiger, T., Muller, M., Monia, B. P., and Fabbro, D. Antitumor activity of a C-raf antisense oligonucleotide in combination with standard chemotherapeutic agents against various human tumors transplanted subcutaneously into nude mice. *Clin. Cancer Res.*, 3: 1179–1185, 1997.
45. Anderson, V., Cooper, M., O'Chei, E., Gilly, J., Falcey, J., and Waksal, H. W. Dose dependent pharmacokinetics of a chimerized monoclonal antibody (C225) against EGFR. *Proc. Am. Assoc. Cancer Res.*, 39: 3561, 1998.
46. Pegram, M. D., Lipton, A., Hayes, D. F., Weber, B. L., Baselga, J., Tripathy, D., Baly, D., Baughman, S. A., Twaddell, T., Glaspy, J. A., and Slamon, D. J. Phase II study of receptor-enhanced chemosensitivity using recombinant humanized anti-p185^{HER2/neu} monoclonal antibody plus cisplatin in patients with HER2/*neu*-overexpressing metastatic breast cancer refractory to chemotherapy treatment. *J. Clin. Oncol.*, 16: 2659–2671, 1998.
47. Slamon, D. J., Leyland-Jones, B., Shak, S., Paton, V., Bajamonde, A., Fleming, T., Eiermann, W., Volter, J., Baselga, J., and Norton, L. Addition of Herceptin (humanized anti-HER2 antibody) to first line chemotherapy for HER2 overexpressing metastatic breast cancer markedly increases anticancer activity: a randomized multinational controlled phase III trial. *Proc. Am. Soc. Clin. Oncol.*, 17: 377, 1998.

Clinical Cancer Research

Antitumor Activity of Sequential Treatment with Topotecan and Anti-Epidermal Growth Factor Receptor Monoclonal Antibody C225

Fortunato Ciardiello, Roberto Bianco, Vincenzo Damiano, et al.

Clin Cancer Res 1999;5:909-916.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/5/4/909>

Cited articles This article cites 45 articles, 20 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/5/4/909.full#ref-list-1>

Citing articles This article has been cited by 59 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/5/4/909.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/5/4/909>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.