

# Wortmannin Inhibits PKB/Akt Phosphorylation and Promotes Gemcitabine Antitumor Activity in Orthotopic Human Pancreatic Cancer Xenografts in Immunodeficient Mice<sup>1</sup>

Sylvia S. W. Ng, Ming-Sound Tsao, Trudley Nicklee, and David W. Hedley<sup>2</sup>

Divisions of Experimental Therapeutics [S. S. W. N., D. W. H.] and Molecular and Cellular Biology [M-S. T.], Ontario Cancer Institute, and Departments of Medical Oncology and Hematology [D. W. H.], Laboratory Medicine and Pathobiology [M-S. T., T. N.], and Medical Biophysics [S. S. W. N., M-S. T., D. W. H.], Princess Margaret Hospital and University of Toronto, Toronto, Ontario, M5G 2 M9, Canada

## ABSTRACT

Pancreatic cancer is resistant to almost all classes of cytotoxic agents. Gemcitabine seems to be the current drug of choice. We have recently reported that inhibition of the phosphatidylinositol 3-kinase-protein kinase B (PKB/Akt) cell survival pathway by wortmannin enhances gemcitabine-induced apoptosis in cultured human pancreatic cancer cells (1). The present study investigated the effects of wortmannin on orthotopic human pancreatic cancer xenografts implanted in severe combined immunodeficient mice. Animals were given single i.v. bolus injections of 0.175, 0.35, or 0.7 mg/kg of wortmannin and killed at 0.5, 1, 2, or 4 h after treatment. Phosphorylated PKB/Akt levels in tumor tissues were measured by fluorescence image analysis. Wortmannin was found to inhibit PKB/Akt phosphorylation in a time- and dose-dependent manner, reaching a plateau at 4 h and at 0.7 mg/kg. The levels of phosphorylated PKB/Akt were maximally decreased by ~50% relative to the vehicle control. Subsequently, the extent of apoptosis in tumors treated with gemcitabine or wortmannin alone or in combination was determined using terminal deoxynucleotidyl transferase-mediated nick end labeling assay and computerized image analysis. Orthotopic tumors exposed to 80 mg/kg gemcitabine for 48 h and then 0.7 mg/kg wortmannin for 4 h showed a 5-fold increase ( $P = 0.002$ ) in apoptosis compared with those treated with each agent alone and with the vehicle control. The combination treatment also significantly ( $P <$

0.001) inhibited tumor growth. Taken together, our findings support the potential of phosphatidylinositol 3-kinase inhibitors as adjuncts to conventional chemotherapy in the treatment of pancreatic cancer.

## INTRODUCTION

Pancreatic adenocarcinoma is the second most common gastrointestinal malignancy and the fifth leading cause of cancer death in North America (2). Surgery remains the sole curative option. However, fewer than 15% of patients are resectable, and recurrence after a successful pancreatectomy occurs in the majority of these patients (3). Chemotherapy or radiotherapy alone or in combination have a limited effect on the overall survival of patients with pancreatic cancer (2, 4). The current drug of choice seems to be the deoxycytidine analogue gemcitabine (2',2'-difluorodeoxycytidine), despite its <20% objective response rate in the clinic (5). Therefore, the development of more effective treatment is urgently needed to improve quality of life and reduce mortality.

Several genetic abnormalities, which include activating *Ki-ras* mutations (6, 7) and overexpression of multiple receptor tyrosine kinases (8–10), occur at very high frequencies in pancreatic cancers. PI3K,<sup>3</sup> a downstream effector of *Ki-ras* and receptor tyrosine kinases (11–14), activates PKB/Akt (15, 16), which, in turn, has been shown to mediate cell survival via the regulation of numerous proteins such as glycogen synthase kinase-3 $\beta$  (17), BAD (18), nuclear factor  $\kappa$ B and Bcl-X<sub>L</sub> (19). Elevated levels of phosphorylated PKB/Akt can protect cells from undergoing apoptosis induced by cytotoxic drugs and contribute to drug resistance. Given that the most prevalent genetic aberrations found in pancreatic cancers are capable of up-regulating the activity of the PI3K-PKB/Akt cell survival pathway, we hypothesize that its perturbation by pharmacological inhibitors may have therapeutic potential.

The fungal metabolite wortmannin and the synthetic compound LY294002 are the two currently known inhibitors that show fairly high specificity for PI3K. Wortmannin binds to the p110 catalytic subunit of PI3K, noncompetitively and irreversibly inhibiting (IC<sub>50</sub>, 2–4 nM) the enzyme (20). It has been shown that wortmannin at 100 nM causes 95% inhibition of PI3K and that the concentration required to induce apoptosis of cells maintained in growth factors correlates closely with that required for PI3K inhibition (21). There are limited studies reporting the *in vivo* antitumor

Received 2/19/01; revised 7/11/01; accepted 7/11/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Supported by the National Cancer Institute of Canada through funds raised by the Terry Fox Run (to D. W. H.) and Medical Research Council of Canada Grant 14359 (to M-S. T.).

<sup>2</sup> To whom requests for reprints should be addressed, at Department of Medical Oncology and Hematology, Princess Margaret Hospital, 610 University Avenue, Toronto, Ontario, M5G 2 M9, Canada. Phone: (416) 946-2262; Fax: (416) 946-6546; E-mail: david.hedley@uhn.on.ca.

<sup>3</sup> The abbreviations used are: PI3K, phosphatidylinositol 3-kinase; PKB, protein kinase B; SCID, severe combined immunodeficient; MTD, maximum tolerated dose; EGFR, epidermal growth factor receptor; HGFR, hepatocyte growth factor receptor; TUNEL, terminal deoxynucleotidyl transferase-mediated nick end labeling; IOD, integrated optical density.

action of wortmannin (22, 23); however, it is unclear whether the observed effects were caused by inhibition of PI3K and subsequent PKB/Akt phosphorylation. We have recently demonstrated that PI3K inhibitors suppress PKB/Akt phosphorylation and enhance gemcitabine-induced apoptosis in human pancreatic adenocarcinoma cell lines (1). The current study examined the *in vivo* effects of wortmannin on PKB/Akt phosphorylation and apoptosis enhancement in orthotopic human pancreatic cancer xenografts implanted in SCID mice.

## MATERIALS AND METHODS

### Orthotopic Model

Human pancreatic adenocarcinoma cells PK1 were obtained from Dr. Masao Kobari (Sendai, Japan). Their genetic characteristics have been described previously (1, 24). All animal experiments were done in accordance with institutional guidelines for animal welfare. In a preliminary study, PK1 cell suspensions were injected both s.c. and orthotopically into SCID mice to determine the take rate. Animals were sacrificed when tumors reached  $\sim 12 \times 12 \times 12$  mm; harvested orthotopic tumors were cut into pieces of  $\sim 3 \times 3 \times 3$  mm. Male SCID mice 6 weeks of age weighing 18–23 g, were anesthetized with isoflurane (Janssen Pharmaceutica, North York, Ontario, Canada). An incision was made in the upper left abdomen, and the pancreas was exposed. Tumor pieces were attached to the pancreas using 4-0 chromic gut suture (Roboz Surgical Instrument Co., Inc., Rockville, MD). The pancreas was then returned to the peritoneum, and the abdominal wall and the skin were closed with 4-0 chromic gut and silk sutures (Roboz Surgical Instrument Co., Inc.), respectively. The animals were allowed to recover.

### Immunohistochemistry

Five- $\mu$ m-thick sections of paraffin-embedded, formalin-fixed s.c. and orthotopic tumor tissues were stained with H&E for histological examination. For detection of phosphorylated PKB/Akt, EGFR, Met/HGFR, and Her2/neu, sections were stained with a polyclonal antibody specific for PKB/Akt phosphorylated at serine 473 (1:10; New England Biolabs, Mississauga, Ontario, Canada), a monoclonal anti-EGFR antibody (1:100; Zymed Laboratories, Inc., San Francisco, CA), polyclonal anti-Met/HGFR, and anti-Her2/neu antibodies (1:100; Santa Cruz Biotechnology, Santa Cruz, CA), respectively. Antigens were visualized using the streptavidin-biotin-peroxidase method.

### Treatment Protocols

Gemcitabine was obtained from Eli Lilly & Co. (Indianapolis, IN) and dissolved at 20 mg/ml in 0.9% NaCl. Wortmannin was purchased from BioMol (Philadelphia, PA), dissolved at 0.4 mg/ml in DMSO, and diluted with 0.9% NaCl before use.

In the first set of experiments, groups of tumor-bearing SCID mice ( $n = 4$  each) were given single bolus injections of 0.7 mg/kg wortmannin via the tail vein and killed at various time points. This dose of wortmannin was reported to be the MTD in mice (22). In a second series of experiments, groups of animals ( $n = 4$  each) were injected i.v. with increasing doses of wortmannin and killed at 4 h. The lungs, heart, brain, and tumors of each animal were harvested for Western blotting. The remaining tumor tissues were embedded in OCT (Miles, Inc., Elkhart, IN)

and snap-frozen in liquid nitrogen for subsequent immunofluorescence staining. In a third series of experiments, tumor-bearing SCID mice were randomly assigned to four groups ( $n = 4$  each). Each group was given i.v. bolus injections of either the drug vehicle ( $\leq 1\%$  DMSO), 80 mg/kg of gemcitabine for 48 h, 0.7 mg/kg of wortmannin for 4 h, or 80 mg/kg of gemcitabine for 48 h and then 0.7 mg/kg wortmannin for 4 h. These schedules were based on *in vitro* experiments treating PK1 cells with the two drugs (1). Harvested tumors were snap-frozen in OCT (Miles, Inc.) in liquid nitrogen and subsequently processed for TUNEL assay.

### Quantification of Phosphorylated PKB/Akt

**Western Blot Analysis.** PKB/Akt phosphorylation was used as the end point for PI3K activation. Briefly, organs and tumors were homogenized in 500  $\mu$ l of lysis buffer [1% Triton X-100, 0.1% SDS, 50 mM Tris (pH 8.0), 150 mM NaCl, 1 mM PMSF, 0.1 mM  $\text{NaVO}_4$ , 0.1 mM benzamidine, 5  $\mu$ g/ml leupeptin, and 5  $\mu$ g/ml aprotinin]. Homogenates were clarified by centrifugation at 15,000 rpm for 15 min at 4°C. Samples were heated in SDS sample buffer for 5 min at 95°C, run on 12% SDS-polyacrylamide gels, and transferred to nitrocellulose membranes (Bio-Rad Laboratories, Mississauga, Ontario, Canada) using the Mini *Trans*-Blot Electrophoresis Transfer Cell (Bio-Rad Laboratories). Membranes were blocked overnight at 4°C with 10% nonfat milk in TBST [10 mM Tris (pH 7.6), 150 mM NaCl, and 0.5% Tween 20] and then exposed to a primary antibody specific for PKB/Akt phosphorylated at serine 473 (New England Biolabs) for 1 h at room temperature. Secondary antibody containing the horseradish peroxidase detection system for chemiluminescence was used as recommended by the manufacturer (New England Biolabs).

**Indirect Immunofluorescence.** Because the tumors were interspersed with variable amounts of necrotic and non-malignant tissues, a quantitative immunofluorescence technique was developed to examine phosphoPKB/Akt levels in viable tumor areas. Three serial sections (5  $\mu$ m thick) were obtained from each frozen tumor, mounted on glass slides, and then fixed in 4% paraformaldehyde. The first section was stained with H&E for transmitted light microscopy. The remaining sections were incubated with a rabbit polyclonal anti-phosphoPKB/Akt (serine 473) antibody (1:10; New England Biolabs) or a sheep polyclonal anti-PKB/Akt (1:50; Upstate Biotechnology, Lake Placid, NY) for 1 h. A Cy3-conjugated donkey antirabbit IgG and a Cy5-conjugated donkey antisheep IgG (1:200; Molecular Probes, Eugene, OR) were used as secondary antibodies for phosphoPKB/Akt and PKB/Akt, respectively. Control samples exposed to secondary antibodies alone showed no specific staining. All sections were counterstained with the DNA-specific dye diamidinophenyl indole (Molecular Probes) at 1  $\mu$ g/ml to outline the nuclear area.

**Computerized Image Analysis.** Composite digital images of the entire H&E sections were captured using a color CCD video camera (Sony DXC 970 MD) mounted on a Zeiss Axioskop transmitted light microscope fitted with an autostage and linked to a MicroComputer Image Device (Imaging Research, Inc., St. Catherine, Ontario, Canada). These images were used to select representative areas of viable tumor tissues for analysis of the fluorescence markers. Viable tumor

tissues labeled with the anti-phosphoPKB/Akt or anti-PKB/Akt antibodies were examined using a second MicroComputer Image Device image analysis system equipped with a Xillix MicroImager (Xillix, Vancouver, British Columbia, Canada) mounted on an Olympus BX50 reflected fluorescence microscope. Each individual field was obtained using a  $\times 40$  objective. Twenty random 0.149-mm<sup>2</sup> fields of the same tumor and four tumors/group were analyzed. The Cy3- and Cy5-conjugated secondary antibodies were excited using 540 nm and 620 nm bandpass filters with emissions collected at 610 nm and 700 nm, respectively. For phosphoPKB/Akt staining, the nuclear image obtained with DAPI was used to create a mask, which was then overlaid on the Cy3 fluorescence image, so that only nuclear phosphoPKB/Akt fluorescence was measured. It has been shown that phosphoPKB/Akt translocates to the nucleus and regulates the transcription of genes important in mediating cell survival (25–27). For PKB/Akt staining, total Cy5 fluorescence was measured. The fluorescence intensities of Cy3 and Cy5 were expressed as mean IODs, which are measures of the amounts of labeled phosphoPKB/Akt and PKB/Akt, respectively, in the tumor sections.

#### Apoptosis Assay

Five serial sections (5  $\mu$ m thick) were obtained for each frozen tumor, mounted on glass slides, and then fixed in 4% paraformaldehyde. The first section was processed for H&E staining. TUNEL assay was performed on the remaining four sections using the ApopTag Red kit according to the manufacturer's instructions (Intergen Co., Purchase, NY). Tissue sections processed in the absence of terminal deoxynucleotidyl transferase served as negative controls. The rhodamine-conjugated secondary antibody was excited using a 540 nm bandpass filter with emission collected at 610 nm.

Using the fluorescence image analysis system described above fitted with a Quantix cooled CCD camera (Photometrix, Inc., Tucson, AZ), tiled images of tumor sections subjected to TUNEL assay were acquired using a  $\times 20$  objective. Four sections of the same tumor and four tumors/group were analyzed. Tumors were traced manually with reference to the parallel H&E sections so as to exclude edges, necrotic, and nonmalignant tissues from analysis. Apoptotic nuclei, often consisting of clusters of discrete nuclear fragments, could be readily defined using image analysis criteria so as to reject artifacts. The extent of apoptosis in each tumor, expressed as proportional area, was calculated from the sum of the TUNEL-positive pixel area divided by the total viable tumor area.

#### Antitumor Activity Study

PK1 cells ( $10^7$ ) were injected into the pancreas of SCID mice. Four weeks after tumor implantation, the mice were assigned to one of the following four treatment groups ( $n = 10$  each): (a) vehicle control; (b) gemcitabine, biweekly treatment 80 mg/kg/injection; (c) wortmannin, biweekly treatment 0.35 mg/kg/injection; (d) gemcitabine plus wortmannin, with gemcitabine on Monday and Thursday and wortmannin on Wednesday and Saturday. All groups received treatment via i.p. injection. Mice in all groups were killed after 5 weeks of treatment. Orthotopic tumors were harvested and weighed.

#### Statistics

All values are presented as mean  $\pm$  SE. Comparisons between control and treatment groups were made with ANOVA and then Dunnett's test, with  $P < 0.05$  as the criterion for statistical significance.

## RESULTS

**Characterization of PK1 Orthotopic Xenografts.** Microscopic tumors were detected  $<1$  month after the injection of PK1 cell suspensions into the pancreas (Fig. 1A). All SCID mice developed palpable tumors (100% take rate) in the s.c. and orthotopic sites within 2 months. Weight loss was observed in mice bearing orthotopic but not s.c. tumors. Ascites and metastases were not evident at the time of death. Tumors at both sites were poorly differentiated carcinoma with similar growth rates. Whereas the orthotopic tumors demonstrated focal glandular cell differentiation, their s.c. counterparts displayed a squamous differentiation; also, phosphorylated PKB/Akt was detected at much higher level in the former than the latter (Fig. 1B). Immunohistochemical staining revealed high levels of EGFR and Met/HGFR, and moderate levels of Her2/neu expression in the orthotopic tumors (Fig. 1C). Therefore, orthotopic xenografts propagated by suturing tumor pieces onto the pancreas were used in all subsequent experiments.

#### *In Vivo* Effects of Wortmannin on PKB/Akt Phosphorylation

**Normal Tissues.** Fig. 2A shows representative Western blots of lung, heart and brain homogenates from SCID mice treated with single i.v. bolus injections of 0.7 mg/kg wortmannin and killed at 0.5, 1, 2, or 4 h. No mortality or acute toxicity in animals given this dose of wortmannin for 24 h was observed in our pilot studies (data not shown). Phosphorylated PKB/Akt was detected in all three organs under basal conditions. In the lungs, the levels of phosphorylated PKB/Akt were markedly decreased, with the effect being evident at 0.5 h and persisting up to 4 h. However, inhibition of PKB/Akt phosphorylation was less prominent and more transient in the heart; the levels of phosphorylated PKB/Akt at 4 h after wortmannin treatment were comparable with those in the vehicle control group. Phosphorylated PKB/Akt levels in the brain remained relatively constant. Illustrated in Fig. 2B are representative Western blots of organ homogenates from SCID mice treated with increasing i.v. doses (0.175, 0.35, and 0.7 mg/kg) of the drug for 4 h. PKB/Akt phosphorylation was inhibited only in the lungs at the highest dose of wortmannin. Lower doses failed to reduce phosphorylated PKB/Akt levels in the organs. Similar results were obtained from organs of four different animals killed at each time point and at each dose. Total PKB/Akt levels in the three organs were not altered by wortmannin (data not shown). These results demonstrate that i.v. administration of wortmannin is able to inhibit PKB/Akt phosphorylation in normal tissues *in vivo*.

**Pancreatic Carcinoma Xenografts.** The levels of phosphorylated PKB/Akt in cryostat sections of vehicle control and treated tumors were determined from immunofluorescence measurements. Bar graphs of the intensities of phosphorylated PKB/Akt staining expressed as mean IODs *versus* the duration and dose of wortmannin treatment are illustrated in Fig. 3, A and B, respectively. PKB/Akt phosphorylation in





Fig. 1 A, H&E staining showing a microscopic tumor in the pancreas of a mouse injected orthotopically with PK1 cell suspensions. B, phosphorylated PKB/Akt immunohistochemical staining of orthotopic and s.c. tumors. C, immunohistochemical staining of orthotopic tumors for epidermal growth factor receptor (EGFR), hepatocyte growth factor receptor (Met/HGFR), and Her2/neu (Her2/neu).

orthotopic tumors was inhibited by wortmannin in a time- and dose-dependent manner, reaching a plateau at 4 h and at 0.7 mg/kg. Total PKB/Akt levels were not different between control and treated tumors (data not shown). The levels of phosphorylated PKB/Akt were significantly decreased by a maximum of ~50% ( $P < 0.05$ ) relative to the vehicle control. Furthermore, Western blotting performed on tumor tissues also demonstrated similar trends of reduction of phosphory-

lated PKB/Akt levels with increasing doses and duration of wortmannin treatment (data not shown).

**Effects of Wortmannin on Gemcitabine-treated Orthotopic Tumors.** On the basis of the effects of wortmannin on PKB/Akt phosphorylation, we then tested its potential in enhancing gemcitabine toxicity *in vivo*. A composite image of a TUNEL-labeled orthotopic tumor treated with 80 mg/kg i.v. gemcitabine for 48 h and then 0.7 mg/kg i.v. wortmannin for 4 h




**Fig. 2** Effects of wortmannin on normal tissues *in vivo*. Representative Western blots showing the levels of phosphorylated PKB/Akt in the lung, heart, and brain of tumor-bearing SCID mice treated with 0.7 mg/kg i.v. wortmannin for increasing duration (A) or with increasing doses of wortmannin for 4 h (B). Similar results were obtained from four separate animals in each group.

is illustrated in Fig. 4A. TUNEL-positive cells in tumor sections presented with fragmented nuclei (Fig. 4A, *close-up*) and were observed throughout the entire tumor. The central area was necrotic. Proportional area, calculated from the sum of the TUNEL-positive pixel area divided by the total viable tumor area, represents the extent of apoptosis in the tumors. Values of proportional area were not significantly different among the four parallel sections from the same tumor (*e.g.*,  $0.023 \pm 0.002$  for one untreated tumor), indicating high intratumoral reproducibility of this parameter. As shown in Fig. 4B, gemcitabine (80 mg/kg for 48 h) or wortmannin (0.7 mg/kg for 4 h) alone did not produce significant increases in apoptosis compared with the vehicle control. However, the extent of apoptosis was significantly increased by 5-fold ( $P = 0.002$ ) in tumors treated with gemcitabine and wortmannin in combination.

***In Vivo* Inhibition of Tumor Growth.** Four, three, and three deaths were noted in the vehicle control, gemcitabine-, and wortmannin-treated groups, respectively, before the end of the 5-week treatment period because of large tumors. Conversely, all mice receiving gemcitabine and wortmannin in combination were alive and exhibited a healthier appearance. Orthotopic tumors were dissected free of surrounding normal tissues and weighed. As shown in Fig. 5, gemcitabine alone and in combination with wortmannin significantly reduced tumor weights by 1.4- and 5-fold ( $P < 0.001$ ) relative to the vehicle control, respectively. The further decrease in tumor weights observed in the combination treatment group was significantly different from the gemcitabine monotherapy group ( $P < 0.001$ ). Wortmannin alone failed to inhibit tumor growth.

## DISCUSSION


Our results demonstrate for the first time that the prototype PI3K inhibitor wortmannin suppresses PKB/Akt phosphory-



**Fig. 3** Effects of increasing time of exposure to 0.7 mg/kg i.v. wortmannin (A) and of increasing doses of wortmannin for 4 h (B) on phosphorylated PKB/Akt levels, expressed as mean IOD  $\pm$  SE, in orthotopic tumors. Mean IODs were obtained from four separate tumors in each group. \*, significantly ( $P < 0.05$ ) different from the respective control.

lation and promotes apoptosis induced by gemcitabine in orthotopic human pancreatic cancer xenografts. More importantly, the increase in apoptosis associated with the combined use of gemcitabine and wortmannin was translated into greater inhibition of tumor growth. The differences in growth pattern observed and the higher levels of phosphorylated PKB/Akt detected at the orthotopic site of implantation are consistent with previous reports (28, 29) showing the importance of trophic factors in tumor growth and the clinical relevance of the orthotopic model in drug evaluation.


Schultz *et al.* (22) have reported previously that wortmannin given *p.o.* at the MTD of 0.75 mg/kg inhibits tumor growth by >60% in the human BxPC-3 pancreatic carcinoma xenograft. Furthermore, LY294002 has been shown to suppress the growth of human ovarian carcinoma xenografts (30). Whether or not the observed *in vivo* antitumor action was caused by PI3K inhibition remains to be elucidated. In the present study, SCID mice bearing orthotopic tumor xenografts were given increasing doses of *i.v.* wortmannin and killed at various time points. Because high levels of PKB/Akt expression have been detected in the lungs, heart, and brain (31–33), Western blotting was performed on homogenates of these organs to determine whether wortmannin possesses any *in*



**Fig. 4** *A*, wide-field image analysis of a TUNEL-labeled orthotopic tumor treated with 80 mg/kg gemcitabine for 48 h and then 0.7 mg/kg wortmannin for 4 h. *Left*, composite image with viable tumor tissue outlined. *Right*, close-up of an individual  $\times 20$  field. *B*, proportional area of tumors undergoing apoptosis in response to the vehicle control, gemcitabine (*gem.*; at 80 mg/kg for 48 h, i.v.) or wortmannin (*wort.*; at 0.7 mg/kg for 4 h, i.v.) alone or in combination. \*, significantly ( $P = 0.002$ ) different from the vehicle control group.

*in vivo* activity. Phosphorylated PKB/Akt was found in all three organs under basal conditions. The blockade of PKB/Akt phosphorylation in the lungs lasted longer than that in the heart. Possible explanations include more rapid metabolism of wortmannin or higher basal levels of PTEN/MMAC1, a negative regulator of the PI3K-PKB/Akt pathway (34, 35), in the latter. The observation that phosphorylated PKB/Akt levels in the brain remained fairly constant suggests the inability of wortmannin to cross the blood brain barrier.

The effects of wortmannin in orthotopic tumors were studied using a quantitative immunofluorescence technique applied to viable tumor tissues in cryostat sections. Phosphorylated PKB/Akt levels were reduced in orthotopic tumors treated with i.v. wortmannin in a time- and dose-dependent manner. However, PKB/Akt phosphorylation was only inhibited by  $\sim 50\%$ , as compared with  $>90\%$  in PK1 monolayer cultures (1). The lower activity of wortmannin in the orthotopic model might be attributed to its unfavorable pharmacokinetic profile or to the development of wortmannin-resistant PI3K *in vivo*, as recently suggested (23). Alternatively, activation of PKB/Akt in the orthotopic model might occur via PI3K-independent pathways such as those involving p38 and MAPKAP kinase 2 (36).



**Fig. 5** Tumor weights in grams (g) in mice treated with the vehicle control, gemcitabine (*gem.*; 80 mg/kg biweekly, i.p.), wortmannin (*wort.*; 0.35 mg/kg biweekly, i.p.) alone or in combination. *a*, significantly ( $P < 0.001$ ) different from the vehicle control group. *b*, significantly ( $P < 0.001$ ) different from the gemcitabine-treated group.

We have recently shown that wortmannin significantly increases apoptosis in gemcitabine-pretreated PK1 cells (1). Having established the ability of wortmannin to suppress PKB/Akt phosphorylation, we therefore asked if gemcitabine-induced apoptosis can be similarly potentiated by wortmannin in the orthotopic model. Fluorescence image analysis of TUNEL-labeled tumor sections revealed that gemcitabine or wortmannin alone did not produce significant increases in apoptotic cells compared with the vehicle control. However, gemcitabine but not wortmannin monotherapy demonstrated some antitumor activity. Contrary to our data, other groups have reported inhibition of tumor growth by PI3K inhibitors (22, 30). Such discrepancies might be attributable to the use of differential doses, schedules, and cell lines. Tumors treated with gemcitabine and wortmannin in combination showed a significant 5-fold increase in apoptosis and decrease in weight *versus* the vehicle control. Our TUNEL measurements demonstrated high intratumoral reproducibility; hence, any observed differences were attributable to intertumor variations consequent to treatments. Taken together, these results suggest a place for PI3K inhibitors as adjuncts in the treatment of pancreatic cancer. Although wortmannin demonstrated efficacy in potentiating apoptosis and tumor growth inhibition in pancreatic cancer *in situ*, its ability to inhibit PKB/Akt phosphorylation in normal tissues remains a concern. Wortmannin administered daily at twice the MTD for 10 days has been reported to cause 100% mortality (22), indicating its narrow therapeutic index. Nonetheless, wortmannin can serve as a lead for the development of PI3K inhibitors with better pharmacokinetic and pharmacodynamic properties.

In summary, we conclude that wortmannin inhibits PKB/Akt phosphorylation and enhances apoptosis as well as tumor growth inhibition induced by gemcitabine in orthotopic human pancreatic cancer xenografts *in vivo*. Our findings further support the preclinical development of treatment protocols that target the PI3K-PKB/Akt cell survival pathway to be used in pancreatic cancer patients.

## ACKNOWLEDGMENTS

We thank Rebecca Fine and James Ho for their technical assistance.

## REFERENCES

- Ng, S. S. W., Tsao, M. S., Chow, S., and Hedley, D. W. Inhibition of phosphatidylinositol 3-kinase enhances gemcitabine-induced apoptosis in human pancreatic cancer cells. *Cancer Res.*, *60*: 5451–5455, 2000.
- Rosenberg, L. Treatment of pancreatic cancer. *Int. J. Pancreatol.*, *22*: 81–93, 1997.
- Regine, W. F., John, W. J., and Mohiuddin, M. Current and emerging treatments for pancreatic cancer. *Drugs Aging*, *11*: 285–295, 1997.
- Kollmannsberger, C., Peters, H. D., and Fink, U. Chemotherapy in advanced pancreatic adenocarcinoma. *Cancer Treat. Rev.*, *24*: 133–156, 1998.
- Burriss, H. A., Moore, M. J., Andersen, J., Greem, M. R., Rothenberg, M. I., Modiano, M. R., Cripps, M. C., Portenoy, R. K., Storniolo, A. M., Tarassoff, P., Nelson, R., Dorr, F. A., Stephens, C. D., and von Hoff, D. D. Improvements in survival and clinical benefit with gemcitabine as first-line therapy for patients with advanced pancreatic cancer: a randomized trial. *J. Clin. Oncol.*, *15*: 2403–2413, 1997.
- Pellegata, N. S., Sessa, F., Renault, B., Bonato, M., Leone, B. E., Solcia, E., and Ranzani, G. N. *K-ras* and *p53* gene mutations in pancreatic cancer: ductal and nonductal tumors progress through different genetic lesions. *Cancer Res.*, *54*: 1556–1560, 1994.
- Hruban, R. H. and van Manfeld, A. D. M. *K-ras* oncogene activation in adenocarcinoma of the human pancreas. *Am. J. Pathol.*, *143*: 545–554, 1993.
- Oikawa, T., Hitomi, J., Kono, A., Kaneko, E., and Yamaguchi, K. Frequent expression of genes of receptor tyrosine kinases and their ligands in human pancreatic cancer cells. *Int. J. Pancreatol.*, *18*: 15–23, 1995.
- Korc, M., Chandrasekar, B., Yamanaks, Y., Friess, H., Bucher, M., and Beger, H. G. Overexpression of the epidermal growth factor receptor in human pancreatic cancer is associated with concomitant increases in the levels of epidermal growth factor and transforming growth factor  $\alpha$ . *J. Clin. Investig.*, *90*: 1352–1360, 1992.
- Korc, M. Role of growth factors in pancreatic cancer. *Surg. Oncol. Clin. N. Am.*, *7*: 25–41, 1998.
- Ullich, A., and Schlessinger, J. Signal transduction by receptors with tyrosine kinase activity. *Cell*, *61*: 203–212, 1990.
- Rodriguez-Viciana, P., Warne, R., Dhand, B., Vanhaesebroeck, B., Gout, I., Fry, M. J., Waterfield, M. D., and Downward, J. Phosphatidylinositol-3-OH kinase as a direct target of Ras. *Nature (Lond.)*, *370*: 527–532, 1994.
- Rodriguez-Viciana, P., Warne, R., Vanhaesebroeck, B., Waterfield, M. D., and Downward, J. Activation of phosphatidylinositol 3-kinase by interaction with Ras and by point mutation. *EMBO J.*, *15*: 2442–2451, 1996.
- van Weering, D. H., de Rooji, J., Marte, B., Downward, J., Bos, J. L., and Burgering, B. M. Protein kinase B activation and lamellipodium formation are independent phosphoinositide 3-kinase-mediated events differentially regulated by endogenous ras. *Mol. Cell. Biol.*, *18*: 1802–1811, 1998.
- Franke, T. F., Yang, S. I., Chan, T. O., Datta, K., Kazlauskas, A., Morrison, D. K., Kaplan, D. R., and Tschlis, P. N. The protein kinase encoded by the *Akt* oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase. *Cell*, *81*: 727–736, 1995.
- Burgering, B. M., and Coffey, P. J. Protein kinase B (c-Akt) in phosphoinositide-3-OH kinase signaling. *Nature (Lond.)*, *376*: 599–602, 1995.
- Crowder, R. J., and Freeman, R. S. Glycogen synthase kinase-3 $\beta$  activity is critical for neuronal death caused by inhibiting phosphatidylinositol kinase or Akt but not for death caused by nerve growth factor withdrawal. *J. Biol. Chem.*, *275*: 34266–34271, 2000.
- Graff, J. R., Konicek, B. W., McNulty, A. M., Wang, Z., Houck, K., Allen, S., Paul, J. D., Hbawi, A., Goode, R. G., Sandusky, G. E., Vessella, R. L., and Neubauer, B. L. Increased AKT activity contributes to prostate cancer progression by dramatically accelerating prostate tumor growth and diminishing p27<sup>Kip1</sup> expression. *J. Biol. Chem.*, *275*: 24500–24505, 2000.
- Jones, R. G., Parsons, M., Bonnard, M., Chan, V. S., Yeh, W. C., Woodgett, J. R., and Ohashi, P. S. Protein kinase B regulates T lymphocyte survival, nuclear factor  $\kappa$ B activation and Bcl-X(L) levels *in vivo*. *J. Exp. Med.*, *191*: 1721–1734, 2000.
- Powis, G., Bonjouklian, R., Berggren, M. M., Gallegos, A., Abraham, R., Ashendel, C., Zalkow, L., Matter, W. F., Dodge, J., and Grindey, G. Wortmannin, a potent and selective inhibitor of phosphatidylinositol-3-kinase. *Cancer Res.*, *54*: 2419–2423, 1994.
- Yao, R., and Cooper, G. M. Requirement for phosphatidylinositol 3-kinase in the prevention of apoptosis by nerve growth factor. *Nature (Lond.)*, *267*: 2003–2006, 1995.
- Schultz, R. M., Merriman, R. L., Andis, S. L., Bonjouklian, R., Grindey, G., Rutherford, P. G., Gallegos, A., Massey, K., and Powis, G. *In vitro* and *in vivo* antitumor activity of the phosphatidylinositol-3-kinase inhibitor, wortmannin. *Anticancer Res.*, *15*: 1135–1140, 1995.
- Lemke, L. E., Paine-Murrueta, G. D., Taylor, C. W., and Powis, G. Wortmannin inhibits the growth of mammary tumors despite the existence of a novel wortmannin-insensitive phosphatidylinositol-3-kinase. *Cancer Chemother. Pharmacol.*, *44*: 491–497, 1999.
- Liu, N., Furukawa, T., Kobari, M., and Tsao, M. S. Comparative phenotype studies of duct epithelial cell lines derived from normal human pancreas and pancreatic carcinoma. *Am. J. Pathol.*, *153*: 263–269, 1998.
- Coffer, P. J., Jin, J., and Woodgett, J. R. Protein kinase B (c-Akt): a multifunctional mediator of phosphatidylinositol activation. *Biochem. J.*, *335*: 1–13, 1998.
- Alessi, D. R., and Cohen, P. Mechanisms of activation and function of protein kinase B. *Curr. Opin. Gene Dev.*, *8*: 55–62, 1998.
- Downward, J. Mechanisms and consequences of activation of protein kinase B/Akt. *Curr. Opin. Cell. Biol.*, *10*: 262–267, 1998.
- Takahashi, Y., Mai, M., Wilson, M. R., Katadai, Y., Bucana, C. D., and Ellis, L. M. Site-dependent expression of vascular endothelial growth factor, angiogenesis and proliferation in human gastric carcinoma. *Int. J. Oncol.*, *8*: 701–705, 1996.
- Mohammad, R. M., Dugan, M. C., Mohamed, A. N., Almatchy, V. P., Flake, T. M., Dergham, S. T., Shields, A. F., Al-Katib, A. A., Vaitekevicius, V. K., and Sarkar, F. H. Establishment of a human pancreatic tumor xenograft model: potential application for preclinical evaluation of novel therapeutic agents. *Pancreas*, *16*: 19–25, 1998.
- Hu, L., Zaloudek, C., Mills, G. B., Gray, J., and Jaffe, R. B. *In vivo* and *in vitro* ovarian carcinoma growth inhibition by a phosphatidylinositol 3-kinase inhibitor (LY294002). *Clin. Cancer Res.*, *6*: 880–886, 2000.
- Coffer, P. J., and Woodgett, J. R. Molecular cloning and characterization of novel putative protein-serine kinase related to the cAMP-dependent and protein kinase C families. *Eur. J. Biochem.*, *201*: 475–481, 1991.
- Jones, P. F., Jakubowicz, T., Pitossi, F. J., Mauer, F., and Hemmings, B. A. Molecular cloning and identification of a serine/threonine protein kinase of the second messenger subfamily. *Proc. Natl. Acad. Sci. USA*, *88*: 4171–4175, 1991.
- Bellacosa, A., Franke, T. F., Gonzalez-Portal, M. E., Datta, K., Taguchi, T., Gardner, J., Cheng, J. Q., Testa, J. R., and Tschlis, P. N. Structure, expression and chromosomal mapping of c-akt: relationship to v-akt and its implications. *Oncogene*, *8*: 745–754, 1993.
- Maehama, T., and Dixon, J. E. PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-triphosphate. *J. Biol. Chem.*, *273*: 13375–13378, 1998.
- Stambolic, V., Suzuki, A., de la Pompa, J. L., Brothers, G. M., Mirtsos, C., Sasaki, T., Ruland, J., Penninger, J. M., Siderovski, D. P., and Mak, T. W. Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN. *Cell*, *95*: 29–39, 1998.
- Alessi, D. R., Andjelkovic, M., Caudwell, B., Cron, P., Morrice, N., Cohen, P., and Hemmings, B. A. Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J.*, *15*: 6541–6551, 1996.

# Clinical Cancer Research

## Wortmannin Inhibits PKB/Akt Phosphorylation and Promotes Gemcitabine Antitumor Activity in Orthotopic Human Pancreatic Cancer Xenografts in Immunodeficient Mice

Sylvia S. W. Ng, Ming-Sound Tsao, Trudey Nicklee, et al.

*Clin Cancer Res* 2001;7:3269-3275.

**Updated version** Access the most recent version of this article at:  
<http://clincancerres.aacrjournals.org/content/7/10/3269>

**Cited articles** This article cites 34 articles, 11 of which you can access for free at:  
<http://clincancerres.aacrjournals.org/content/7/10/3269.full#ref-list-1>

**Citing articles** This article has been cited by 36 HighWire-hosted articles. Access the articles at:  
<http://clincancerres.aacrjournals.org/content/7/10/3269.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://clincancerres.aacrjournals.org/content/7/10/3269>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.