

Advances in Brief

Expression Analysis of Estrogen Receptor α Coregulators in Breast Carcinoma: Evidence That *NCOR1* Expression Is Predictive of the Response to Tamoxifen¹

Igor Girault, Florence Lerebours, Samira Amarir, Sengül Tozlu, Michèle Tubiana-Hulin, Rosette Lidereau, and Ivan Bièche²

Laboratoire d'Oncogénétique, Institut National de la Santé et de la Recherche Médicale E0017 [I. G., F. L., S. T., R. L., I. B.] and Département de Médecine [M. T.-H.], Centre René Huguenin, F-92211 St-Cloud, and Laboratoire de Génétique Moléculaire, Unité Propre de Recherche et d'Enseignement Supérieur JE 2195, Faculté des Sciences Pharmaceutiques et Biologiques, Université René Descartes, Paris V, F-75006 Paris [I. B.], France

Abstract

Purpose: Dysregulated expression of steroid receptor transcriptional coactivators and corepressors has been implicated in tamoxifen resistance, especially in estrogen receptor (ER) α -positive breast cancer patients. Therefore, expression analysis of these ER α coregulators may identify new predictors of the response to tamoxifen treatment.

Experimental Design: We measured mRNA levels of 16 coactivator and 11 corepressor genes with a real-time quantitative reverse transcription-PCR method in 14 ER α -positive breast tumors. Three selected coactivator genes (*TIF2*, *AIB1*, and *GCN5L2*) and two corepressor genes (*NCOR1* and *MTAIL1*) were additionally investigated in a well-characterized series of ER α -positive unilateral invasive primary breast tumors from 99 postmenopausal patients who only received tamoxifen as adjuvant hormone therapy after primary surgery. We sought relationships between mRNA levels of the coregulators and those of molecular markers, including ER α , ER β , *CCND1*, and *ERBB2*.

Results: ER α coregulator expression was unrelated to age, histological grade, lymph node status, and macroscopic tumor size. The relationship between mRNA expression of the coregulators, and ER α and β only showed a significant positive correlation between *GCN5L2* and ER α ($P = 0.015$). mRNA levels of *CCND1* correlated with those of all of the coregulators studied ($P < 0.05$ or trend), whereas *ERBB2* mRNA levels only correlated with *AIB1* mRNA levels ($P =$

0.011). Low *NCOR1* expression (versus intermediate and high) was associated with significantly shorter relapse-free survival (log-rank test; $P = 0.0076$). The prognostic significance of low *NCOR1* expression persisted in Cox multivariate regression analysis ($P = 0.043$).

Conclusions: These findings point to *NCOR1* as a promising independent predictor of tamoxifen resistance in patients with ER α -positive breast tumors.

Introduction

ER α belongs to the superfamily of steroid nuclear receptor transcriptional factors. It regulates the proliferation and differentiation of many tissues, especially reproductive tissues (1). On binding to specific DNA sequences such as EREs, estrogen-ER α complexes activate or repress target gene transcription. The biological activity of estrogen is now realized to be more complex than initially thought, with the discovery of a second ER named ER β (2). The ER-mediated transcriptional activity of estrogen is influenced by several regulatory factors known as coactivators and corepressors, which activate or repress the transcription of ER-responsive genes (1). Up to now, the enhancer activity of these coregulators has mainly been studied with ER α , but it may differ between the two ERs, providing additional regulatory steps in estrogen signaling (3). Transcriptional activation is triggered, via histone acetyltransferase activity, by protein-protein interactions between the receptor and cofactors, and also through interactions with the SWI/SNF chromatin remodeling complex; together, these mechanisms result in nucleosome disruption and allow RNA polymerase complex transcription activity to take place (4). Transcription silencing is also based on protein-protein interactions, this time between the receptor and corepressors, the latter belonging to larger protein complexes with histone deacetylase activity, which promotes nucleosome condensation (5).

Tamoxifen is the most common endocrine agent used at all stages of breast cancer and particularly for the treatment of postmenopausal patients. ER α status has been used to identify breast cancer patients who are likely to respond to tamoxifen, but resistance nonetheless occurs in 50% of treated ER α -positive breast cancer patients (6). This tamoxifen resistance could be because of dysregulation of ER α coregulator expression in breast tumors (7). Indeed, in addition to the competitive antag-

Received 4/16/02; revised 9/6/02; accepted 9/16/02.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by the Comité Régional des Hauts-de-Seine de la Ligue Nationale Contre le Cancer.

² To whom requests for reprints should be addressed, at Laboratoire d'Oncogénétique, Institut National de la Santé et de la Recherche Médicale E0017, Centre René Huguenin, 35 rue Dailly, F-92211 St-Cloud, France. Phone: 33-1-47-11-15-66; Fax: 33-1-47-11-16-96; E-mail: i.bièche@stcloud-huguenin.org.

³ The abbreviations used are: ER, estrogen receptor; ERE, estrogen-responsive element; RT-PCR, reverse transcription-PCR; TIF2, transcriptional intermediary factor 2; AIB1, augmented in breast cancer 1; GCN5L2, general control of amino acid synthesis, yeast, homologue-like 2; NCOR1, nuclear receptor corepressor 1; MTA1L1, metastasis-associated 1-like 1; PR, progesterone receptor; Ct, cycle threshold; TBP, TATA box-binding protein; RFS, relapse-free survival.

Table 1 Coregulator genes mRNA levels in 14 ER α -positive breast tumors

Genes	Genbank accession no.	Mean \pm SD	Range
Coactivators			
<i>SRC1 (NCOA1)</i>	NM_003743	1.7 \pm 0.6	1–2.8
<i>TIF2^a (NCOA2)</i>	NM_006540	6.0 \pm 4.7	1–17.6
<i>AIB1 (NCOA3)</i>	NM_006534	1.8 \pm 0.6	1–3.1
<i>ARA70 (NCOA4)</i>	NM_005437	2.3 \pm 0.8	1–4.4
<i>ARA54 (RNF14)</i>	NM_004290	2.2 \pm 1.0	1–4.6
<i>TIF1</i>	NM_003852	3.1 \pm 1.8	1–6.9
<i>CARM1</i>	XM_032719	3.7 \pm 2.3	1–8.9
<i>SRCAP</i>	NM_006662	2.4 \pm 1.1	1–5.1
<i>P300 (EP300)</i>	NM_001429	2.3 \pm 1.5	1–5.8
<i>CBP (CREBBP)</i>	NM_004380	3.5 \pm 1.9	1–7.7
<i>GCN5L2</i>	NM_021078	4.0 \pm 3.4	1–13.5
<i>PCAF</i>	NM_003884	4.2 \pm 4.1	1–4.9
<i>BRG1 (SMARCA4)</i>	NM_003072	1.7 \pm 0.4	1–2.5
<i>hBRM (SMARCA2)</i>	NM_003070	3.1 \pm 2.0	1–9.3
<i>SNF5 (SMARCB1)</i>	NM_003073	3.1 \pm 1.0	1–4.4
<i>BAF60b (SMARCD2)</i>	NM_003077	5.4 \pm 4.9	1–8.9
Corepressors			
<i>NCOR1</i>	NM_006311	5.7 \pm 3.4	1–15.5
<i>SMRT (NCOR2)</i>	NM_006312	2.9 \pm 1.2	1–5.2
<i>REA</i>	NM_007273	4.5 \pm 2.6	1–12.2
<i>HDAC1</i>	NM_004964	2.1 \pm 0.7	1–3.2
<i>HDAC2</i>	NM_001527	2.9 \pm 2.4	1–11.1
<i>HDAC3</i>	NM_003883	2.1 \pm 1.2	1–6.4
<i>SIN3B (KIAA0700)</i>	XM_050561	2.2 \pm 1.4	1–6.8
<i>SAP18</i>	NM_005870	2.4 \pm 1.2	1–5.5
<i>SAP30</i>	NM_003864	2.2 \pm 1.5	1–7.0
<i>MTA1</i>	NM_004689	2.2 \pm 1.0	1–4.9
<i>MTAIL1</i>	NM_004739	6.2 \pm 3.9	1–14.5

^a Bold characters: coregulator genes selected to be additionally studied in the series of 99 ER α -positive breast tumors.

onistic effect of tamoxifen for the ER α ligand-binding site, the main mechanism underlying the antiproliferative activity of this drug (8), tamoxifen could also inhibit cell proliferation by promoting apoptosis, as shown *in vitro* and *in vivo* (9, 10). Tamoxifen-induced apoptosis may be associated with the recruitment of coregulators, which do not normally interact with estrogen-ER α complexes, leading to abnormal transcriptional regulation of ER α target genes (11).

To identify the ER α coregulators involved in tamoxifen resistance, we used real-time quantitative RT-PCR assays to analyze the expression of a large panel of coregulators (16 coactivators and 11 corepressors; Table 1) in 14 ER α -positive breast tumors. We then additionally investigated five coregulators of interest, comprising three coactivators, *TIF2* (12), *AIB1* (13), and *GCN5L2* (14) and two corepressors, *NCOR1* (15) and *MTAIL1* (16), in a well-characterized series of 99 ER α -positive unilateral invasive primary breast tumors from postmenopausal patients who exclusively received tamoxifen as adjuvant hormone therapy after primary surgery. Relationships between the mRNA levels of these coregulators, and clinical and pathological parameters, including RFS, were then studied. Finally, we sought relationships between mRNA levels of the coregulators and those of well-known molecular markers in breast cancer (ER α , ER β , and PR), and two candidate predictors of the response to endocrine therapy, *CCND1* (17) and *ERBB2* (18), of which the expression has been shown recently to be regulated by ER α coactivator genes (19, 20).

Table 2 Characteristics of the 99 postmenopausal patients with ER α -positive breast tumors and relation to RFS

	RFS		<i>P</i> ^b
	Number of patients	Number of events ^a (%)	
Age			NS ^c
≤70	50	20 (40.0)	
>70	49	13 (26.5)	
Histological grade ^d			0.0012
I	13	2 (15.4)	
II	64	17 (26.6)	
III	21	13 (61.9)	
Lymph node status			0.0011
0	16	2 (12.5)	
1–3	56	15 (26.8)	
>3	27	16 (59.3)	
Macroscopic tumor size ^e			0.015
≤30 mm	67	18 (26.9)	
>30 mm	30	14 (46.7)	

^a First relapses (local and/or regional recurrences, and/or metastases).

^b *P* (log-rank test).

^c NS, not significant.

^d Scarff Bloom Richardson classification. Information available for 98 patients.

^e Information available for 97 patients.

Materials and Methods

Patients and Samples

We analyzed samples of primary breast tumors excised from 99 women at Centre René Huguenin from 1980 to 1994. The samples were examined histologically for the presence of tumor cells. A tumor sample was considered suitable for this study if the proportion of tumor cells was >60%. Immediately after surgery, the tumor samples were placed in liquid nitrogen until total RNA extraction.

The patients (mean age, 70.7 years; range, 54–86) met the following criteria: primary unilateral ER α -positive nonmetastatic postmenopausal breast carcinoma; complete clinical, histological, and biological information available; no radiotherapy or chemotherapy before surgery; and full follow-up at Centre René Huguenin. The histological type and the number of positive axillary nodes were established at the time of surgery. The malignancy of infiltrating carcinomas was scored according to the histoprosthetic system of Bloom and Richardson (21). ER α -positive status was determined at the protein level by using biochemical methods (dextran-coated charcoal method until 1988 and enzymatic immuno-assay thereafter) and confirmed by ER α real-time quantitative RT-PCR assay. Standard prognostic factors are reported in Table 2. Thirty patients had modified radical mastectomy, and 69 had breast-conserving surgery plus locoregional radiotherapy. The patients underwent physical examinations and routine chest radiography every 3 months for 2 years, then annually. Mammograms were done annually. The median follow-up was 6 years (range, 1.5–17.5 years). All of the patients received postoperative adjuvant endocrine therapy (tamoxifen, 20 mg daily for 3–5 years) and no other treatment. Thirty-three patients relapsed. The first relapse events consisted of local and/or regional recurrences in 3 patients, metastases in 26 patients, and both events in 4 patients.

Table 3 Oligonucleotide primer sequences

Gene	Oligonucleotide	Sequence	PCR product size (bp)
<i>TIF2</i>	Upper primer	5'-GCT GGG AGG ACC TGG TAA GAA-3'	117
	Lower primer	5'-TGA ATG CCA ATC CTT GTC TCA G-3'	
<i>AIB1</i>	Upper primer	5'-GAC CGC TTT TAC TTC AGG CAT T-3'	125
	Lower primer	5'-TGT GTT AAC CAG GTC CTC TTG CT-3'	
<i>GCN5L2</i>	Upper primer	5'-CTT CAG TCA GTG CAG CGG TTG-3'	123
	Lower primer	5'-TCC TCT TCT CGC CTG GCA TAG-3'	
<i>NCOR1</i>	Upper primer	5'-CCC AGC AAC GAG AGG AAT CA-3'	91
	Lower primer	5'-GTC CAT GGG AGG AGT GCT TGT-3'	
<i>MTA1L1</i>	Upper primer	5'-CCG ACG GCC TTA TGC TCC T-3'	145
	Lower primer	5'-CTG GGC CAC CAG ATC TTT GAC-3'	
<i>TBP</i>	Upper primer	5'-TGC ACA GGA GCC AAG AGT GAA-3'	132
	Lower primer	5'-CAC ATC ACA GCT CCC CAC CA-3'	

mRNA expression of the 27 coregulators was first determined in 14 ER α -positive samples from among the 99 postmenopausal breast tumors.

Real-Time RT-PCR

Theoretical Basis. Reactions are characterized by the point during cycling when amplification of the PCR product is first detected, rather than the amount of PCR product accumulated after a fixed number of cycles. The higher the starting quantity of the target molecule, the earlier a significant increase in fluorescence is observed. The parameter Ct is defined as the fractional cycle number at which the fluorescence generated by SYBR Green dye-amplicon complex formation passes a fixed threshold above baseline.

The precise amount of total RNA added to each reaction mix (based on absorbance) and its quality (lack of extensive degradation) are both difficult to assess. Therefore, we also quantified transcripts of the gene coding for the TBP (a component of the DNA-binding protein complex TFIID) as the endogenous RNA control and normalized each sample on the basis of its TBP content (22).

The relative target gene expression level was also normalized to the expression in the breast tumor sample (calibrator) from the two series ($n = 14$ or $n = 99$), which contained the smallest amount of target gene mRNA.

Final results, expressed as N-fold differences in target gene expression relative to the *TPB* gene and the calibrator, and termed "N target," were determined as follows: $N \text{ target} = 2^{(\Delta Ct \text{ calibrator} - \Delta Ct \text{ sample})}$, where ΔCt values of the calibrator and sample are determined by subtracting the Ct value of the target gene from the Ct value of the TBP gene.

Primers and PCR Consumables. The primers for the chosen coregulator genes and *TBP* were chosen with the assistance of Oligo 4.0 software (National Biosciences, Plymouth, MN). We conducted BLASTN searches against dbEST and nr (the nonredundant set of the GenBank, EMBL, and DDBJ database sequences) to confirm the total gene specificity of the chosen nucleotide sequences and the absence of DNA polymorphisms. The primer nucleotide sequences for the 5 coregulator genes (and *TBP*) analyzed in the full panel of 99 breast tumors are shown in Table 3; the other coregulator gene primer sequences are available on request. The primer sequences for

ER α , *ER β* , *PR*, *ERBB2*, and *CCND1* have been published elsewhere (22–24). To avoid amplification of contaminating genomic DNA, one of the two primers was placed at the junction between two exons or in a different exon. For example, the upper primer of *TBP* was placed at the junction between exon 5 and 6, and the lower primer in exon 6.

RNA Extraction. Total RNA was extracted from tissue specimens by using the acid-phenol guanidium method (25). The quality of RNA samples was determined by electrophoresis through agarose gels and staining with ethidium bromide; the 18S and 28S RNA bands were visualized under UV light.

cDNA Synthesis. RNA was reverse-transcribed in a final volume of 20 μ l containing 1 \times RT-PCR buffer [500 mM each deoxynucleotide triphosphate, 3 mM MgCl₂, 75 mM KCl, and 50 mM Tris-HCl (pH 8.3)], 10 units of RNasin RNase inhibitor (Promega, Madison, WI), 10 mM DTT, 50 units of Superscript II RNase H⁻ reverse transcriptase (Life Technologies, Inc.), 1.5 mM random hexamers (Pharmacia, Uppsala, Sweden), and 1 μ g of total RNA (patient samples). Samples were incubated at 20°C for 10 min and 42°C for 30 min, and reverse transcriptase was inactivated by heating at 99°C for 5 min and cooling at 5°C for 5 min.

PCR Amplification. All of the PCR reactions were performed using a ABI Prism 7700 Sequence Detection System (Perkin-Elmer Applied Biosystems, Foster City, CA).

For each PCR run, a master mix was prepared on ice with 1 \times SYBR Green buffer, 5 mM MgCl₂, 200 μ M dATP, dCTP, and dGTP, and 400 μ M dUTP, 300 nM each primer, and 1.25 units of AmpliTaq Gold DNA polymerase (Perkin-Elmer Applied Biosystems). Five μ l of each diluted cDNA solution sample was added to 20 μ l of the PCR master-mix. The thermal cycling conditions comprised an initial denaturation step at 95°C for 10 min, and 50 cycles at 95°C for 15 s and 65°C for 1 min.

Statistical Analysis

RFS was determined as the interval between initial diagnosis and detection of the first relapse (local and/or regional recurrence, and/or metastasis). Mean ER α coregulator mRNA levels were compared by using the Kruskal-Wallis test. Spearman's rank correlation test was used to study relationships between continuous variables. Survival distributions were esti-

Table 4 Mean, median, and range of coregulator genes mRNA levels in 99 ER α -positive breast tumors

	Coactivators			Corepressors	
	<i>TIF2</i>	<i>AIB1</i>	<i>GCN5L2</i>	<i>NCOR1</i>	<i>MTAIL1</i>
Mean \pm SD	7.7 \pm 4.5	2.6 \pm 1.5	6.9 \pm 5.3	5.6 \pm 2.9	4.3 \pm 2.0
Median	6.5	2.3	5.1	5.2	4.1
Range	1–31.3	1–12.2	1–35.3	1–16.9	1–18.5

mated by using the Kaplan-Meier method (26), and the significance of differences between survival rates was determined using the log-rank test. Multivariate analysis using Cox's proportional hazards model was used to assess the independent contribution of each variable to RFS (27). Differences between groups were judged significant at confidence levels $>95\%$ ($P < 0.05$).

Results

Expression Analysis of 27 Coregulator Genes in 14 ER α -Positive Breast Tumors

We measured mRNA expression levels of 27 ER α coregulator genes in a series of 14 ER α -positive breast tumor samples (Table 1). The widest ranges of values for a coactivator gene and a corepressor gene were, respectively, from 1 to 17.6 for *TIF2* and from 1 to 15.5 for *NCOR1*. We then additionally studied selected genes in a larger series of 99 ER α -positive breast tumors. The selection was based on: (a) the widest range of expression in the initial series of 14 breast tumor samples; (b) selection of two genes from each coactivator and corepressor type of genes; and (c) coregulator genes reported previously to be dysregulated in breast cancer (*AIB1*; Ref. 13).

Thus, we selected two coactivators that interact directly with ER α (*TIF2* and *AIB1*), a coactivator involved in nucleosome disruption (*GCN5L2*), a corepressor that interacts with ER α (*NCOR1*), and a corepressor belonging to a histone deacetylase complex (*MTAIL1*).

Expression Analysis of Five Coregulator Genes in 99 Postmenopausal ER α -Positive Breast Tumors

Coregulator mRNA Steady-State Levels in Breast Tumor Samples. Table 4 shows the mean, median, and range of *TIF2*, *AIB1*, *GCN5L2*, *NCOR1*, and *MTAIL1* mRNA levels in the 99 postmenopausal ER α -positive breast tumors. The range of *TIF2* and *GCN5L2* expression exceeded 30-fold, whereas that of the other three coregulators did not exceed 20-fold.

Relationships among Coregulator mRNA Levels. We then sought links between mRNA values for each of the coregulators by using the Spearman rank correlation test (Table 5). Among the coactivators, *TIF2* mRNA levels correlated with both *AIB1* and *GCN5L2* mRNA levels ($P = 0.00057$, $r = +0.346$, and $P = 0.0029$, $r = +0.298$, respectively). A positive correlation was also found between *AIB1* and *GCN5L2* mRNA levels ($P = 0.027$, $r = +0.220$). Regarding the two corepressors, a weak positive correlation was observed between their mRNA levels ($P = 0.024$, $r = +0.225$). The following strong positive correlations were observed between coactivators and corepressors: *AIB1* (coactivator) and *MTAIL1* (corepressor);

Table 5 Relationships between coregulator genes mRNA levels in 99 ER α -positive breast tumors

	Coactivators			Corepressors	
	<i>TIF2</i>	<i>AIB1</i>	<i>GCN5L2</i>	<i>NCOR1</i>	
<i>AIB1</i>	+0.346 ^a 0.00057 ^b				
<i>GCN5L2</i>	+0.298 0.0029	+0.220 0.027			
<i>NCOR1</i>	+0.371 0.00022	+0.073 NS ^c	+0.398 0.000077		
<i>MTAIL1</i>	+0.240 0.016	+0.378 0.00017	+0.174 NS (0.080)	+0.225 0.024	

^a Spearman correlation coefficient.

^b P , Spearman rank correlation test.

^c NS, not significant.

$P = 0.00017$, $r = +0.378$), *TIF2* and *NCOR1* ($P = 0.00022$, $r = +0.371$), and *GCN5L2* and *NCOR1* ($P = 0.000077$, $r = +0.398$). A weak positive correlation was also found between *TIF2* and *MTAIL1* mRNA levels ($P = 0.016$, $r = +0.240$).

Relationships between Coregulator mRNA Levels, and Clinical and Pathological Parameters. Coregulator mRNA levels were compared among patient subgroups defined by clinical and pathological parameters (age, lymph node status, histological grade, and macroscopic tumor size) using the Kruskal-Wallis test (Table 6). No difference in mean coregulator mRNA levels was found.

Relationships between Coregulator and ER α , ER β , PR, CCND1, and ERBB2 mRNA Levels. We then looked for relationships between mRNA levels of coregulators and those of classical molecular markers, *i.e.*, ER α , ER β , PR, CCND1, and ERBB2, determined previously using the same methodology (Table 6; Ref. 28).⁴ A positive correlation was only observed between ER α and the coactivator *GCN5L2* (Spearman rank test; $P = 0.015$, $r = +0.243$). A trend toward statistical significance was observed between mRNA levels of ER α and *NCOR1*, a corepressor ($P = 0.061$, $r = +0.187$). No correlation was observed between coregulator and ER β mRNA levels. mRNA levels of PR (a well-known ER α -responsive gene) correlated positively with those of *TIF2* ($P = 0.0083$, $r = +0.263$) and *NCOR1* ($P = 0.0011$, $r = +0.328$), and a trend toward significance was observed with *GCN5L2* ($P = 0.054$, $r = +0.192$). Highly significant positive correlations were found between mRNA levels of CCND1 and four coregulators, *AIB1* only showing a trend toward significance ($P = 0.10$). The only significant correlation found for ERBB2 was with *AIB1* ($P = 0.011$, $r = +0.254$).

Prognostic Value of Coregulator mRNA Levels. For the prognostic analysis of each gene, the patient population was divided into tertiles (three groups of 33), corresponding to low, intermediate, and high mRNA levels. Only *NCOR1* showed

⁴ F. Spyrtos, M. Labroquère, M. Tubiana-Hulin, S. Tozlu, M. Vidaud, K. Hacène, R. Lidereau, V. Becette, and I. Bièche. Expression of estrogen receptor beta mRNA in postmenopausal primary breast cancer patients receiving adjuvant tamoxifen, submitted for publication.

Table 6 Relationships between coregulator genes mRNA levels and clinical, pathological, and molecular parameters in ER α -positive breast tumors of 99 postmenopausal patients

		Coactivators			Corepressors						
		<i>TIF2</i>	<i>AIB1</i>	<i>GCN5L2</i>	<i>NCOR1</i>	<i>MTA1L1</i>					
Age			NS ^a	NS	NS	NS	NS	NS	NS	NS	
≤70	50 ^b	8.3 ± 5.4 ^c		3.7 ± 2.3		6.7 ± 5.4		5.8 ± 3.4		4.3 ± 2.1	
>70	49	7.0 ± 3.3		3.4 ± 1.7		7.1 ± 5.3		5.4 ± 2.3		4.0 ± 1.8	
Histological grade ^d			NS	NS	NS	NS	NS	NS	NS	NS	
I	13	8.1 ± 3.8		2.7 ± 0.9		5.4 ± 1.8		6.1 ± 2.6		4.3 ± 1.6	
II	64	8.0 ± 4.6		3.8 ± 2.3		7.2 ± 5.6		5.6 ± 2.6		4.1 ± 2.0	
III	21	6.7 ± 4.5		3.3 ± 1.4		6.6 ± 5.8		5.3 ± 4.0		3.8 ± 1.6	
Lymph node status			NS	NS	NS	NS	NS	NS	NS	NS	
0	16	8.1 ± 6.7		3.2 ± 1.6		7.6 ± 5.9		5.0 ± 2.2		3.6 ± 1.6	
1–3	56	7.5 ± 3.5		3.4 ± 2.2		5.4 ± 6.9		6.0 ± 2.9		4.3 ± 2.1	
>3	27	7.9 ± 4.7		4.0 ± 1.9		4.9 ± 6.8		5.3 ± 3.3		4.1 ± 1.8	
Macroscopic tumor size ^e			NS	NS	NS	NS	NS	NS	NS	NS	
≤30 mm	67	8.0 ± 4.7		3.6 ± 2.1		6.4 ± 4.2		5.7 ± 2.7		3.8 ± 1.4	
>30 mm	30	6.9 ± 4.0		3.4 ± 1.9		7.5 ± 7.2		5.2 ± 3.1		4.7 ± 2.6	
mRNA levels											
<i>ERα</i>	99	+0.125 ^f	NS	-0.075	NS	+0.243	0.015	+0.187	NS (0.06)	+0.008	NS
<i>ERβ</i>	99	-0.047	NS	+0.153	NS	+0.109	NS	+0.170	NS (0.09)	+0.170	NS (0.09)
<i>PR</i>	99	+0.263	0.0083	-0.002	NS	+0.192	NS (0.05)	+0.328	0.0011	+0.135	NS
<i>CCND1</i>	99	+0.328	0.0011	+0.162	NS (0.10)	+0.258	0.0097	+0.272	0.0064	+0.295	0.0031
<i>ERBB2</i>	99	+0.020	NS	+0.254	0.011	+0.153	NS (0.13)	-0.109	NS	+0.05	NS

^a *P*, Kruskal-Wallis test for clinical and pathological parameters and Spearman rank correlation test for molecular mRNA levels.

^b Case number.

^c Mean mRNA levels ± SD.

^d Scarff Bloom Richardson classification. Information available for 98 patients.

^e Information available for 97 patients.

^f Spearman correlation coefficient.

prognostic value. Five-year RFS rates were 58.7% (49.8–67.6%) in the subgroup of low *NCOR1* mRNA levels, and 80.8% (73.7–87.9%) and 83.1% (76.1–90.1%) in the subgroups with intermediate and high mRNA levels, respectively. Given the similar values in the subgroups with intermediate and high *NCOR1* mRNA levels, we compared full follow-up RFS between the low *NCOR1* subgroup (33 patients) and the combined intermediate and high subgroups (66 patients). Univariate analysis (log-rank test) showed that the outcome of the former 33 patients was significantly worse than that of the remaining 66 patients ($P = 0.0076$; Fig. 1).

Using a Cox proportional hazards model, we then assessed the prognostic significance of the four parameters that were significant in univariate analysis (histopathological grade, lymph node status, macroscopic tumor size, and *NCOR1* status; Table 2; Fig. 1). The prognostic significance of histopathological grade, lymph node status, and *NCOR1* status persisted, whereas that of macroscopic tumor size disappeared (Table 7).

Discussion

To gain more insight into the roles of the different ER α coregulator genes in breast tumors, we have chosen a two-step strategy to measure mRNA expression of coregulator genes by using real-time quantitative RT-PCR. The first step was aimed to quantify the mRNA expression of a large panel of genes ($n =$




Fig. 1 RFS in the patient subgroup with low *NCOR1* mRNA levels (33 patients), and the combined subgroups with intermediate and high *NCOR1* mRNA levels (66 patients).

27) coding for coregulators in a small homogeneous series of 14 ER α -positive breast tumors and, thus, to distinguish expressed from unexpressed coregulators in breast tumors. Given the high number of coregulator genes, it is suspected that related genes may show functional redundancy (29). Therefore, in a second step, in addition to *AIB1* (reported previously as being dysregulated in breast cancer; Ref. 13), two other coactivators (*TIF2* and

Table 7 Multivariate analysis of relapse-free survival (RFS)

	RFS		
	Regression coefficient	Relative risk (95% CI) ^a	P
Histological grade	0.83		0.0098
I		1.0	
II		2.3 (1.2–4.3)	
III		5.2 (1.5–18.4)	
Lymph node status	0.83		0.0078
0		1.0	
1–3		2.3 (1.2–4.2)	
>3		5.2 (1.5–17.6)	
Macroscopic tumor size	0.44		0.24
\leq 30 mm		1.0	
>30 mm		1.6 (0.8–3.2)	
NCOR1 mRNA levels	0.75		0.043
Low		2.1 (1.0–4.3)	
Intermediate and high		1.0	

^a 95% confidence interval.

GCN5L2) and two corepressors (*NCOR1* and *MTAIL1*) were selected for their wide range of mRNA expression levels (>10-fold), and were additionally analyzed in a well-characterized series of 99 ER α -positive breast tumors from postmenopausal patients treated exclusively with tamoxifen after surgery.

mRNA levels of the three coactivators correlated positively with one another as well as mRNA levels of the two corepressor genes (Table 5). Positive correlations were also observed in all of the pairwise comparisons of the coactivators and corepressors, except for *AIB1* with *NCOR1* and *GCN5L2* with *MTAIL1*. These correlations may be explained by: (a) transcriptional activation of all five of the genes by a common regulatory pathway; (b) specific up-regulation of one coactivator directly involved in transcriptional activation of the correlated coactivator and/or corepressor; and (c) up-regulation of corepressors aimed at controlling coactivators overexpression.

With respect to clinical and pathological parameters, ER α coregulator expression was unrelated to age, histological grade, lymph node status, and macroscopic tumor size (Table 6). Previous studies of breast tumors have shown a significant relationship between *AIB1* overexpression and high tumor grade (30), and also higher *TIF2* expression levels in node-positive than node-negative breast tumors (31). These discrepancies with our study could be explained in large part by differences in the study populations. Indeed, we studied a well-defined series of postmenopausal patients with ER α -positive tumors.

The only significant relationship between ER α coregulator gene expression, and ER α and ER β expression was a positive correlation between *GCN5L2* and ER α , suggesting an interaction between the transcriptional regulation of the two genes, despite the lack of a consensus ERE sequence in the promoter region of the *GCN5L2* gene (data not shown). Expression of the *TIF2* coactivator and the *NCOR1* corepressor was significantly linked to expression of the PR gene, which contains an ERE element in its promoter, suggesting a predominant role of these two genes in the transcriptional regulation of ER α -responsive genes mediated by ERE.

It has been shown recently that transcriptional regulation of two candidate predictors of the response to endocrine therapy,

the genes *CCND1* and *ERBB2*, can be exerted by ER α coactivators, despite the lack of an ERE in the promoter region of the two genes (19, 20). *AIB1* was found to enhance *CCND1* expression when incubated with estrogen, whereas antiestrogen treatment had no activating or inhibiting effect on *CCND1* transcription. In our study, significant positive correlations were observed between mRNA expression of *CCND1* and that of all of the coregulators but only a trend toward significance with *AIB1*, suggesting that *AIB1* is not a major coactivator involved in *CCND1* transcriptional regulation during breast tumorigenesis. Regarding the *ERBB2* gene, Newman *et al.* (20) showed that p160 coactivators (of which *AIB1* is a member) can modulate its enhancer activity. In our series, *ERBB2* mRNA levels correlated significantly with *AIB1* mRNA levels but not with the other four ER α coregulators. The same relationship was reported at the protein level by Bouras *et al.* (30), who found that tumors overexpressing *AIB1* showed strong ERBB2-positive staining. These results suggest that transcriptional coactivators may establish cross-talk between ERE-dependent and -independent metabolic pathways involved in breast tumorigenesis.

The breast tumor samples studied here were obtained from a well-characterized series of patients treated exclusively with tamoxifen as adjuvant endocrine therapy. Therefore, we were able to evaluate the potential of the five selected ER α coregulator genes as predictors of tamoxifen resistance. The hypothesis of *NCOR1* being a potential marker of antiestrogen hormone-therapy (32) is conceivably related to *in vitro* studies that have established that *NCOR1* protein binds ER α (whether or not the interaction is tamoxifen-dependent is controversial; Refs. 33–35), and inhibits the partial agonistic activity of tamoxifen and its physiological metabolite (4-hydroxy-tamoxifen; Refs. 35, 36). Likewise, in a mouse model of breast cancer, decreased *NCOR1* protein expression correlated with acquired tamoxifen resistance (33).

In human breast tumors, Chan *et al.* (37) did not show significant difference in the level of *SMRT* mRNA (a corepressor related to *NCOR1*) between tamoxifen-resistant breast tumor samples and untreated patients. Conversely, Kurebayashi *et al.* (31) found higher *NCOR1* expression levels in tumors from patients without recurrence compared with patients with recurrence supporting its role as a tamoxifen resistance mechanism *in vivo*.

Here, we found that *NCOR1* mRNA expression status (low versus intermediate and high) was associated with shorter RFS (log-rank test; $P = 0.0076$), and this prognostic significance persisted in Cox multivariate regression analysis ($P = 0.043$). Nevertheless, validation of the predictive value of *NCOR1* parameter in the response to tamoxifen endocrine therapy in breast cancer patients needs a prospective randomized study to show that this parameter does influence the outcome only in patients who received adjuvant tamoxifen compared with untreated patients.

We showed previously, in the same patients series, that *ERBB2* overexpression was a marker of poor prognosis (28). By combining *NCOR1* and *ERBB2* expression status, we identified four separate prognostic groups. The patients with the best prognosis had high *NCOR1* expression and normal *ERBB2* expression (log-rank test; $P = 0.00064$), suggesting that the analysis of the two molecular markers together, affecting ta-

moxifen efficiency, may provide a more accurate prediction of hormone responsiveness (38).

In conclusion, *NCOR1* emerged from a series of 27 ER α coregulator genes as a promising independent prognostic marker of tamoxifen resistance in patients with ER α -positive breast tumors. Therefore, *NCOR1* real-time RT-PCR assay should be valuable in the breast cancer clinical setting to better assess endocrine treatment efficiency.

Acknowledgments

We thank the staff of Centre René Huguenin for assistance with specimen collection and patient care. We also thank Dr. Kamel Hacène (Département de Statistiques Médicales, Centre René Huguenin, St-Cloud, France) for his helpful contribution.

References

- Klinge, C. M. Estrogen receptor interaction with co-activators and co-repressors. *Steroids*, 65: 227–251, 2000.
- Mosselman, S., Polman, J., and Dijkema, R. ER β : identification and characterization of a novel human estrogen receptor. *FEBS Lett.*, 392: 49–53, 1996.
- Wammark, A., Almlöf, T., Leers, J., Gustafsson, J. A., and Treuter, E. Differential recruitment of the mammalian mediator subunit TRAP220 by estrogen receptors ER α and ER β . *J. Biol. Chem.*, 276: 23397–23404, 2001.
- Lemon, B. D., and Freedman, L. P. Nuclear receptor cofactors as chromatin remodelers. *Curr. Opin. Genet. Dev.*, 9: 499–504, 1999.
- Alland, L., Muhle, R., Hou, H., Potes, J., Chin, L., Schreiber-Agus, N., and DePino, R. A. Role for N-CoR and histone deacetylase in Sin3-mediated transcriptional repression. *Nature (Lond.)*, 387: 49–55, 1997.
- McGuire, W. L., Clark, G. M., Dressler, L. G., and Owens, M. A. Role of steroid hormone receptors as prognostic factors in primary breast cancer. *NCI Monogr.*, 1: 19–23, 1986.
- Takimoto, G. S., Graham, J. D., Jackson, T. A., Tung, L., Powell, R. L., Horwitz, L. D., and Horwitz, K. B. Tamoxifen resistant breast cancer: coregulators determine the direction of transcription by antagonist-occupied steroid receptors. *J. Steroid. Biochem. Mol. Biol.*, 69: 45–50, 1999.
- Katzenellenbogen, B. S., Montano, M. M., Ekena, K., Herman, M. E., McInerney, E. M., and William, L. McGuire Memorial Lecture. Antiestrogens: mechanisms of action and resistance in breast cancer. *Breast Cancer Res. Treat.*, 44: 23–38, 1997.
- Ellis, P. A., Saccani-Jotti, G., Clarke, R., Johnston, S. R., Anderson, E., Howell, A., A'Hern, R., Salter, J., Detre, S., Nicholson, R., Robertson, J., Smith, I. E., and Dowsett, M. Induction of apoptosis by tamoxifen and ICI 182780 in primary breast cancer. *Int. J. Cancer*, 72: 608–613, 1997.
- Perry, R., Kang, Y., and Greaves, B. Effects of tamoxifen on growth and apoptosis of estrogen-dependent and independent human breast cancer cells. *Ann. Surg. Oncol.*, 2: 238–245, 1995.
- Smith, C. L., Nawaz, Z., and O'Malley, B. W. Coactivator and corepressor regulation of the agonist/antagonist activity of the mixed antiestrogen, 4-hydroxytamoxifen. *Mol. Endocrinol.*, 11: 657–666, 1997.
- Voegel, J. J., Heine, M. J., Zechel, C., Chambon, P., and Gronemeyer, H. TIF2, a 160 kDa transcriptional mediator for the ligand-dependent activation function AF-2 of nuclear receptors. *EMBO J.*, 15: 3667–3675, 1996.
- Anzick, S. L., Kononen, J., Walker, R. L., Azorsa, D. O., Tanner, M. M., Guan, X. Y., Sauter, G., Kallioniemi, O. P., Trent, J. M., and Meltzer, P. S. AIB1, a steroid receptor coactivator amplified in breast and ovarian cancer. *Science (Wash. DC)*, 277: 965–968, 1997.
- Wang, L., Mizzen, C., Ying, C., Candau, R., Barlev, N., Brownell, J., Allis, C. D., and Berger, S. L. Histone acetyltransferase activity is conserved between yeast and human GCN5 and is required for complementation of growth and transcriptional activation. *Mol. Cell. Biol.* 17: 519–527, 1997.
- Horlein, A. J., Naar, A. M., Heinzl, T., Torchia, J., Gloss, B., Kurokawa, R., Ryan, A., Kamei, Y., Soderstrom, M., and Glass, C. K. Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. *Nature (Lond.)*, 377: 397–404, 1995.
- Futamura, M., Nishimori, H., Shiratsuchi, T., Saji, S., Nakamura, Y., and Tokino, T. Molecular cloning, mapping, and characterization of a novel human gene. MTA1-L1, showing homology to a metastasis-associated gene, MTA1. *J. Hum. Genet.*, 44: 52–56, 1999.
- Barnes, D. M., and Gillett, C. E. Cyclin D1 in breast cancer. *Breast Cancer Res. Treat.*, 52: 1–15, 1998.
- Yamauchi, H., Stearns, V., and Hayes, D. F. When is a tumor marker ready for prime time? A case study of c-erbB-2 as a predictive factor in breast cancer. *J. Clin. Oncol.*, 19: 2334–2356, 2001.
- Planas-Silva, M. D., Shang, Y., Donaher, J. L., Brown, M., and Weinberg, R. A. AIB1 enhances estrogen-dependent induction of cyclin D1 expression. *Cancer Res.*, 61: 3858–3862, 2001.
- Newman, S. P., Bates, N. P., Vernimmen, D., Parker, M. G., and Hurst, H. C. Cofactor competition between the ligand-bound oestrogen receptor and an intron 1 enhancer leads to oestrogen repression of ERBB2 expression in breast cancer. *Oncogene*, 19: 490–497, 2000.
- Bloom, H. J. G., and Richardson, W. W. Histological grading and prognosis in breast cancer. *Br. J. Cancer*, 11: 359–377, 1957.
- Bieche, I., Onody, P., Laurendeau, I., Olivieri, M., Vidaud, D., Lidereau, R., and Vidaud, M. Real-time reverse transcription-PCR assay for future management of ERBB2-based clinical applications. *Clin. Chem.*, 45: 1148–1156, 1999.
- Bieche, I., Parfait, B., Laurendeau, I., Girault, I., Vidaud, M., and Lidereau, R. Quantification of estrogen receptor α and β expression in sporadic breast cancer. *Oncogene*, 20: 8109–8115, 2001.
- Bieche, I., Franc, B., Vidaud, D., Vidaud, M., and Lidereau, R. Analyses of MYC, ERBB2, and CCND1 genes in benign and malignant thyroid follicular cell tumors by real-time polymerase chain reaction. *Thyroid*, 11: 147–152, 2001.
- Chomczynski, P., and Sacchi, N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.*, 162: 156–159, 1987.
- Kaplan, E. L., and Meier, P. Nonparametric estimation from incomplete observations. *J. Am. Stat. Assoc.*, 53: 457–481, 1958.
- Cox, D. R. Regression models and life-tables. *J. R. Stat. Soc.*, 34: 187–220, 1972.
- Bieche, I., Onody, P., Lerebours, F., Tozlu, S., Hacene, K., Andrieu, C., Vidaud, M., Tubiana-Hulin, M., Spyrtos, F., and Lidereau, R. ERBB2 status and benefit from adjuvant tamoxifen in ER α -positive postmenopausal breast carcinoma. *Cancer Lett.*, 174: 173–178, 2001.
- Shang, Y., Hu, X., DiRenzo, J., Lazar, M. A., and Brown, M. Cofactor dynamics and sufficiency in estrogen receptor-regulated transcription. *Cell*, 103: 843–852, 2000.
- Bouras, T., Southey, M. C., and Venter, D. J. Overexpression of the steroid receptor coactivator AIB1 in breast cancer correlates with the absence of estrogen and progesterone receptors and positivity for p53 and HER2/neu. *Cancer Res.*, 61: 903–907, 2001.
- Kurebayashi, J., Otsuki, T., Kunisue, H., Tanaka, K., Yamamoto, S., and Sonoo, H. Expression levels of estrogen receptor- α , estrogen receptor- β , coactivators, and corepressors in breast cancer. *Clin. Cancer Res.*, 6: 512–518, 2000.
- Cottone, E., Orso, F., Biglia, N., Sismondi, P., and De Bortoli, M. Role of coactivators and corepressors in steroid and nuclear receptor signaling: potential markers of tumor growth and drug sensitivity. *Int. J. Biol. Markers*, 16: 151–166, 2001.
- Lavinsky, R. M., Jepsen, K., Heinzl, T., Torchia, J., Mullen, T. M., Schiff, R., Del-Rio, A., Ricote, L., Ngo, M., Gemsch, S., Hilsenbeck, J., Osborne, S. G., Glass, C. K., Rosenfeld, C. K., and Rose, D. W. Diverse signaling pathways modulate nuclear receptor recruitment of N-CoR

- and SMRT complexes. *Proc. Natl. Acad. Sci. USA*, *95*: 2920–2925, 1998.
34. Yamamoto, Y., Wada, O., Suzawa, M., Yogiashi, Y., Yano, T., Kato, S., and Yanagisawa, J. The tamoxifen-responsive estrogen receptor α mutant D351Y shows reduced tamoxifen-dependent interaction with corepressor complexes. *J. Biol. Chem.*, *276*: 42684–42691, 2001.
35. Zhang, X., Jeyakumar, M., Petukhov, S., and Bagchi, M. K. A nuclear receptor corepressor modulates transcriptional activity of antagonist-occupied steroid hormone receptor. *Mol. Endocrinol.*, *12*: 513–524, 1998.
36. Jackson, T. A., Richer, J. K., Bain, D. L., Takimoto, G. S., Tung, L., and Horwitz, K. B. The partial agonist activity of antagonist-occupied steroid receptors is controlled by a novel hinge domain-binding coactivator L7/SPA and the corepressors N-CoR or SMRT. *Mol. Endocrinol.*, *11*: 693–705, 1997.
37. Chan, C. M. W., Lykkesfeldt, A. E., Parker, M. G., and Dowsett, M. Expression of nuclear receptor interacting proteins TIF-1, SUG-1, receptor interacting protein 140, and corepressor SMRT in tamoxifen-resistant breast cancer. *Clin. Cancer Res.*, *5*: 3460–3467, 1999.
38. Kurokawa, H., Lenferink, A. E., Simpson, J. F., Pisacane, P. I., Sliwkowski, M. X., Forbes, J. T., and Arteaga, C. L. Inhibition of HER2/neu (erbB-2) and mitogen-activated protein kinases enhances tamoxifen action against HER2-overexpressing, tamoxifen-resistant breast cancer cells. *Cancer Res.*, *60*: 5887–5894, 2000.

Clinical Cancer Research

Expression Analysis of Estrogen Receptor α Coregulators in Breast Carcinoma: Evidence That NCOR1 Expression Is Predictive of the Response to Tamoxifen

Igor Girault, Florence Lerebours, Samira Amarir, et al.

Clin Cancer Res 2003;9:1259-1266.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/9/4/1259>

Cited articles This article cites 37 articles, 12 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/9/4/1259.full#ref-list-1>

Citing articles This article has been cited by 15 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/9/4/1259.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/9/4/1259>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.